***Lysionotus chatungii* – a new species of Gesneriaceae from Arunachal Pradesh in North-Eastern India**

Momang Taram, A. P. Das and Hui Tag¹

Plant Systematic and Ethnobotanical Research Laboratory, Department of Botany, Rajiv Gandhi
University, Rono Hills, Doimukh-791112, Arunachal Pradesh, India

¹*Communicating author; E-mail: huitag2008rgu@gmail.com*

[Received 01.12.2019; Revised ..12.2019; Accepted ..12.2019; Published 31.12.2019]

Abstract

Lysionotus chatungii M.Taram, A.P Das & H.Tag, a new species of Gesneriaceae is described from Geku in Upper Siang District of Arunachal Pradesh in North-Eastern India. The new species is presented here along with its detailed description, habitat structure and distribution along with a plate of color photographs for its easy recognition.

Key words: Gesneriaceae, *Lysionotus chatungii*, New species, Arunachal Pradesh.

INTRODUCTION

The genus *Lysionotus* D.Don (1822) is distributed from North India and Nepal eastwards through South China, Taiwan, Bhutan, Laos, Myanmar, North Thailand, North Vietnam to South Japan (Moller *et al.* 2017). According to Moller *et al.* (2017) 6 species of *Lysionotus* are found in India and 3 species of *Lysionotus* in North East India (Sinha & Dutta 2016). Giri *et al.* (2008) recorded 3 species (*L. atropurpurea* H.Hara, *L. palinensis* G.D. Pal and *L. serratus* D.Don). Recently two new taxa of *Lysionotus* viz. *L. gamosepalous* W.T Wang var. *biflorus* A.Joe *et al.* (2017) and *L. bijantiae* D.Borah & A.Joe (2018) have been described from Arunachal Pradesh. In this article another new species of *Lysionotus* is presented from Arunachal Pradesh in North Eastern India. The new species is described here and illustrated with a plate of color photographs. With this new species described herein, there are now five species and two varieties of *Lysionotus* recorded from Arunachal Pradesh as well as from Northeastern India.

Taxonomy

Lysionotus chatungii M.Taram, A.P. Das & H.Tag, *sp.nov.* [PLATE – I]

Lysionotus chatungii is morphologically similar to *Lysionotus chingii* Chun ex W.T. Wang but can be easily distinguished by a number of characters like lamina sub-coreaceous lamina (vs. papery/chartaceous); ovate-elliptic to rounded-ovate (vs. elliptic-oblong); base obliquely-cordate to rounded (vs. cuneate); calyx divided from base (vs. calyx divided up to the middle) (Table 1)

Type: INDIA, Arunachal Pradesh: Upper Siang District, Geku. 28° 27' 48.3" N and 95° 82' 10.3" E; Elevation 873 m; 15 April 2019, *Momang Taram 0301* [Holotype CAL; Isotypes ASSAM, ARUN]

PLATE - I. *Lysionotus chatungii* M.Taram, A.P. Das & H.Tag: **A.** Twig with alternate leaves; **B.** Twig with opposite leaves and 4-nate in prophyll bearing portion; **C.** Lamina with unequal and slightly cordate base; **D & E.** Prophylls; **F.** Flower in lateral view; **G.** Flower in front view; **H.** Emergence of peduncle from axiles of petiole and prophyll; **I.** Union of peduncle and pedicel with articulation showed by arrow (bract absent); **J.** Sepal lobes; **K.** Corolla throat inside with two calli; **L.** Two stamens with adherant anthers; **M.** Pistil with disc; **N.** Mature follicles; **O.** Seed.

Epiphytic, or on rocky-walls, root-climbing or creeping sub-shrub, 150 – 300 cm long, terete, much branched, glabrous, young stem green with purple irregular blotches, mature stem brown; internodes 2 – 12 cm. Leaves mostly opposite, but may be alternate, even ternate or in whorl of 4; sometimes two leaves of normal shape and size are alternating with prophylls; 4 or more prophylls occur at every node in lower non-leafy part of branches; Petioles 0.5 – 1 cm, glabrous, purple; Lamina ovate-elliptic to rounded ovate, 7 – 20 × 5 – 12 cm, nearly entire to minutely dentate to obscurely serrate, shortly caudate-acuminate, base rounded to shallowly cordate, generally oblique, lateral veins sub-opposite 4 – 8 on each side of midrib, adaxially dark green, abaxially whitish green with purple blotches, glabrous, sub-coreaceous. Inflorescence axillary, 2 – 4 in each node, 1-flowered (cymose-solitary) each; peduncle slender, 3.0 – 4.0 cm long, terete, green with purple irregular blotches, glabrous, attachment with pedicel prominently articulated; pedicel slender but slightly thicker, 2.0 – 2.5 cm long, terete, purple, glabrous. Peduncles not arising from leaf-axile are with a ovate to oblong prophyll (0.4 – 1.0 cm x 0.1 – 0.3 cm) at base. Calyx divided into 5 lobes from base; each lobe slender-linear to linear-subulate, anterior 3 longer [1 - 2 cm x 0.1 - 0.2 cm], posterior 2 shorter [0.7 – 1.3 cm x 0.1 – 0.2 cm], or sometimes all equal, entire, narrowly acuminate, glabrous, purple. Corolla bilabiate, 3 + 2, anterior lip deeply 3-lobed, 0.6 – 1.2 cm long, sub-equal, ovate, more or less entire, rounded; posterior lip obscurely 2-lobed, 0.4 – 0.8 cm long, glabrous, white-purplish with deep-purple reticulation, tube 1.6 – 2.3 cm long, slightly inflated upwards, 4 – 4.3 cm long, glabrous, purplish with deep-purple stripes and 2 yellow elongated calli at throat alternating with anterior lobes. Stamens 2, included, filaments 0.5 – 0.7 cm long, attached to corolla tube along the two margins of the anterior lip, terete, yellow, glabrous, curved and blackish at the tip, connected to the middle of anther; anthers broadly oblong (oblate), cohering each-other face-to-face, white,

Table 1. Comparison of *Lysionotus atropurpureus* and *Lysionotus chingii* with the new species *Lysionotus chatungii*

Characters	<i>Lysionotus atropurpureus</i> Hara	<i>Lysionotus chingii</i> Chun ex W.T. Wang	<i>Lysionotus chatungii</i> sp. nov.
Habit	Epiphyte	Climbing sub-shrubs	Climbing sub-shrubs
Stem length	Up to 25 cm	Up to 9 m	Up to 3 m
Petiole	0.5 – 1.5 cm	Longer, up to 2.3 cm	Up to 1.0 cm
Lamina shape	Oblong lanceolate to narrowly elliptic	Elliptic to oblong	Ovate-elliptic to rounded-ovate
texture	Papery	Papery	Sub-coriaceous
margin	Denticulate	Entire to denticulate with small glands	Entire to minutely dentate to sub-serrate without glands
base	Cuneate	Broadly cuneate to cuneate	Rounded to obliquely and shallowly cordate
apex	Acuminate	Acuminate	Shortly caudate-acuminate
Inflorescence	Cymes 1 – 4 flowered	Solitary	Solitary
Bracts	Linear	Orbicular- ovate	Ebracteate
Peduncle	5.5 – 7.5 cm	1.4 – 2.8 cm	2.7 – 4.0 cm
Pedicel	0.7 – 2 cm	0.2 – 0.7 cm	1.4 – 2.5 cm
Calyx	Divided from base	Divided above middle; lobes equal; tube 1.2 – 1.8 cm	Divided from base; posterior 2 lobes shorter; tube absent
Corolla	Purple	Tinged greenish; 3.3 cm long	White-purplish with deep-purple reticulation; tube 1.6 – 2.3 cm long
Disc	Ring-like with entire margin	Ring-like with entire margin	4-lobed, lobes oblong with conical tips
Flowers & fruits	July – October	July – September	March – August

dehiscing longitudinally, unappendaged; staminodes 2, 0.1 – 0.3 cm long, linear, yellow. Ovary 2 cm long, shortly stipitate, cylindrical, greenish, glabrous; style 0.5 cm long, terete, glabrous; stigma slightly bilamellate. Nectary 4-lobed, lobes erect, oblong with conical tip. Capsules 10 – 12 cm long, linear, glabrous, dehiscing longitudinally; Seeds 0.3 – 0.4 cm long, linear, with along slender funiculus at base and long brownish hair-like appendage at the tip.

Affinities: The habit and solitary inflorescence of new species *Lysionotus chatungii* is similar with *Lysionotus chingii*. However, the new specie is distinguished from it by much shorter stem length; sub-coriaceous leaves; ovate-elliptic to rounded-ovate lamina with rounded to obliquely-cordate base; calyx segments divided to the base.

Another somewhat similar looking species, *Lysionotus atropureus* Hara is also available in North-east India including Arunachal Pradesh (Giri *et al.* 2008). However, structurally/ morphologically it is a completely different plant as in numerous important characters like habit, leaf structure, inflorescence, flower structure etc. the new species is completely unique.

Table 1 shows the differences of the new species, *Lysionotus chatungii* with that of *L. chingii* and *L. atropureus*.

Flowering and Fruiting: Flowering start from March and fruiting ends in August.

Etymology: This species is named after Late **Taro Chatung**, the first and the most famous journalist of Arunachal Pradesh whose contribution in journalism for this remote state is immense.

Distribution, Threat-status and Ecology: So far, the new species known only from its Type-locality in the Upper Siang district of Arunachal Pradesh in Northeast India and can be treated as endemic. The plant was found growing on rocks and on tree-trunks in moist shaded areas along the margin of a narrow but perennial stream. The estimated area of its occurrence is less than 3 km² and the population is thought to be less than 100 individual. It was growing in association with Mosses, *Asplenium* sp., some epiphytic ferns, *Aeschynanthus parviflorus* (D. Don) Spreng., *Codonanthe devosiana* Lem., etc.

Acknowledgements

The authors are thankful to Dr. Sudeshna Datta of Botanical Survey of India for her useful comments on the specimen and to Mr. Dipankar Borah for his help in the laboratory. They are also thankful to Mr. Ojar Taku, Mr. Bosong Taram, Olom Tosung and Mr. Gemin Taku for their logistic supports during the field works.

LITERATURE CITED

- Borah, D. & Joe, A. 2018. A new species of *Lysionotus* (Gesneriaceae) from Northeastern India. *Taiwania* 63(3): 232 – 234.
- Don, D. 1822. Description of two new genus of Nepal plants. *Edinburgh Phil. J.* 7: 82 – 86.
- Giri, G.S.; Pramanik, A. & Chowdhery, H.J. 2008. *Materials for the Flora of Arunachal Pradesh*. Vol. II. Botanical Survey of India, Kolkata.
- Joe, A.; Hareesh, V.S. & Sabu, M. 2017. A new taxon of *Lysionotus* (Gesneriaceae) from Northeastern India. *Taiwania* 62(4): 337 – 339.
- Moller, M.; Nampy, S.; Janeesha, A.P. & Weber, A. 2017. The Gesneriaceae of India: Consequences of updated generic concepts and new family classification. *Rheedea* 27(1): 23 – 41.
- Sinha, B.K. & Datta, S. 2016. Taxonomic account on the family Gesneriaceae in Northeast India. *Nelumbo* 58: 1 – 43