CLINICAL HEALTH PROMOTION

Research & Best Practice for patients, staff and community

PhD Thesis by Mette Rasmussen

Research and Best Practice

Intensive Smoking Cessation Interventions in Denmark

- Based on data from the Danish Smoking Cessation Database

Editorial Office

WHO-CC, Clinical Health Promotion Centre
Health Sciences, Lund University,
Sweden

The Official Journal of

The International Network of Health Promoting
Hospitals and Health Services

CLINICAL HEALTH PROMOTION

FACULTY OF HEALTH AND
MEDICAL SCIENCES
COPENHAGEN UNIVERSITY

PhD thesis by Mette Rasmussen, 2018.

This thesis has been submitted to the Graduate School of The Faculty of Health and Medical Sciences, Clinical Research, University of Copenhagen.

Thesis submitted July 15th 2018.

WHO-CC

Bispebjerg/Frederiksberg University Hospital
Nordre Fasanvej 57
Hovedvejen, indg. 5, 2. sal
DK-2000 Frederiksberg
Mette.Rasmussen.03@regionh.dk

Supervisors

Professor Vibeke Backer, MD DMSci
University of Copenhagen
and Respiratory Research Unit,
Department of Respiratory Medicine, Copenhagen University Hospital
Bispebjerg & Frederiksberg, Denmark

Professor Hanne Tønnesen, MD DMSci
Clinical Health Promotion Centre, Copenhagen University Hospital
Bispebjerg & Frederiksberg, Denmark
and Department of Health Sciences
Lund University, Skåne University Hospital, Sweden

Head of Committee

Professor Charlotta Holm Pisinger
University of Copenhagen

Examiners

Professor Ole Hilberg
University of Southern Denmark

Professor Erik Börje Bjelke
University of Oslo

Clinical Health Promotion is an open access journal and all issues can be downloaded free of charge at www.clinhp.org

Copying and other reproduction of material is permitted with proper source reference.

© WHO-CC, Clinical Health Promotion Centre, Bispebjerg & Frederiksberg Hospital, University of Copenhagen, Denmark & Health Sciences, Lund University, Sweden 2018.

ISSN 2226-5864

PhD Thesis

Mette Rasmussen, MSc

Intensive Smoking Cessation Interventions in Denmark

Based on data from the Danish Smoking Cessation Database

This thesis has been submitted to the Graduate School of the Faculty of Health and Medical Sciences,
University of Copenhagen on 15 July 2018

Included studies

- I. Mette Rasmussen, Bolette Pedersen, Susanne Vahr Lauridsen, Vibeke Backer, Hanne Tønnesen (Manuscript)
Efficacy of intensive face-to-face interventions for smoking cessation – a systematic review and meta-analysis.
- II. Mette Rasmussen, Hanne Tønnesen (Published)
The Danish Smoking Cessation Database
Clin Health Promot 2016; 6:36-41
doi: 10.29102/clinhp.16006
- III. Mette Rasmussen, Esteve Fernández, Hanne Tønnesen (Published)
Effectiveness of the Gold Standard Programme compared with other smoking cessation interventions in Denmark: a cohort study
BMJ Open Feb 2017, 7 (2) e013553
doi: 10.1136/bmjopen-2016-013553
- IV. Mette Rasmussen, Mads Klinge, Jesper Krogh, Merete Nordentoft, Hanne Tønnesen (Published)
Effectiveness of the Gold Standard Programme (GSP) for Smoking Cessation on smokers with and without a severe mental disorder—a Danish cohort study
BMJ Open Jun 2018, 8 (6) e021114
doi: 10.1136/bmjopen-2017-021114

Acknowledgement

This thesis was made possible by the many patients who agreed to let me use their data and experiences, by all staff at the participating smoking cessation units, and the coordinating office. I owe you all special thanks.

Likewise, I want to thank my supervisor, Hanne Tønnesen (principal investigator), WHO-CC Clinical Health Promotion Centre, for academic guidance, critical and supportive feedback and inspiring scientific discussions throughout the process. Also, thanks to my supervisor Vibeke Backer, for all your insight and helpful advice.

I want to thank Esteve Fernández, WHO-CC for Tobacco Control in Barcelona, and his entire crew, who all generously welcomed me in the department for two months and made me feel at home from the very first day. Esteve, your support and involvement in my research is much appreciated. Also, thanks to Antoni Gual, Hospital Clínic de Barcelona, for introducing me to the process of implementation of smoke-free psychiatry.

Thanks to the co-authors of the included papers in this thesis, who have provided great time and effort: Vibeke Backer, Esteve Fernández, Mads Klinge, Jesper Krogh, Susanne Vahr Lauridsen, Merete Nordentoft, Bolette Pedersen and Hanne Tønnesen.

These years would not have been the same without my fellow PhD students Anne-Birgitte Hjuler Ammari, Julie Weber Melchior Egholm, Helle Juul-Larsen, Amélié Cléo Keller, Susanne Vahr Lauridsen, Bolette Pedersen, and Jeff Kirk Svane. I have enjoyed every single scientific and technical discussions throughout the years. Your curiosity and interest has been invaluable and much appreciated, as has your friendships. I look forward to working together in the future.

I also owe thanks to my great colleagues at the WHO-CC Clinical Health Promotion Centre, in Denmark and Sweden, for their continued support, help, interest and input.

Last, but not least, I want to thank my encouraging, helpful and accepting family and friends.

Mette Rasmussen
Copenhagen, Oktober 2018

Contents

Included studies	3
Acknowledgement.....	4
Contents.....	5
Definitions and abbreviations.....	6
English summary	8
Danish summary / Dansk resumé	9
Introduction	10
Background	11
The smoking epidemic	12
Smoking cessation interventions	15
Smoking cessation interventions in Denmark.....	17
Efficacy and effectiveness research	20
Objective and hypothesis	21
Materials and methods.....	22
Trial profile for the thesis	22
Methods of the included studies	23
Results and discussion	24
Summary of findings	24
Discussion	28
Bias and limitations.....	31
Ethical considerations.....	32
Conclusion and impact.....	33
Perspectives.....	33
References	35
Appendix/Papers.....	43
Manuscript I.....	44
Paper II	62
Paper III	69
Paper IV	80

Definitions and abbreviations

Throughout this thesis, the following definitions and abbreviations will be used.

Definitions	
Intensive smoking cessation intervention	Intensive smoking cessation interventions (ISCIs) are defined in accordance with the <i>Clinical Practice Guideline for Treating Tobacco Use and Dependence: 2008 Update</i> ¹ . These interventions include at least four scheduled in-person meetings, with each session lasting >10 minutes. The content of the interventions includes an element of patient education that covers basic information, danger situations, and coping skills. The ISCI can be either individually based or group based, and no restrictions on the use of pharmacotherapy are imposed. ^{Manuscript I}
Shorter interventions	Shorter interventions (SI) are defined as ‘interventions delivered in-person and/or by telephone, including a maximum of 3 in-person contacts and/or no more than 1 hour of total contact time’. ^{Manuscript I}
Continuous abstinence	‘ <i>Continuous abstinence</i> refers to abstinence between quit day and a follow-up time’. ²
Point prevalence abstinence	‘ <i>Point prevalence</i> abstinence refers to the prevalence of abstinence during a time window immediately preceding follow-up’. ²
Real life	These are interventions that are already implemented into a normal daily routine in a smoking cessation clinic.

Abbreviations	
CA	Continuous Abstinence
COPD	Chronic Obstructive Pulmonary Disease
CPR	Central Personal Register (social security number)
EBM	Evidence-Based Medicine
FCTC	Framework Convention for Tobacco Control
GSP	Gold Standard Programme
ICD-10	International Classification of Diseases, 10 th edition
ISCI	Intensive Smoking Cessation Intervention
NCD	NonCommunicable Diseases
PP	Point Prevalence
RCT	Randomised Clinical Trial
SC	Smoking Cessation
SCDB	The National Danish Smoking Cessation Database
SCI	Smoking Cessation Interventions
SI	Shorter Intervention
SMD	Severe Mental Disorder
SUD	Substance Use Disorder
WHO	World Health Organization

English summary

Background

Tobacco smoking is by far the most deadly and preventable lifestyle factor for public health, both globally and nationally. Nevertheless, it is estimated that worldwide, more than 1.1 billion adults are smokers. Today, 16% (0.8 million) of the adult population in Denmark are daily smokers, while 23% (1.1 million) consider themselves smokers, and one in four deaths are smoking related. The most effective method of reducing smoking prevalence is by employing a two-tier strategy to 1) prevent adolescents from initiating smoking and 2) offer effective help to smokers to quit.

The aim of this thesis was to evaluate the effect of intensive smoking cessation interventions (ISCI) on successful quitting through different study designs, synthesised from cohort studies and a review of randomised controlled trials. The aim was fulfilled through 4 scientific studies, each with their own objective.

Studies

A systematic review with meta-analysis (I) was set up to compare ISCI and shorter interventions in randomised controlled trials (RCT). Initially, 9,569 unique papers were identified, and, after screening for relevant studies, 18 RCTs were included, investigating a total of 10,131 smokers. The studies were primarily conducted in Western Europe and the USA. The body of evidence for three of four outcomes were graded moderate, and the last outcome was graded low. A random effect meta-analysis revealed that smokers randomised to ISCI were significantly more likely to be successful quitters compared to shorter interventions for all four outcomes. The primary outcome (continuous abstinence in the short and long term) increased the chance of success 3-fold.

A methodological study (II) of the Danish National Smoking Cessation Database (SCDB) was undertaken. The development of the SCDB and the most commonly registered smoking cessation interventions (SCI) were described, as were the data collection and validity.

Two cohort studies based on the data from the SCDB were conducted to evaluate the effectiveness of SCI in real life. One study investigated the effectiveness of five different SCIs (III) and found that the intensive Gold Standard Programme' (GSP - Standard SCI in Denmark) was the only intervention that was effective for both men and women. The other cohort study was set up to investigate the effectiveness of the GSP in a vulnerable subgroup of smokers diagnosed with severe mental disorder (SMD) (IV). The study showed that smokers with SMD were significantly less likely to become successful quitters compared to smokers without mental disorders. However, one in every four smokers with SMD managed to stay continuously smoke-free for at least 6 months compared to one in three in the comparison group. The data were analysed using a mixed-effect logistic regression model. In both cohort studies, compliance was identified as the strongest predictor of a successful outcome.

Conclusion

This thesis adds to the evidence on intensive smoking cessation interventions (ISCI) through different study designs. The results from both study designs favoured ISCI, although the GSP was significantly less effective in smokers with SMD compared to smokers without mental illness. To develop effective SCI in the future, it is important to evaluate the effects in an RCT and to follow-up the post-implementation effects in real life (e.g., through a national SCDB).

Danish summary / Dansk resumé

Baggrund

Rygning er ubetinget den mest dødelige men forebyggelige livsstilsfaktor for folkesundheden både globalt og nationalt. Dette til trods anslås det, at der findes mere end 1,1 milliarder voksne rygere på verdensplan. I Danmark ryger 16 % (0,8 millioner), af den voksne befolkning dagligt, og 23 % (1,1 millioner) betegner sig selv som rygere. Hvert fjerde dødsfald i Danmark kan tilskrives rygning. Den mest effektive måde at reducere antallet af rygere er ved at anvende en to-strengs strategi med de formål at 1) forebygge at unge begynder at ryge, og 2) tilbyde effektiv hjælp til rygestop.

Formålet med denne afhandling var at evaluere effekten af intensive rygestopprogrammer målt ved succesfuldt rygestop. Dette blev undersøgt gennem forskellige studie-design i form af kohortestudier og en systematisk litteraturgennemgang af randomiserede studier. Formålet blev opfyldt via 4 videnskabelige studier med hver deres forskningsspørgsmål.

Studier

I det systematiske review (I) blev intensive rygestopprogrammer sammenlignet med kortere programmer i randomiserede forsøg. I litteratursøgningen blev 9.569 artikler identificeret, og efter en grundig gennemgang blev 18 randomiserede studier, med i alt 10.131 rygere, inkluderet. Studierne var hovedsagligt gennemført i Vesteuropa og USA. Kvaliteten af evidensen blev vurderet som moderat for tre af fire effektmål og lav for ét. Meta-analyser viste at intensive rygestopprogrammer var signifikant mere effektive end kortere programmer for alle fire effektmål. Intensive rygestopprogrammer var 3 gange så effektive som kortere programmer, målt som kontinuert rygestop på kort og lang sigt (primære effektmål).

Der blev gennemført et metodestudie (II) af den nationale danske rygestopbase. Her blev udviklingen af Rygestopbasen, de mest almindelige rygestopprogrammer, procedurer for dataindsamling samt validiteten af databasen beskrevet.

Baseret på data fra Rygestopbasen blev der udført to kohorte studier for at evaluere effekten af rygestopprogrammer implementeret i den kliniske hverdag. Ét studie undersøgte effekten af fem forskellige rygestopprogrammer (III), og viste at det intensive danske 'guld standard program' (GSP) var det eneste program med en signifikant effekt for både mænd og kvinder. Det andet kohorte studie undersøgte effekten af GSP i en sårbar gruppe af rygere diagnosticeret med svær mental sygdom. Studiet viste at rygere med svær mental sygdom havde signifikant lavere effekt af GSP end rygere uden mental sygdom. På trods af den lavere effekt, var hver fjerde ryger med svær mental sygdom stadig kontinuert røgfri efter 6 måneder, sammenlignet med hver tredje ryger uden mental sygdom. Kohorte studierne blev analyseret med en logistisk regressionsmodel. I begge studier blev gennemførelse af rygestopprogrammet identificeret som den vigtigste prædiktør for et succesfuldt rygestop.

Konklusion

Denne afhandling bibringer ny viden om intensive rygestopprogrammer gennem forskellige studie-design. Resultaterne fra begge anvendte design peger på intensive rygestopprogrammer som de mest effektfulde, på trods af en signifikant lavere effekt blandt rygere med svær mental sygdom. For i fremtiden at kunne udvikle effektive rygestopprogrammer er det vigtigt at evaluere effekten af programmerne i randomiserede forsøg, for efterfølgende at monitorere effekten efter de er blevet implementeret i den kliniske hverdag, for eksempel via nationale effekt-databaser som den danske rygestopbase.

Introduction

Smoking is still the number one single and preventable factor of greatest importance to the health of the population nationally and globally, with more than 1.1 billion smokers worldwide ³. In 2015, smoking caused more than 7.2 million deaths globally ⁴, corresponding to one in every ten deaths ⁵. In Denmark, the smoking prevalence has dropped by approximately 1% per year from 2003-2012; since then, the prevalence has stagnated at approximately 16%, corresponding to 0.8 million daily smokers ⁶, and 13,600 smoking related deaths occur every year ⁷.

A reduction of the smoking prevalence is best achieved through two means: 1) preventing adolescents from starting smoking and 2) offering smokers help with quitting; the latter method is the primary focus of this thesis.

While the damages from tobacco smoking ⁸⁻¹¹, the tobacco epidemic ^{3,12,13} and the benefits after smoking cessation ^{14,15} are well known and well described, knowledge of how to achieve successful quitting is more sparse. It is clear that successful quitting is influenced by multiple factors. However, these factors are not all directly related to the included studies. Some of these factors, such as tobacco control policies, nicotine addiction, vulnerable groups, and the composition of smoking cessation interventions, will be covered in the background, as they can all potentially influence the motivation for and the outcomes of a quitting attempt in some manner. Finally, the use of quality assessment tools to monitor and follow-up the effects of evidence-based interventions implemented into a daily routine (i.e., the 'effectiveness in real life') are introduced, along with the concepts of effectiveness and efficacy research.

This thesis investigates the impact of intensive smoking cessation interventions through four scientific studies, which are referred to using roman numerals. The investigation is performed by systematically reviewing intensive smoking cessation interventions based on previously published literature (I). Since 2001, data on smoking cessation interventions implemented in the daily routine in Denmark have been collected through the National Smoking Cessation Database, which is described in a methodological study (II). This database offers a unique opportunity to explore the effects of smoking cessation interventions after implementation in a normal daily routine (i.e., 'effectiveness in real life'). The thesis also includes two cohort studies: one evaluating the effectiveness of different smoking cessation interventions (III) and the other specifically evaluating the effectiveness of an intensive smoking cessation intervention in smokers diagnosed with a severe mental disorder (IV).

The overall aim of this thesis was to evaluate the effect of intensive smoking cessation interventions (ISCI) on successful quitting through different study designs synthesised from cohort studies, as well as reviewing randomised controlled trials.

Background

Globally, the prevalence of smoking is approximately 18% (21.0% among men and 6.2% among women), corresponding to more than 1.1 billion smokers worldwide ³. In Denmark, the smoking prevalence is similar among men and women; 16.1% of men and 15.6% of women smoke on a daily basis, and a total of 23% considers themselves to be smokers ¹⁶. In absolute numbers, these percentages corresponds to 0.8 million daily smokers and 1.1 million smokers in the Danish population aged 15 years or older ⁶.

Despite the established evidence on the harmful effects of tobacco smoking ^{8–11} (figure 1), smoking continues to be the single most damaging and preventable public health lifestyle factor, globally and nationally ^{17–19}. It is well known that smoking causes the most common noncommunicable diseases (NCD), such as cardiovascular diseases, cancers and other respiratory diseases, including chronic obstructive pulmonary disease (COPD) ²⁰. In addition, smoking deteriorates a number of other conditions. Pregnancy is a clear example of this deterioration, as smoking causes a tremendous range of complications for both the mother and child ^{21–30}. Other clear examples include asthma patients ^{31–34} and surgical patients, for whom smoking doubles the risk of post-operative complications ^{35,36}.

Figure 1: Damage from tobacco smoking. COPD: Chronic obstructive pulmonary disease. Based on ^{15,37} and with the courtesy of 'Klinisk Enhed for Sygdomsforebyggelse' 2003.

Approximately half of all lifetime smokers die from smoking-related causes ^{17,38}. In addition, the longevity of lifetime smokers is 7-10 years shorter than that of never smokers ^{38,39}, and smokers have poorer self-rated health ⁴⁰. Every year, 13,600 people in Denmark die as a consequence of smoking ⁷, corresponding to approximately one in every four deaths ⁴¹, and these numbers have not changed much over the past decade ²⁰. Worldwide, in 2015, smoking caused more than 7.2 million deaths annually ⁴, corresponding to one in every ten deaths ⁵.

Globally, most smokers start smoking before the age of 20 ^{3,42}. Therefore, it is important that major efforts are directed towards prevention among young people, for example, through bans and taxation ⁴². In theory, these programmes are easily implemented; however, they heavily depend on political will. Thus, an increase in the proportion of young smokers is currently being observed in Denmark ^{41,43}. The proportion of daily smokers among those 16-24 years of age has increased to 15.6%, from 14.0% in 2013 ⁴¹. In comparison, the current prevalence of young smokers in Norway is just 3% compared to 23% 12 years ago, highlighting the importance of effective preventive strategies ⁴². Overall, there is great potential in reducing smoking and related illnesses.

The smoking epidemic

Despite differences among countries, the pattern of the smoking epidemic and the challenges related to fighting it are ubiquitous ³. In 1994, Lopez et al. described the basic elements of the smoking epidemic based on historical data from developed countries. The epidemic was characterised by 4 stages ¹² (table 1). The model is generic, prevalence may vary between countries, and local exceptions may exist.

	Stage I	Stage II	Stage III	Stage IV
Timeframe	1-2 decades	2-3 decades	3 decades	>4 decades
♂ prevalence	<15% <i>Towards the end</i> prevalence rises rapidly	Rapid rise to a peak of 50-80%	Begins to decline to ≈40% at the end of this stage. Lower among middle- aged and older adults	Declines slowly 2 decades in: ≈33-35% 4 decades in: <20%
♀ prevalence	Rarely exceeds 10% Could be <5%	Rapid increase, delayed by 1-2 decades compared to men	<i>Towards the end</i> , an initial decline in pre- valence following a long plateau of ≈35-45%. Higher prevalence among young women, and <10% among 55-60 year olds	Declines slowly 2 decades in: ≈30% 4 decades in: <20%
Typical smoker	Educated groups	Similar among socio- economic groups	Few educated	Mostly lower social- economic groups
Death due to smoking	Not yet evident with few †	♂†: Rising to 10% at the end of the phase ♀†: Few	†: Rapid rise ♂: 25-30% ♀: 5% <i>Towards the end</i> ♂† rates will have peaked	♂†: Peaks early in the period at 30-35% of all deaths, and 40-45% among middle aged adults. After a decade † decreases to <30% and progressively declines from here. ♀†: Rapid rise, until it peaks at 20-25% of all deaths 2-3 decades into this stage. Hereafter, it progressively declines.
Public opinion	Smoking becomes socially acceptable	Few ex-smokers	Knowledge about risks is well-spread	Demand smoke-free work-places and smoke- free personal environments.
Tobacco control strategies	Underdeveloped	Not well developed Risk of smoking not widely understood	Implementation of comprehensive tobacco control law. Smoke-free public places and transport. Smoke-free work-places are not yet common	Legislation regarding smoke-free work-places and policies to address nicotine-dependent smokers wanting to quit.

Table 1: The smoking epidemic described in 4 stages. Inspired by Lopez ^{12,13}.

A key feature of the smoking epidemic is the 30-40 year delay from onset of smoking to smoking-related mortality ^{12,44}. In addition, there is a marked difference between men and women; due to the significantly lower cumulative exposure to tobacco smoking in women, the mortality rate is correspondingly lower and appears approximately 2 decades later.

Like most developed countries, Denmark is well into the fourth stage of the tobacco epidemic (table 1), with both men and women having reached the peak of smoking-related deaths and a slow decrease of smoking prevalence ^{13,45}. Three decades ago, it was predicted that the current proportion of smoking related mortality among men and women would be approximately 33% and 20%, respectively ¹², based on the smoking prevalence in the Danish population in the 1980s, when 40% of women and 60% of men smoked ⁴⁶. This finding corresponds to the actual numbers ^{7,47}. These are disproportionally high death rates, which can be reduced by an increased focus on prevention and smoking cessation treatment.

International tobacco control

The nature of tobacco control is to reduce the prevalence of smokers. This reduction is best achieved through a two-tier strategy. First, it is necessary to prevent people from starting to smoke; however, given the high prevalence of people who already smoke, it is also important to provide adequate support to those willing to quit.

In response to the global tobacco epidemic and to reduce the related morbidity and mortality, the World Health Organization Framework Convention on Tobacco Control (WHO FCTC) ⁴⁸ was put into action in 2005 ^{49,50} after many years in the making. In fact, the need for an international strategy was voiced as early as 1979, and, 10 years later, the work was actually initiated ⁴⁹. In 1993, the notion of an international convention was proposed, although initially criticised by WHO for being overly ambitious ⁵¹. It was not until Gro Harlem Brundtland took office as the Director General of WHO in 1998, declaring tobacco control one of her two major priorities, that concrete negotiations began. Five years later and after considerable efforts and obstruction attempts by the tobacco industry, the convention was ready. It built on an alliance between the WHO, World Bank, United Nations International Children's Emergency Fund (UNICEF), researchers and other relevant partners ^{49,51}. Two years after opening for signatures by member states, the WHO FCTC was put into action, with 168 of the 192 member states having signed. This implementation made the WHO FCTC the fastest growing and most wide-ranging convention ever within the United Nations ^{49,51,52}. Today, 181 countries have adopted the convention ⁵².

Shortly following the WHO FCTC, the MPOWER measures, under the Bloomberg initiative ¹¹, were released in 2008 ^{3,37,49} (figure 2), as a practical tool to further reduce tobacco use.

Figure 2: The WHO MPOWER measures.

Even before these initiatives, many countries, including Denmark, had already introduced restrictive laws and strategies to reduce smoking ⁹. Since the 1990s, several initiatives to help smokers quit have been launched in Denmark, with smoking cessation programmes being just one among many initiatives. Additionally, policy measures such as higher tobacco taxes, and restrictions on sale and advertising have been implemented.

Today, there are an array of programmes helping smokers to quit, such as a national quit line, web support, campaigns, training materials, etc. Furthermore, Denmark signed the WHO global action plan for the prevention and control of NCDs 2013-2020, which includes the reduction of smoking by 30% in 2025 ⁵³.

Despite all these tobacco control initiatives, however, Denmark is still far away from full implementation of the MPOWER measures with the 'Protect people from tobacco smoke' at the lowest level of implementation, and only 3 in 6 policies at the highest level (figure 3), with 'Offer help to quit tobacco use' being one of them ⁵.

Figure 3: The Danish fulfilment in 2016 of the WHO FCTC reported by the MPOWER measures, based on the results reported by WHO ⁵.

A recent report on the 'Capacity Assessment on the Implementation of Effective Tobacco Control Policies in Denmark' has further elaborated on the inadequacy of the current level of Danish tobacco control ⁴³. Another report from 2015, using the Tobacco Control Scale ⁵⁴, placed Denmark at number 23 among 35 European countries, 8 places down from the 2013 ranking ⁵⁵.

Even though the development of international tobacco politics has been long and winding thus far, new and more ambitious initiatives are still put into action, the latest being the Tobacco Endgame, which aims to completely phase out smoking or any other use of tobacco ⁵⁶. For practical reasons, the involved countries (e.g., Finland, Scotland, Ireland and New Zealand) have decided on a smoking frequency of a few percentages as their goal ⁵⁷.

The abovementioned policies and actions have all contributed to a general change in the smoking culture and tradition, as well as a decrease in the prevalence by approximately 1% annually in Denmark from 2003-2012 (figure 4). From 2012 onwards, the prevalence has stagnated at approximately 16% daily smokers ⁵⁸.

To further reduce the smoking prevalence, it is necessary to institute more restrictive smoke-free policies and improve other tobacco control elements, such as in Norway, where only 11% of the population 15 years or older are daily smokers, decreasing from 16% in 2012 and 31% in 2000 ⁵⁹.

In contrast to the inadequate tobacco control policies, Denmark has developed a national standardised intensive smoking cessation intervention (ISCI) ^{1,60-63}, internationally described as the Gold Standard Programme (GSP), in addition to an open quit line ⁶⁴. The GSP is freely accessible without referral in the municipalities.

Figure 4: Smoking prevalence in Denmark 2003-2017 ⁵⁸. The drop in 2009 coincided with a change in the execution of the survey.

While the monitoring of the smoking epidemic is well implemented, the knowledge and monitoring of the effect of smoking cessation intervention (SCI) is more lacking.

Smoking cessation interventions

Intensive smoking cessation interventions

In 2000, a practice guideline was published in the USA ⁶⁵. This work was based on a comprehensive literature review, including more than 50 meta-analyses, within the field of smoking cessation. This guideline proposed a definition of ISCI, based on a number of single factors that were associated with successful quitting. The guideline was updated in 2008, with minor changes in the ISCI definition ^{1,66}. According to this international guideline, an SCI can be regarded an ISCI if it contains at least 4 in-person sessions, each lasting >10 minutes, and if it includes an element of patient education (as a minimum covering recognition of danger situations, coping skills and basic information). ISCI can be both individual or group-based, and it may be with or without elements of pharmacotherapy ¹.

Shorter interventions

An inspection of multiple Cochrane reviews ⁶⁷⁻⁷² on smoking cessation clarifies that there is no single definition of short interventions. Short interventions can, however, be divided into three categories: dialogues (e.g., motivational interviewing or brief intervention), monologues (e.g., advice or very brief advice [VBA]) or 'no personal contact' (self-help materials, e.g., written, web-based, apps, or videos). Based on the non-existence of a common definition throughout this thesis, shorter interventions (SI) are defined as 'interventions delivered in-person and/or by telephone, including a maximum of 3 in-person contacts and/or no more than 1 hour of total contact time', thus including both dialogues and monologues possibly in combination with self-help materials. This definition is also used by the WHO in the international standards for health promotion and health care ⁷³.

Nicotine addiction

More than 9,500 organic and inorganic components have been identified in cigarette smoke ⁷⁴, at least 70 of which can cause cancer ⁷⁵. Nicotine is likely the best-known component in tobacco smoke. Though not carcinogenic, nicotine is responsible for the physical addiction to smoking

^{76,77}. Nicotine addiction is one of the reasons why successful quitting is hard to achieve and is the main reason why tobacco smoking occurs worldwide ^{76,78}. Nicotine addiction can be described as taking place in 3 phases; ‘acquisition and maintenance of nicotine-taking behaviour, withdrawal symptoms upon cessation of nicotine intake and vulnerability to relapse’ ⁷⁸.

The biochemical mechanisms of nicotine addiction are complex and involve different subtypes of nicotine-receptors in the brain, however, in line with other addictions, it is basically related to the reward systems of the brain, such as dopamine release ⁷⁸. After having smoked regularly for a period of time, the body adapts to an increased level of nicotine, and hence, starts craving nicotine to avoid withdrawal symptoms, such as irritability, anxiety, and depressive mood ⁷⁸. When lighting up a cigarette, the smoker immediately reports the effect as being relaxing, with increased concentration, and stress reduction ⁷⁶. It is likely that these nicotine effects are actually the result of relief from addiction symptoms rather than from the nicotine itself ⁷⁶. After smoking cessation, the nicotine craving can remain for weeks or years, which makes the smoker vulnerable to relapse ⁷⁸.

In addition to the direct nicotine addiction, smoking is also associated with social and cultural factors, all adding to the difficulty of quitting ⁷⁸. Whether someone starts smoking, attempts to quit, or continues to smoke is also related to the smoking culture in their surroundings ⁷⁶. Furthermore, smoking is often coupled to behavioural routines, such as drinking a cup of coffee, taking a break, ending a meal, or meeting with certain friends. By repeating these routines, they become associated with smoking and trigger the urge to smoke ⁷⁶.

The Fagerström test for nicotine dependence (FTND) was developed to score addiction level using a 10-point score. In clinical routine, it is used to guide nicotine replacement therapy during smoking cessation interventions. A value lower than 2 points seldom indicates a need for medicine, while a value higher than 6 points frequently indicates the need for treatment, most effective if combining slow and fast released nicotine products (e.g., patches and spray) ⁷⁹.

Benefits of smoking cessation

However, difficult as it may seem, there are benefits related to smoking cessation, both in the short term and in the long term. Figure 5 lists the improvements after smoking cessation.

Figure 5: Improvements after smoking cessation. Based on ^{15,80}.

Smoking cessation intervention for vulnerable groups – exemplified by severe mental disorder

Different groups of smokers might have different needs and outcomes, which may also be the case in regard to severe mental illness. Internationally, mental illness is high on the agenda at the WHO ⁸¹, as well as the European Union (EU) ⁸². The USA has recommend ISCI to all smokers ¹, and the National Institute for Health and Care Excellence (NICE) has special smoking cessation recommendations for ‘acute, maternity and mental health services’ ⁸³, specifically stating to ‘provide intensive support for people using acute and mental health services’.

Patients diagnosed with mental disorders have a higher premature mortality ^{84,85}. Smoking prevalence is high in this vulnerable group ^{84,86–89}; these individuals are more likely to be heavy smokers ⁸⁴, and smoking is an important factor in their 15–20-year reduced life expectancy ⁸⁴. Smoking cessation improves physical and mental health ⁹⁰, reduces morbidity, and contributes to longevity ⁸⁶. Furthermore, 61–71% of smokers with a mental disorder want to quit ^{84,91} and a recent review concluded that smoking cessation treatments appear to be equally effective in psychiatric patients with severe mental disorder (SMD) as in the general public ⁹².

Nationally, mental illness is one of the most common chronic diseases, with 1 in 5 experiencing symptoms of one mental illness or more every year ⁹³. In the 2017 national cross sectional health profile, 9.6% reported sustaining a mental illness lasting longer than 6 months ⁴¹. Compared with the Danish population, the prevalence of smokers with mental disorder is twice as high, and they are 2.5 times more likely to be heavy smokers ⁸⁴. In contrast, the difference in proportions between those who wants to quit in the two groups is only 6% ⁸⁴ (table 2).

	People with mental disorders		Danish population (age 15 or older)	
	Proportion	Number	Proportion	Number
Population size	~10%	478,000	100%	4,780,000
Smoking prevalence	38.8%	185,464	19.8%	946,440
Heavy smokers	25.9%	48,035	10.1%	95,590
Want to quit	68.3%	126,672	74.5%	705,098

Table 2: Smoking characteristics in people with mental disorders compared with the Danish population as a whole. Based on ⁸⁴.

Research of real-life data shows nearly the same effect for disadvantaged groups and heavy smokers as for the background population: 26–29% versus 33% remain smoke-free six months after a GSP ^{94,95}. It is appropriate to examine whether this result also applies to patients with SMD or whether there is a need to develop new programmes for this important subgroup of smokers.

Smoking cessation interventions in Denmark

Smoking cessation programmes were first launched in the 1990s in Denmark to help smokers quit. Since then, these programmes have been a constant element in the national smoking prevention, corresponding to the “Offer help to quit tobacco use” in the WHO MPOWER measurements.

The GSP has been the standard SCI for groups since 1995 ⁶², and in 2003, a corresponding concept for individual smoking cessation (SC) counselling followed ⁶¹. Training curricula for smoking cessation counsellors and teaching materials has been continuously developed and revised through the Cancer Society ^{61–63}.

Over time, there have been changes in terms of to the providers of the interventions and how many smokers were recruited. The most pervasive change was a national healthcare reform

implemented in 2007 ^{96,97}. The reform caused major changes, when responsibility for citizen-oriented smoking prevention was moved from the former counties to the new municipalities ⁹⁸. Prior to the national healthcare reform, the SCI was evenly distributed between four providers: municipalities, hospitals, counties and pharmacies ⁹⁸. After the reform, the municipalities managed to maintain their own activities as well as to pick up on the countries' efforts with no association to the hospitals. In contrast, all activity in the hospitals was virtually shut down by the end of 2010 ⁹⁸. Consequently, the number of smokers attending an SCI dropped from 7,320 smokers in 2006 to 6,119 smokers in 2010, representing a decrease of 16.4% ⁹⁸. In 2012, the activities of the municipalities were no longer maintained, with only 4,461 smokers registered.

National smoking cessation databases

Some countries have developed a national system for monitoring the effect of SCI. A Google-search revealed two national smoking cessation databases, one in Scotland and one in Denmark. A survey of existing smoking cessation databases (conducted in 33 European countries in 2009) identified 9 databases. In addition to Scotland and Denmark, databases were established in the Czech Republic, Hungary, Iceland, Luxembourg, Malta, Romania and Spain. Five countries (France, Croatia, Poland, Slovenia, and Ukraine) did not respond to the enquiries ⁹⁹. In addition to the Scottish and Danish databases being the only publicly displayed databases, they were also found to be the most extensive in terms of registered smokers and smoking cessation clinics, and they were the most developed regarding the user interface, securing the data quality, and extracting outcomes data.

Figure 6: Regions and municipalities reporting data to the SCDB. The red areas do not report to the database. Based on ¹⁰⁰.

In Denmark, the national smoking cessation database (SCDB) was established in 2001. Today, approximately 185 smoking cessation providers from all five health regions of Denmark report data from their smoking cessation programmes to the SCDB (figure 6). Municipalities, counties, regional facilities, hospitals, pharmacies, GPs and other private actors within the smoking cessation area report data. The SCDB holds information on almost 120,000 smokers regarding the programme, smokers and setting. In addition, approximately 10% of all smokers attending a quality assured SCI (i.e., using a smoking cessation clinic reporting to the SCDB for monitoring and quality assessment of their activities) do not consent to have their data registered in the database.

The SCDB primarily serves as a quality database, reporting activities implemented in the clinical daily routine, according to five quality indicators and standards. In the quality approach, the effect is measured according to the number of smokers completing the intervention given. In addition to being a quality database, the SCDB also maintains a strong focus on research, and in the research approach the intention-to-treat principle applies ¹⁰¹ (i.e., the nominator is given by the number included regardless of compliance). This terminology is illustrated in figure 7, which shows that 10,272 smokers were registered in the SCDB in 2016.

Figure 7: Example of a flowchart of participants registered in the SCDB in 2016. The five quality indicators used in the SCDB are marked with an *. The participants are divided based on compliance ¹⁰⁰. Translated and adjusted with permission from the Smoking Cessation Database Secretariat.

Efficacy and effectiveness research

Clinical trials are often classified as ‘effectiveness’ or ‘efficacy’ research based on the approach ¹⁰² (table 3). Throughout this thesis, the term ‘effectiveness research’ will be used to label research based on interventions taking place in a real-life setting, and ‘efficacy research’ that will be used to categorise interventions held under specific conditions, such as randomised clinical trials (RCTs).

Although it is possible to divide studies into categories of efficacy or effectiveness research, the terms could be considered as a spectrum rather than as dichotomous terms ^{102–107}.

	Experimental interventions	Real-life interventions
Also known as	Efficacy trials or explanatory trials	Effectiveness trials or pragmatic trials
Answers the question	Can this work? (i.e., measurable effect)	Does this work in practice? (i.e., utility in routine practice)
Study conditions	Ideal and highly controlled setup	Real-life setup
Study sample	Homogeneous / Selected	Heterogeneous / Reflects routine practice
Comparison group	Often no-treatment or placebo	Often usual care
Compliance	Intensive resources dedicated to obtaining high compliance	Not necessarily high / Reflects routine practice
Clinician skills	Control of clinicians’ skill levels	Normal clinical staff
Validity	High internal validity	High external validity
Treatment protocol	Standardised treatment protocol	Less standardised treatment
Outcome measure	Closely linked to the intervention	Important to the patient
Analysis	Intention to treat / Per protocol	Intention to treat / Adjusted analysis

Table 3: Differences between efficacy and effectiveness research. Based on ^{102–108}.

This thesis builds on the model described in figure 8, illustrating how treatments based on the evidence-based medicine (EBM) approach are implemented into daily clinical routines ^{109,110}. After implementation, it is clinically relevant to follow-up the effects of the implemented treatments, which can be performed using effect-databases and quality assessment tools, such as the SCDB, from where data can ultimately be used to conduct effectiveness research, thereby procuring evidence for the EBM approach.

Figure 8: Process of evidence-based medicine (EBM) being implemented in a daily routine; hopefully, the effects of new procedures are followed up through effect-databases (quality assessment).

Objective and hypothesis

Study I:

Objective: To review the efficacy of intensive smoking cessation interventions (ISCI) in adult daily smokers compared with shorter interventions (SI), measured as continuous abstinence in the short and long term.

Hypothesis: ISCI is more effective than SI.

Study II:

Objective: To describe the Danish Smoking Cessation Database (SCDB) and the most common smoking cessation interventions registered.

Hypothesis: No hypothesis

Study III:

Objective: To compare the effectiveness of the GSP with other in-person smoking cessation interventions in Denmark after implementation in real life and to identify the factors associated with successful quitting.

Hypothesis: GSP will be the most effective intervention after 6 months.

Study IV:

Objective: To compare the effectiveness of the GSP in smokers with and without a diagnosed severe mental disorder (SMD) and to identify the factors associated with successful quitting.

Hypothesis: Smokers with a severe mental disorder will be less likely to be continuously smoke-free after six months than smokers without a mental disorder.

Materials and methods

The overall aim of this thesis (to evaluate the effect of ISCI on successful quitting) was accomplished through four studies. The study flow is illustrated in figure 9.

Trial profile for the thesis

Figure 9: Trial profile for the studies included in the thesis. HP: Health Promotion.

Methods of the included studies

The designs and methods of the scientific studies are described in table 4.

	Manuscript I	Paper II	Paper III	Paper IV
Title	Efficacy of intensive face-to-face interventions for smoking cessation – a systematic review with meta-analyses	The Danish Smoking Cessation Database	Effectiveness of the Gold Standard Programme compared with other smoking cessation interventions in Denmark: a cohort study	Effectiveness of the Gold Standard Programme (GSP) for Smoking Cessation on smokers with and without a severe mental disorder—a Danish cohort study
Design	Systematic review with meta-analysis	Methodology paper describing a national health register	Prospective cohort study	Prospective cohort study
Data collection	Summation of data from 18 RCTs published between 1988-2017	Data validity evaluated based on summary data from smokers registered in the SDCB	Analysis of data from the SCDB, collected on interventions initiated from 2001-2013	Analysis of data from the SCDB initiated 2006-2016 and cross-referenced with the National Patient Register.
Participants	10,131 adult (≥18 years) daily smokers. 6,177 randomised to ISCI, and 3,668 to SI. 168 smokers were excluded, as they did not receive either ISCI or SI	Smokers registered in the SDCB	46,287 smokers of ≥15 years of age, undergoing an in-person smoking cessation intervention in a real-life setting	38,293 adult smokers (≥18 years) with or without a mental disorder, registered in the SCDB after undergoing a GSP in a real-life setting
Main outcome	Successful quitting measured as CA or PP on short- (end of intervention) and long-term (12 or 6-months)	Description of national SCI, method of data collection, coverage, data completeness and validity of the register	Self-reported continuous abstinence at the 6-month follow-up	Self-reported continuous abstinence at the 6-month follow-up
Analysis	Meta-analyses using a random effects model	Descriptive statistics	Descriptive statistics and mixed-effect logistic regression	Descriptive statistics and mixed-effect logistic regression
Publication	Submitted	Published	Published	Published

Table 4: A concise description of the methods used in the four papers included in this thesis.

CA: Continuous abstinence, GSP: Gold Standard Programme, ISCI: Intensive Smoking Cessation Intervention, PP: Point Prevalence, RCT: Randomised Controlled Trial, SCDB: Smoking Cessation Database, SCI: Smoking Cessation Intervention, SI: Shorter Intervention.

Results and discussion

In this section, a short summary of the results of each study will be reported, followed by a discussion across studies. The section is concluded with a description of potential bias and limitations.

Summary of findings

Efficacy of intensive smoking cessation interventions

The literature search in manuscript I resulted in a total of 9,569 papers, after removing duplicate studies. After a systematic screening, 18 relevant studies remained to be included in the analyses 111,112,121–128,113–120. To retrieve the needed information, seven additional publications were used 129–135.

Characteristics of the included studies

The 18 included RCTs were published between 1988 and 2017, and they included a total of 10,131 smokers, ranging from 30 – 5,887 smokers. Most studies were conducted in Western Europe (7 studies) or in the USA (8 studies). The remaining 3 studies were performed in Syria, Qatar and Australia. Men accounted for 55% of the smokers; most included studies were population studies, and some included patients only.

Quality assessment

The risk of bias in the individual studies was assessed using the Cochrane Collaboration tool 136. Overall, the risk of bias was moderate (figure 10). Even though it is not possible to blind participants and personnel when evaluating behavioural interventions, such as smoking cessation interventions, all included studies were assessed a high risk of bias, as it is not possible to eliminate the possibility that the awareness of the randomisation influenced the outcome.

Figure 10: Summary of the assessment of risk of bias of the studies included in the systematic review.

The body of the evidence was rated using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) approach 137. Four outcomes were assessed: continuous abstinence (CA) and point prevalence (PP), both in the short- and the long term. The evidence was considered moderate for CA-short, CA-long, and PP-long, while it was ranked low for PP-short. The short term was defined as the end of the intervention; the long term primarily used a 12-month follow-up, and if that was not available, then a 6-month follow-up was used.

Successful quitting

Many methods are used to define successful quitting 2,138,139. The primary outcome was CA, and the secondary outcome was PP, and the definitions proposed by Hughes et al 2 was applied. Both CA and PP were evaluated in the short and long term. Nine studies reported CA, 7 studies reported on the short term 112–114,119,120,127,128, and 7 studies reported on the long term 112,113,118,119,123,127,128. PP was reported in 16 studies, 11 on the short term 112,113,128,114–

Four meta-analyses were conducted using a random effects model. The summary results are shown in figure 11.

Figure 11: Results of the four meta-analyses conducted in the systematic review (I).

In all analyses, heterogeneity was evident, with I^2 values exceeding 50%, prompting an exploration of the heterogeneity. Three analyses were conducted testing 1) studies with a high risk of bias (in addition to the ‘blinding of participants and personnel’ category), 2) studies with a low (or no) follow-up rate, and 3) studies with low (or no) compliance rates. The first two analyses only to a minor extent explained the heterogeneity. The last analysis, however, drastically reduced the heterogeneity, leading to $I^2 = 0$ in the primary outcomes.

Effectiveness of smoking cessation interventions in real life

In this section, papers II-IV will be introduced, as they are all based on the SCDB. Study II is a methodology paper that describes the SCDB and the most common SCI registered in the database. Papers III and IV are both prospective cohort studies, based on data from the SCDB. The main focus in study III was to estimate the effect of different SCIs, while study IV aimed to evaluate successful quitting among smokers diagnosed with SMD. The proportion of successful quitting was evaluated, and modifiable and non-modifiable factors associated with a positive outcome were identified after controlling for confounders.

The Danish Smoking Cessation Database

Paper II is a methodology study, presenting the SCDB as a national health register. The database was established in response to the smoking cessation activities emerging nationally in the 1990s. It was initiated as a research project to ensure a systematic data collection to evaluate the effectiveness and cost-effectiveness of SCI. From 2006, the database was transformed from a research project to a quality assessment database, and accordingly, five quality indicators and standards (table 5) were developed and implemented.

Indicator	Standard
1) Compliance with smoking cessation intervention	80%
2) Smoke-free at the end of the programme	80%
3) Follow-up rate	80%
4) Smoke-free after 6 months	50%
5) Satisfaction	90%

Table 5: The five indicators and corresponding quality standard developed and implemented in the SCDB in 2006.

Since 2001, the inception data were collected at baseline, with a mandatory follow-up on the effect. During the research project (2001–2005), follow-ups were conducted twice after 6 and 12 months. Since 2006, follow-up has only been mandatory after 6 months (figure 12). In addition, in 2006, the registration of the CPR number (Central Personal Register, a unique number that includes the date of birth and sex and is assigned to all Danes at birth, as well as to immigrants)

was implemented in the SCDB. The addition of the CPR number ensured linkage to other central registers in Denmark.

Figure 12: 'Data collection in the SCDB over time (CPR: a unique number including date of birth and gender assigned to all Danes at birth and to immigrants)'. Reprint with permission from the Clinical Health Promotion Journal.

Based on publicly available homepages and course descriptions, it was estimated that the SCI registered in the SCDB represents approximately 80-90% of all SCIs with personal attendance that take place in Denmark. The data completeness was rated as high as 100% of the questions on the SCI, and at least 95% of the questions on the basic- and follow-up questionnaire were completed. Automated and manual data validation routines have been implemented to ensure a high quality of the data registered in the SCDB.

Differences in SC interventions

Paper III included data collected from 2001-2013, and the study cohort consisted of 46,287 smokers 15 years of age and older undergoing an in-person smoking cessation intervention in a real-life setting. Five face-to-face interventions were compared: GSP (n = 40,651), health promotion counselling (n = 140), crash courses (n = 183), Come & Quit (n = 1,592), and other (3,721). The GSP and the Come & Quit interventions were ISCIs, as were most of the programmes categorised as others'.

Of the 46,287 included smokers, 26% were lost to follow-up, leaving 34,235 participants to be included in the outcome analysis. The crude quit-rates are shown in Figure 13. The minimal effect (i.e., the worst case scenario) was calculated using the Russell criteria ¹³⁸, in which non-respondents are presumed to have relapsed. The high end of the successful quitting range was calculated using the RECORD criteria ¹⁴⁰, in which non-respondents are assumed to resemble respondents.

Overall, 33% of the smokers were continuously smoke-free after 6 months, 24% when using the Russell standards. The crude quit-rates for women and men are shown in figure 13.

The final adjusted logistic regression model was run for men and women separately, as the fully adjusted model showed that women were significantly less likely to be CA after 6 months compared to men: OR: 0.83 (95% CI, 0.79-0.87).

Figure 13: Crude quit-rates for women and men, according to the five interventions compared. The overall, crude quit-rate for all interventions and across sex is indicated by the light blue area.

Four predictors were associated with the outcome for both men and women (figure 14). Compliance was the most pronounced predictor for successful quitting in both sexes, increasing the odds by more than 3.5 times.

Figure 14: Four predictors were associated with outcome. Compliance was the most pronounced predictor for successful quitting in both sexes.

Severe mental disorder and smoking cessation

Paper IV utilised data collected from 2006-2016, and the study cohort consisted of 36,945 smokers of 18 years of age or older who attended a GSP. Smokers diagnosed with an SMD (7,162) were compared to smokers without mental disorder (29,783). To identify smokers with a diagnosis of SMD, all participants were cross-referenced with data from the National Patient Register. Smokers diagnosed with any of the International Classification of Diseases, 10th edition (ICD-10) diagnoses F00-F99, except F17 'Mental and behavioural disorders due to use of tobacco', were classified as having a mental disorder. SMD was defined as having one or more of the following diagnoses: schizophrenia spectrum (F20-F29), affective disorders (F30-F34), anxiety (F40-F42, F43.1), personality disorders (F60-F69), and substance use disorder (F10-16, F18-F16).

Smokers with SMD had a reduced chance of staying continuously abstinent after six months, OR: 0.74 (95% CI, 0.68-0.80), and all subgroups of smokers with SMD were less likely to be successful, however, the result was not statistically significant for the subgroup of smokers diagnosed with anxiety. The crude quit rate was 38.0% in smokers without a mental disorder and 29.4% in smokers with SMD. The predictors associated with continuous abstinence are presented in figure 15.

Finally, the impact of dual diagnosis, defined as having any of the SMD diagnoses in combination with substance use disorder (SUD), was examined. The chance of a successful outcome was reduced by 18-43% in all subgroups of smokers with a dual diagnosis.

Figure 15: Factors associated with outcome in smokers with SMD versus smokers without a mental disorder.

Discussion

Together, these studies have added new knowledge to the effect of intensive smoking cessation interventions on successful quitting. The three intervention studies (I, III, IV) showed that smokers benefit from ISCI. The summary effect of the systematic review (I) showed a 3-fold increase in the likelihood of staying continuously abstinent when randomised to an ISCI compared with SI. The effectiveness of ISCI was recovered when evaluating the GSP after implementation in real life. The magnitude of the effect of ISCI differed between the two study designs, with the highest effect observed in the cohort study (III), with 25% of participants being continuously abstinent after 6 months, compared to 15% in the systematic review (I), indicating that it is possible to effectively implement ISCI into a clinical daily routine on a national scale.

The higher magnitude in the cohort study was surprising, as the effectiveness in real life was expected to be less pronounced compared with the effect obtained in RCTs. Due to the nature of the study designs, RCTs (characterised by, for example, optimal trial conditions, patient selection and increased staff resources and competences) often overestimate the effect compared with interventions implemented in daily clinical practice ¹⁰³. Part of the explanation could be that only the most motivated smoker would turn to a smoking cessation clinic for help and accept inclusion in a national database.

Regarding the systematic review (I), similar conclusions on the comparison of ISCI and shorter interventions have been reached in previous reviews ^{69,70,72}, however, all used different definitions of interventions. In contrast, reviewing other types of SCIs identified only minor effect. Internet-based interventions showed the largest effect among 'Internet plus behavioural support' with 13% successful quitters vs. 8% in the control group ¹⁴¹. Telephone counselling showed a similar effect when testing additional proactive calls, with a quit rate of 11% in the interventions group versus 8% in the control group ¹⁴². The outcomes used in both reviews were a mix of continuous abstinence (CA) and point prevalence (PP) ^{141,142}. Compared to these interventions, using the same definition for the measured outcomes, ISCI proved to be highly effective, with 30% successful quitters versus 11% in the SI.

Recruitment

It is important to consider the recruitment procedures when comparing outcomes of the different studies, particularly how they differ from the patients treated in the daily clinical routine. Some of the smokers in manuscript I were recruited through the media, while others were recruited consecutively as they entered (e.g., a lung-clinic). It would be expected that the participants recruited through the media would be more motivated, as they would have to actively join the trial. However, the trials using consecutive inclusion were actually the interventions with the

largest outcome in general. This finding could, however, be explained by the increased focus and resources in the clinical setting in the project period and the fact that these patients were offered inclusion as they entered the hospital to receive treatment for another disease, which was possibly affected by smoking.

It is a clinical experience that indications and guidelines for treatments may alter during the daily routine, meaning that treatments tested in one group of patients are changed and transferred to other groups of patients. Therefore, it is imperative to monitor the effect in real life to assess and improve the quality of a treatment.

However, both study designs are important. Register-based cohort trials are often based on real-life data, however, they can only point out the possibility of an association between exposure and outcome. Hence, in the absence of RCTs, it is difficult to interpret whether an observed effect can be attributed to the intervention given or if it is a result of other contextual factors ¹⁰⁸. To establish the association between intervention and outcome, it is necessary to test the intervention using an RCT approach.

Gender perspective

Concerning the gender perspective, contradictory results have been found in multiple studies, with some studies showing no differences between men and women ^{143–145} and others confirming that women are less successful in quitting than men ^{146–148}. Bohadana et al. have suggested that these findings might have to do with gender differences in dependence, both related to nicotine and behaviour ¹⁴⁸. The cohort study (III) showed women were significantly less likely to be successful, though the GSP was effective for both men and women. This difference was also seen, and even more pronounced, in regard to the 'health promotion counselling', which seemed to be highly effective for men, though based on a very small group (constituting 140 smokers or 0.3% of the cohort). Even so, this finding raises an interesting hypothesis that short interventions could be more effective than intensive interventions among men. Based on a meta-analysis finding that low-intensity counselling (3-10 minutes) only has an increased effect of 6% compared with no contact ⁶⁶, it is likely that the results reflect that there might be a highly selected and specific group of men who benefit more from short interventions. This possibility should be investigated further.

Compliance

The most important predictor identified in the cohort studies (III-IV) was compliance, which was not surprising because a strong dose-response has been observed between compliance to the programme and successful quitting ¹⁴⁹. A similar dose-response effect was seen when comparing interventions divided into categories according to the number of sessions ⁶⁶. It could be argued that the effect is observed because those who relapse discontinue the programme; however, it might be those confident that they can manage on their own who leave the programme. Another likely explanation of the impact of compliance could be that by attending the planned smoking cessation sessions, the smokers actually gain knowledge and tools to handle the challenges in staying abstinent. Either way, it would be relevant to investigate new ways to keep people in the programme.

Vulnerable and disadvantaged groups

Patients diagnosed with SMD constitute a special group of smokers. They had a continuous abstinence rate of 28% after 6 months, compared to 38% in the comparison group (IV), hence being less successful. Despite of the lower chance of being successful, more than one in four smokers with an SMD succeed in their quitting attempt.

Thus far, there are few studies of SCI and SMD. A recent review based on 4 studies noted that the

effect of specialist smoking cessation programmes for smokers with severe mental illness is still uncertain ¹⁵⁰, and another review investigating the effect of smoking cessation advice in this vulnerable group, did not find any studies on the topic ⁶⁸. Furthermore, reviews exploring subgroups of patients with mental illness have identified few studies of behavioural smoking cessation treatment ¹⁵¹⁻¹⁵³. However, an RCT evaluating the efficacy of initiating smoking cessation treatment in patients admitted to a psychiatry ward has reported promising results ¹⁵⁴, as has an observational study on depressive smokers that used routinely collected data from a smoking cessation clinic in the Czech Republic and showed quit-rates of 25-33%, depending on the severity of depression when entering into the SCI, as well as improvements in the depression level in abstinent patients ¹⁵⁵. In addition, a recent real-world study on clinical practice showed no significant difference in abstinence rates between smokers with or without a mental disorder at the end of the 6-week treatment ¹⁵⁶. Unfortunately, the study only included a short-term follow-up, and, therefore, it is not possible to determine whether the results would be maintained in the long term ¹⁵⁶.

Interestingly, SUD alone did not seem to impact the quit rate compared to other subgroups of mental disorders (IV). Nevertheless, a dual diagnosis lowered the proportion of successful quitters (IV), and it seemed to have some impact on the ability to quit smoking. A review has demonstrated a small effect of SC supportive pharmacotherapy with or without counselling in people in treatment for or recovery from SUD, with a combination of treatments being the most effective, resulting in 16% becoming abstinent compared to 9% in the control group ¹⁵⁷. Counselling alone showed no effect ¹⁵⁷. It would be relevant to also examine the effect of SCI on the outcome of treatment for dual diagnoses.

Contrary to the assumption that patients with SMD might be reluctant to consent to be registered in a database noting their SCI, an inspection of the data showed that this assumption does not seem to be the case. The proportion of smokers in the Danish population with a mental disorder can be estimated at 19.6% ⁸⁴. In comparison, a similar level (19.4%) of smokers registered in the SCDB had a diagnosis of at least one mental disorder. This finding may be limited to Danish conditions, which are characterised by a long tradition of registration of health data.

Worldwide, patients with psychiatric diagnoses do not seem to get the same attention and offers in relation to SCI ^{154,158}, in spite of their relatively high motivation to quit (nearly comparable to the background population) ^{84,91}. It is a shame, as this finding seems to be a waste of a major potential on multiple levels, a potential that has been harvested elsewhere (e.g., in Catalonia). Here, enhancement of tobacco control in the mental health setting has been high on the agenda for the last decade ¹⁵⁸. Likewise, promising results have been observed in an RCT initiating SCI in an inpatient setting in the USA ¹⁵⁴. Due to the physical and mental ⁹² improvements after smoking cessation, it is important that this group of smokers is offered help to quit, and it would be relevant to transfer these experiences to other mental health care settings, while carefully monitoring the effects.

ISCI, such as GSP, has an effect across socioeconomic and vulnerable groups, including smokers diagnosed with SMD. Previous studies on GSP have found similar results in disadvantaged ⁹⁴, heavy ⁹⁵, pregnant ¹⁵⁹, elderly ¹⁶⁰ and, most recently, young ¹⁶¹ groups of smokers, and the programme has also proven to be effective in randomised and controlled trials ^{127,162}.

Therefore, an expansion of the implementation of GSP or other ISCI should be considered among lower socioeconomic groups and other vulnerable groups, which have the highest prevalence of smokers.

Policy

Although, Denmark was rewarded the highest level of implementation of the MPOWER measure “Offer help to quit tobacco use”, still no more than 1% of the smokers in Denmark receive a face-to-face SCI per year based on 2016 numbers ¹⁰⁰. This result is far less than the nationally and internationally recommended 5%. In England, 5-10% of the smokers are treated by the Stop Smoking Service every year ¹⁴⁴. Full implementation of the national guidelines would have a high impact in Denmark.

Another challenge is the general recommendation from WHO and many countries to implement shorter intervention, particularly the shortest one (brief intervention) or just advices, instead of ISCI. The shorter interventions are the cheaper choice, and they have proven to be somewhat effective, even though the absolute quit rates are low (13% versus 11% in the no contact control group) ⁶⁶.

It has been established that tobacco control policies, such as taxes and bans, are indeed effective, especially in preventing adolescents from starting to smoke ⁴², even so, they have still not been implemented in Denmark. As a result of the evasive political decisions, other initiatives are put into action by parties outside the political system, the latest being a declaration from parts of the retail industry to implement under the counter sale of tobacco in multiple supermarkets effectuated by the end of 2018 ¹⁶³.

Overall, the potential for tobacco control, including both prevention and treatment of current smokers, has not been fulfilled in Denmark.

Bias and limitations

A strength of RCTs is that participants are randomly assigned to a treatment, thus aiming at evening out known or, equally important, unknown confounders, which is not the case in cohort studies, where dealing with such factors is a ticklish subject. Known predictors and confounders can be taken into account by adjusting for them in the analysis, however, only as long as data are available, not to mention the factors that are still unknown.

Regarding the data sources, the SCDB contains information about smokers who attend a face-to-face smoking cessation intervention. Nationally, it is recommended but not mandatory to report smoking cessation interventions to the SCDB, therefore, it does not cover all activities (II). Based on publicly available information, such as webpages and course descriptions it was estimated that approximately 80-90% of all face-to-face interventions were reported, and only 3 of the 98 Danish municipalities never reported data to the database (II). Because all smokers in Denmark have access to smoking cessation clinics free of charge and without referral, the selection bias is considered limited.

Non-respondents

In the cohort studies, 25-31% of the smokers did not respond after 6 months, therefore, it was important to consider if and how the non-respondents differed from the respondents (III-IV). Bias introduced by non-respondents is hard to address, however, using baseline data to describe the differences between the groups might provide some insight to how the non-respondents affect the results. In both cohort studies (III-IV), compliance was the factor that differed most between respondents and non-respondents, the first being more likely to be compliant with the programme. As compliance is the strongest predictor associated with successful outcomes the results based on the RECORD criteria ¹⁴⁰ are most likely overestimating the quit rate if extrapolated to the non-respondents (III-IV).

In contrast to the non-responders, only small differences were observed regarding the group of smokers who were followed-up and those who were intentionally not followed-up by the smoking cessation clinic due to a local administrative decision, not in line with the agreement signed by the smoking cessation clinic and the SCDB (III-IV). However, it might only influence the estimation to a minor degree.

Reporting of smoking cessation

In the RCTs included in the systematic review (I), smoking cessation was reported in many different manners. The reporting differs in multiple regards, such as the type of abstinence (e.g., CA or PP), the time interval from cessation to follow-up and the validation. The fact that trials do not use a common standard introduces both bias and limitations for the individual RCT, as well as for the meta-analysis. The smoking status in the cohort studies (III-IV) were based on non-validated self-reporting, which can be considered as a limitation due to potentially overestimation of the quit-rates. Several studies reported different results regarding this subject. It has been shown that patients are likely to overestimate their self-reported success by 3-6%, compared with outcomes validated by carbon monoxide (CO) ¹⁶⁴⁻¹⁶⁶, while a Canadian study showed no significant difference between self-reporting and validation by urinary cotinine levels ¹⁶⁷. Even so, the groups would be comparable as there was no reason to assume that an overestimation would be unevenly distributed among the different groups of smokers.

Data sources

Data from both Danish registers provide a high completeness and precision, and the amount of missing data was expected to be low (II) ¹⁶⁸. Misclassification on all levels can occur in both registers, though the occurrence was expected to be low, as the SMD diagnosis originates from psychiatric departments and hospitals reporting to the National Patient Register as a code of practice ¹⁶⁸, and a structured smoking profile is required for inclusion in the SCDB.

When using routinely collected health data in research one should rightly contemplate the implications in doing so ¹⁴⁰. The SCDB was established as a research database to answer questions about the effectiveness of specific programmes or specific subgroups of smokers. In spite of a shift towards quality assessment in 2006, the SCDB still maintained a strong focus on the research approach. Therefore, the implications of using data from the SCDB for research projects within the scope of the database (III-IV) are considered minimal.

Ethical considerations

None of the studies included in this thesis had any direct patient contact.

The SCDB was approved by the Danish Data Protection Agency (2000-54-0013) since the inception of the data registration. Every patient registered in the SCDB provided informed consent to the registration and follow-up. The patients filled in the registration papers themselves, with personally identifiable data for the registration.

Furthermore, the research project (including III-IV) was approved by the Data Protection Agency (Ref. 2010-41-5463) and the National Board of Health, as particularly sensitive data from the National Patient Register are used (IV). The research project (III-IV) was also registered with the National Committee on Health Research Ethics (H-C-FSP-2010-049).

The systematic review (I) was conducted in agreement with a pre-submitted protocol (PROSPERO 2017 CRD42071059879).

Conclusion and impact

This thesis adds to the evidence on intensive smoking cessation interventions (ISCI) through different study designs. The results supported the use of ISCI. The systematic review (I) showed that ISCI significantly increased the chance of successful quitting for all four predefined outcomes. The cohort studies found that ISCI, in the form of the Danish Gold Standard Programme (GSP), was the only intervention that was effective for both men and women. The GSP was found to be significantly less effective for patients diagnosed with severe mental disorder (SMD); still, one in four participants of this vulnerable subgroup of smokers remained successful quitters after 6 months.

For the continuous development of better SCI, it is important to evaluate the effects using a high-quality design such as RCT and to follow-up participants to observe the effects after implementation in real life (e.g., in a national SCDB).

Perspectives

Implications for practice

Based on the evidence generated in this thesis combined with the proportion of smokers who want to quit, ISCI should be recommended and implemented wherever possible. In a Danish context, it would be relevant to turn up the number of offers to at least reach the national and international goal of treating 5% of smokers every year.

The studies in the clinical setting showed the highest quit-rates. Therefore, it would be highly relevant to continue to use this setting for implementation to achieve high impact among smoking patients in primary and specialist care, including psychiatric care.

Other important settings for SCI are the local health care centres in the municipalities and pharmacies, where it is possible to reach out to a broader group of smokers.

Internationally, implementation of ISCI should be considered in relation to cultural traditions, staff competences, patient preferences and the healthcare structure amongst others, as the geographic distribution of the included studies were limited to Denmark and other Western nations.

Implications for smokers

For the individual smoker, it is paramount to be both informed and able to choose the most effective SCI for obtaining successful quitting, and thereby have a better chance to reduce the high risk of related morbidity and pre-mortality. This process would also benefit their families and surrounding people in reducing exposure to second-hand smoke. Furthermore, being familiar with the results of ISCI, including that compliance to the programme may be of great importance, could add to the motivation to quit.

Implications for society

In the future, offering the most effective SCI would not only be beneficial to individual smokers but also to the general health of the public, as well as to society at large. In the light of the effectiveness of ISCI it would be of great impact in Denmark to fully implement the national guideline targeting treatment of 5% of smokers each year, which would lower the otherwise stagnated smoking prevalence, and, in the long term, result in a reduction in morbidity and mortality from smoking. In addition, an increased focus on disadvantaged and other vulnerable groups of smokers would reduce health inequality.

Implications for research

In this thesis, the effect of ISCI as defined in the 'Clinical Practice Guideline for Treatment of Tobacco Use and Dependence'¹ was investigated. ISCI is a package of many different elements and it was beyond the scope of this thesis to look further into the influence of the different components of the interventions. The systematic review did, however, reveal a wide range of different interventions all nested within the ISCI definition. The interventions differed in regard to multiple aspects, such as the time interval, number and length of sessions, use of pharmacotherapy, intervention method (e.g., counselling, motivational interviewing, cognitive behavioural therapy), and format (group or individual sessions). It would, therefore, be interesting to investigate whether it is possible to identify the most effective elements (individual or in combination). It would also be relevant to evaluate the effect of an IT-solution based on ISCI, with and without personal contact.

Even though the GSP has demonstrated effects across many subgroups of smokers, it would still be interesting to investigate if this is also the case for other groups of patients, such as patients with cancer, asthma, or diabetes, as smoking cessation would have a major impact in reducing the development and/or aggravation of these diseases.

Based on the solid experiences of using a national quality database to assure the quality of SCI, it would be obvious to explore the feasibility of using the same approach to develop and implement similar databases to assess and improve the quality of other health promoting interventions for lifestyle risk factors (e.g., hazardous alcohol consumption, malnutrition, overweight/obesity, as well as insufficient physical activity).

References

1. Clinical Practice Guideline Treating Tobacco Use and Dependence 2008 Update Panel, Liaisons and S. A Clinical Practice Guideline for Treating Tobacco Use and Dependence: 2008 Update. *Am J Prev Med.* 2008;35(2):158-176. doi:10.1016/j.amepre.2008.04.009.
2. Hughes JR, Keely JP, Niaura RS, Ossip-Klein DJ, Richmond RL, Swan GE. Measures of abstinence in clinical trials: issues and recommendations. *Nicotine Tob reasearc.* 2003;5(1):13-25.
3. MacKay J, Schluger N. Global tobacco epidemic. In: Loddenkemper R, Kreuter M, eds. *The Tobacco Epidemic, Ed 2, Rev. and Ext.* Vol 42. Basel: Karger; 2015:19-26. doi:10.1159/000369320.
4. Forouzanfar MH, Afshin A, Alexander LT, et al. Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks, 1990–2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet.* 2016;388(10053):1659-1724. doi:10.1016/S0140-6736(16)31679-8.
5. *WHO Report on the Global Tobacco Epidemic, 2017. Monitoring Tobacco Use and Prevention Policies.* World Health Organization; 2017. http://www.who.int/tobacco/global_report/2017/en/. Accessed May 7, 2018.
6. Danmarks Statistik. www.statistikbanken.dk/FOLK1.
7. *Sygdomsbyrden i Danmark.* Sundhedsstyrelsen; 2016. <https://sundhedsstyrelsen.dk/da/sygdom-og-behandling/~media/C3ACA2467BEE41B49726532872563FFA.ashx>.
8. Stanhope B-J, Burdette WJ, Cochran WG, et al. *Smoking and Health. Report of the Advisory Committee to the Surgeon General of the Public Health Service.* Public Health Service Publication No. 1103; 1964.
9. Eriksen M, Mackay J, Schluger NW, Gomeshtapeh FI, Drope J. *The Tobacco Atlas.* 5th ed. Atlanta: The American Cancer Society, Inc.; 2015.
10. *Smoking and Health. A Report of the Royal College of Physicians of London on Smoking in Relation to Cancer of the Lung and Other Diseases.* London; 1962.
11. Wipfli H, Samet JM. One Hundred Years in the Making: The Global Tobacco Epidemic. *Annu Rev Public Health.* 2016;37(1):149-166. doi:10.1146/annurev-publhealth-032315-021850.
12. Lopez AD, Collishaw NE, Piha T. A descriptive model of the cigarette epidemic in developed countries. *Tob Control.* 1994;3(3):242-247. doi:10.1136/tc.3.3.242.
13. Thun M, Peto R, Boreham J, Lopez AD. Stages of the cigarette epidemic on entering its second century. *Tob Control.* 2012;21(2):96-101. doi:10.1136/tobaccocontrol-2011-050294.
14. Novello A. Surgeon General's report on the health benefits of smoking cessation. *Public Health Rep.* 1990;105(6):545-548.
15. Cahn WZ, Drope J, Hamill S, et al. *The Tobacco Atlas.* 6th ed. (Drope J, Schluger NW, eds.). American Cancer Society, Inc.; 2018. www.tobaccoatlas.org.
16. Sundhedsstyrelsen, Kræftens Bekæmpelse, Danmarks Lungeforening, Hjerteforeningen. *Danskernes Rygevaner 2017.* <https://www.sst.dk/da/udgivelser/2018/danskernes-rygevaner-2017>. Accessed May 11, 2018.
17. *WHO Report on the Global Tobacco Epidemic, 2011. Warning about the Dangers of Tobacco.* World Health Organization; 2011. http://www.who.int/tobacco/global_report/2011/en/.
18. Juel K, Sørensen J, Brønnum-Hansen H. *Risikofaktorer Og Folkesundhed i Danmark.* Statens Institut for Folkesundhed; 2006.
19. Lim SS, Vos T, Flaxman AD, et al. A comparative risk assessment of burden of disease and injury attributable to 67 risk factors and risk factor clusters in 21 regions, 1990-2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2013;380(9859):2224-2260. doi:10.1016/S0140-6736(12)61766-8.
20. Kjølner M, Juel K, Kamper-Jørgensen F. Rygning. In: *Folkesundhedsrapporten Danmark 2007.* Statens Institut for Folkesundhed; 2007:221-234.
21. Andres RL, Day MC. Perinatal complications associated with maternal tobacco use. *Semin Neonatol SN.* 2000;5(3):231-241. doi:10.1053/siny.2000.0025.
22. Benowitz NL, Dempsey DA. Pharmacotherapy for smoking cessation during pregnancy. *Nicotine Tob Res.* 2004;6(Suppl. 2):189-202. doi:10.1080/14622200410001669169.
23. Yang Q, Wen SW, Phillips K, Oppenheimer L, Black D, Walker MC. Comparison of maternal risk factors between placental abruption and placenta previa. *Am J Perinatol.* 2009;26(4):279-286.
24. Wisborg K, Henriksen TB, Obel C, Skajaa E, Ostergaard JR. Smoking during pregnancy and hospitalization of the child. *Pediatrics.* 1999;104(4):e46.
25. Wisborg K, Kesmodel U, Henriksen TB, Olsen SF, Secher NJ. A prospective study of smoking during pregnancy and SIDS. *Arch Dis Child.* 2000;83(3):203-206.

26. McDonnell-Naughton M, McGarvey C, O'Regan M, Matthews T. Maternal smoking and alcohol consumption during pregnancy as risk factors for sudden infant death. *Ir Med J.* 2012;105(4):105-108.
27. Wisborg K, Kesmodel U, Henriksen TB, Olsen SF, Secher NJ. Exposure to tobacco smoke in utero and the risk of stillbirth and death in the first year of life. *Am J Epidemiol.* 2001;154(4):322-327.
28. Wisborg K, Henriksen TB. *Tobak Og Graviditet. Er Der Behov for Forebyggelse? Findes Der Effektive Metoder?* Sundhedsstyrelsen; 2003.
<https://www.sst.dk/da/udgivelser/2004/~media/37BCB6A15904416C88B4B9582E9A40A9.ashx>.
29. Cnattingius S. The epidemiology of smoking during pregnancy: smoking prevalence, maternal characteristics, and pregnancy outcomes. *Nicotine Tob Res Off J Soc Res Nicotine Tob.* 2004;6 Suppl 2:S125-40.
30. Schneider S, Huy C, Schütz J, Diehl K. Smoking cessation during pregnancy: a systematic literature review. *Drug Alcohol Rev.* 2010;29(1):81-90. doi:10.1111/j.1465-3362.2009.00098.x.
31. McLeish AC, Zvolensky MJ. Asthma and cigarette smoking: a review of the empirical literature. *J Asthma.* 2010;47(4):345-361. doi:10.3109/02770900903556413.
32. Thomson NC, Chaudhuri R, Livingston E. Asthma and cigarette smoking. *Eur Respir J Off J Eur Soc Clin Respir Physiol.* 2004;24(5):822-833. doi:10.1183/09031936.04.00039004.
33. Althuis M, Sexton M, Prybylski D. Cigarette smoking and asthma symptom severity among adult asthmatics. *J Asthma.* 1999;36(3):257-264.
34. Polosa R, Thomson NC. Smoking and asthma: dangerous liaisons. *Eur Respir J Off J Eur Soc Clin Respir Physiol.* 2012. doi:10.1183/09031936.00073312.
35. Tønnesen H, Nielsen PR, Lauritzen JB, Møller AM. Smoking and alcohol intervention before surgery: evidence for best practice. *Br J Anaesth.* 2009;102(3):297-306. doi:10.1093/bja/aen401.
36. Grønkjær M, Eliassen M, Skov-Ettrup LS, et al. Preoperative Smoking Status and Postoperative Complications. *Ann Surg.* 2014;259(1):52-71. doi:10.1097/SLA.0b013e3182911913.
37. WHO Report on The Global Tobacco Epidemic 2008. *The MPOWER Package.* World Health Organization; 2008. <http://www.who.int/tobacco/mpower/2008/en/>.
38. Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ.* 2004;328(7455):1519. doi:10.1136/bmj.38142.554479.AE.
39. Christensen AI, Ekholm O, Davidsen M, Juel K. 5.2 Rygning. In: *Sundhed Og Sygelighed i Danmark 2010 & Udviklingen Siden 1987.* Statens Institut for Folkesundhed, Syddansk Universitet; 2012:146-157.
40. Christensen AI, Ekholm O, Davidsen M, Juel K. *Sundhed Og Sygelighed i Danmark 2010 & Udviklingen Siden 1987.* Statens Institut for Folkesundhed, Syddansk Universitet; 2012.
41. *Danskernes Sundhed - Den Nationale Sundhedsprofil 2017.* Sundhedsstyrelsen; 2018.
<https://www.sst.dk/da/udgivelser/2018/danskernes-sundhed-den-nationale-sundhedsprofil-2017>.
42. Vestbo J, Pisinger C, Bast LS, Gyrd-Hansen D. *Forebyggelse Af Rygning Blandt Børn Og Unge. Hvad Virker?* Vidensråd for forebyggelse; 2018. http://www.vidensraad.dk/sites/default/files/vff-rapport_forebyggelse_af_rygning_hvad_virker_indhold_og_forside_2.pdf.
43. *Capacity Assessment on the Implementation of Effective Tobacco Control Policies in Denmark.* Copenhagen: WHO regional Office for Europe; 2018.
<https://www.cancer.dk/dyn/resources/File/file/5/7055/1517908352/implementation-of-effective-tobacco-control-policies-in-denmark-.pdf>.
44. Fagerström K. The epidemiology of smoking: health consequences and benefits of cessation. *Drugs.* 2002;62(Suppl 2):1-9.
45. Ezzati M, Lopez AD. Estimates of global mortality attributable to smoking in 2000. *Lancet.* 2003;362(9387):847-852. doi:10.1016/S0140-6736(03)14338-3.
46. Kræftens Bekæmpelse. Voksnes rygevaner. <https://www.cancer.dk/forebyg/undga-roeg-og-rygning/fakta-om-rygning/voksnes-rygevaner/>. Accessed June 9, 2018.
47. *Dødsårsagsregisteret: Tal Og Analyse.* Sundhedsstyrelsen; 2015.
<http://sundhedsdatastyrelsen.dk/da/tal-og-analyser/analyser-og-rapporter/andre-analyser-og-rapporter/doedsaarsagsregisteret>.
48. *WHO Framework Convention on Tobacco Control.* World Health Organization; 2005.
<http://apps.who.int/iris/bitstream/handle/10665/42811/9241591013.pdf;jsessionid=9C7CAAE69A7A3EE1E62B5D92EF34EB85?sequence=1>. Accessed June 11, 2018.
49. Pötschke-Langer M, Schotte K, Szilagyi T. The WHO Framework Convention on Tobacco Control. In: Loddenkemper R, Kreuter M, eds. *The Tobacco Epidemic, Ed 2, Rev. and Ext.* Vol 42. Basel: Karger; 2015:149-157. doi:10.1159/000369441.

50. World Health Organization. WHO Framework Convention on Tobacco Control. WHO. 2017. <http://www.who.int/fctc/cop/about/en/>. Accessed June 11, 2018.
51. Roemer R, Taylor A, Lariviere J. Origins of the WHO Framework Convention on Tobacco Control. *Am J Public Health*. 2005;95(6):936-938. doi:10.2105/AJPH.2003.025908.
52. World Health Organization. Parties to the WHO Framework Convention on Tobacco Control. WHO. 2017. http://www.who.int/fctc/signatories_parties/en/. Accessed June 14, 2018.
53. World Health Organization. Draft comprehensive global monitoring framework and targets for the prevention and control of noncommunicable diseases (A66/8). 2013. http://apps.who.int/gb/ebwha/pdf_files/WHA66/A66_8-en.pdf. Accessed May 3, 2016.
54. Joossens L, Raw M. The Tobacco Control Scale: a new scale to measure country activity. *Tob Control*. 2006;15(3):247-253. doi:10.1136/tc.2005.015347.
55. Joossens L, Raw M. *The Tobacco Control Scale 2016 in Europe*. Association of European Cancer Leagues; 2017. <http://www.tobaccocontrolscalescale.org/wp-content/uploads/2017/03/TCS-2016-in-Europe-COMLETE-LoRes.pdf>. Accessed June 14, 2018.
56. Thomson G, Edwards R, Wilson N, Blakely T. What are the elements of the tobacco endgame? *Tob Control*. 2012;21(2):293-295. doi:10.1136/tc.2010.040881.
57. Tobacco Endgame - Tobaksfakta. <http://www.tobaksfakta.se/tobacco-endgame/>. Accessed June 26, 2018.
58. Sundhedsstyrelsen, Kræftens Bekæmpelse, Danmarks Lungeforening, Hjerteforeningen. Danskernes rygevaner. <https://www.sst.dk/da/sundhed-og-livsstil/tobak/udgivelser/danskernes-rygevaner>. Accessed February 19, 2018.
59. Percentage smokers (per cent), by sex, age, contents and year. Statbank Norway. <https://www.ssb.no/en/statbank/table/05307/tableViewLayout1?rxid=fba52324-e745-43b1-8740-058b118535f6>. Accessed June 14, 2018.
60. Hilding-Nørkjær H, Koefoed BG, Falk J, Tværmose P. *Metoder Til Rygeafvænning - Dokumentation Og Anbefalinger*. Sundhedsstyrelsen og Nationalt Center for Rygestop; 2003. <https://www.sst.dk/da/udgivelser/2003/~media/5255EA0B02B9439FACBA393751EDFAF3.ashx>. Accessed June 26, 2018.
61. Lind M, Jaspers K. *Manual Til Individuel Rygeafvænning*. 5. udgave. Kræftens Bekæmpelse; 2011. https://www.cancer.dk/dyn/resources/File/file/7/1637/1385430538/1_forord_litt.pdf.
62. Lind M, Jaspers K. *Manual Til Rygeafvænning i Grupper*. 3. udgave. Kræftens Bekæmpelse; 2013. https://www.cancer.dk/dyn/resources/File/file/5/3605/1399893540/01startdel_web.pdf.
63. Lind M, Jaspers K, Kjaer NT. *Rygeafvænning - Teori Og Praksis for Professionelle Rådgivere*. 1. udgave. Kræftens Bekæmpelse; 2009.
64. Stoplinien. <http://stoplinien.dk/>. Accessed June 26, 2018.
65. The Tobacco Use and Dependence Clinical Practice Guideline Panel, Staff, and Consortium Representatives S, Representatives and C. A Clinical Practice Guideline for Treating Tobacco Use and Dependence: A US Public Health Service Report. *JAMA J Am Med Assoc*. 2000;283(24):3244-3254. doi:10.1001/jama.283.24.3244.
66. Tobacco Use and Dependence Guideline Panel. *Treating Tobacco Use and Dependence: 2008 Update*. Rockville (MD): US Department of Health and Human Services; 2008 May.; 2008. <https://www.ncbi.nlm.nih.gov/books/NBK63952/>.
67. Lindson-Hawley N, Thompson TP, Begh R. Motivational interviewing for smoking cessation. *Cochrane Database Syst Rev*. 2015. doi:10.1002/14651858.CD006936.pub3.
68. Khanna P, Clifton A V, Banks D, Tosh GE. Smoking cessation advice for people with serious mental illness. In: Clifton A V, ed. *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd; 2016. doi:10.1002/14651858.CD009704.pub2.
69. Thomsen T, Villebro N, Møller AM. Interventions for preoperative smoking cessation. *Cochrane database Syst Rev*. 2014;3:CD002294. doi:10.1002/14651858.CD002294.pub4.
70. Rigotti NA, Clair C, Munafò MR, Stead LF. Interventions for smoking cessation in hospitalised patients. *Cochrane Database Syst Rev*. 2012;5:CD001837. doi:10.1002/14651858.CD001837.pub3.
71. Stead LF, Buitrago D, Preciado N, Sanchez G, Hartmann-Boyce J, Lancaster T. Physician advice for smoking cessation. *Cochrane Database Syst Rev*. 2013. doi:10.1002/14651858.CD000165.pub4.
72. Lancaster T, Stead LF. Individual behavioural counselling for smoking cessation. *Cochrane Database Syst Rev*. 2017;3:CD001292. doi:10.1002/14651858.CD001292.pub3.
73. Tonnesen H, Juhl CB, Lauritzen JB, Egholm JW, Adami J. Systematic review and meta-analyses: clinical health promotion and postoperative complications. PROSPERO 2018 CRD42018094537. https://www.crd.york.ac.uk/prospéro/display_record.php?RecordID=94537. Accessed June 26, 2018.

74. Wiebel FJ. Chemistry and primary toxicity of tobacco and tobacco smoke. In: Loddenkemper R, Kreuter M, eds. *The Tobacco Epidemic, Ed 2, Rev. and Ext.* Vol 42. Basel: Karger; 2015:37-46. doi:10.1159/000369323.
75. Kræftens Bekæmpelse. Fakta om rygning. <https://www.cancer.dk/forebyg/undga-roeg-og-rygning/fakta-om-rygning/>. Accessed June 20, 2018.
76. Jarvis MJ. Why people smoke. *BMJ*. 2004;328(7434):277-279. doi:10.1136/bmj.328.7434.277.
77. Hansen NS. *Tobakkens Biologi*. 1. udgave. Tobaksskaderådet; 1994. http://sundhedsstyrelsen.dk/publ/PUBL1994/TOBAKKENS_BIOLOGI.PDF.
78. Zaveri NT, Rollema H, Swan GE. Nicotine Dependence. In: Loddenkemper R, Kreuter M, eds. *The Tobacco Epidemic, Ed 2, Rev. and Ext.* Vol 42. Basel: Karger; 2014:47-57. doi:10.1159/000369324.
79. Mundt K, Kann A. Forebyggelsessamtaler om tobak og alkohol. *Sygeplejersken*. 2000;(2):34-40.
80. Mundt K, Fugleholm AM, Hedegaard AM, Jepsen JM, eds. *Rygeophør Op Sygehus - Fakta, Metode Og Anbefalinger*. Klinisk Enhed for Sygdomsforebyggelse og Netværk af forebyggende sygehus i Danmark; 2001.
81. *Mental Health Action Plan 2013-2020*. World Health Organization; 2013. http://apps.who.int/iris/bitstream/10665/89966/1/9789241506021_eng.pdf?ua=1%5Chttp://apps.who.int/iris/bitstream/10665/89966/1/9789241506021_eng.pdf.
82. EU joint action on mental helath and wellbeing. *European Framework for Action on Mental Health and Wellbeing*. Brussels; 2016. https://ec.europa.eu/health/sites/health/files/mental_health/docs/ev_20161006_co07_en.pdf.
83. NICE National Institute for Health and Care Excellence. Public health guideline: Smoking: acute, maternity and mental health services (PH48). <https://www.nice.org.uk/guidance/ph48/chapter/1-Recommendations#recommendation-3-provide-intensive-support-for-people-using-acute-and-mental-health-services>. Accessed September 7, 2017.
84. Nordentoft M, Krogh J, Lange P, Moltke A. *Psykisk Sygdom Og Ændringer i Livsstil*. Vidensråd for forebyggelse; 2015. http://www.vidensraad.dk/sites/default/files/vidensraad_for_forebyggelse_psykisk_sygdom_aendringer_i_livsstil_2015.pdf.
85. Nordentoft M, Wahlbeck K, Hällgren J, et al. Excess mortality, causes of death and life expectancy in 270,770 patients with recent onset of mental disorders in Denmark, Finland and Sweden. *PLoS One*. 2013;8(1):e55176. doi:10.1371/journal.pone.0055176.
86. Bobes J, Arango C, Garcia-Garcia M, Rejas J. Healthy lifestyle habits and 10-year cardiovascular risk in schizophrenia spectrum disorders: an analysis of the impact of smoking tobacco in the CLAMORS schizophrenia cohort. *Schizophr Res*. 2010;119(1-3):101-109. doi:10.1016/j.schres.2010.02.1030.
87. Holma IAK, Holma KM, Melartin TK, Ketokivi M, Isometsä ET. Depression and smoking: a 5-year prospective study of patients with major depressive disorder. *Depress Anxiety*. 2013;30(6):580-588. doi:10.1002/da.22108.
88. Lasser K, Boyd JW, Woolhandler S, Himmelstein DU, McCormick D, Bor DH. Smoking and Mental Illness. *JAMA*. 2000;284(20):2606. doi:10.1001/jama.284.20.2606.
89. de Leon J, Diaz FJ. A meta-analysis of worldwide studies demonstrates an association between schizophrenia and tobacco smoking behaviors. *Schizophr Res*. 2005;76(2-3):135-157. doi:10.1016/j.schres.2005.02.010.
90. Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P. Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ*. 2014;348:g1151. doi:10.1136/bmj.g1151.
91. *Smoking and Mental Health*. Report CR1. London: Royal College of Physicians and Royal College of Psychiatrists; 2013. https://cdn.shopify.com/s/files/1/0924/4392/files/smoking_and_mental_health_-_full_report_web.pdf?7537870595093585378.
92. Banham L, Gilbody S. Smoking cessation in severe mental illness: what works? *Addiction*. 2010;105(7):1176-1189. doi:10.1111/j.1360-0443.2010.02946.x.
93. Vendsborg P, Lindhart A, Gram T, Bentzen J. *Tal Til Psyken - Fakta Om Psykisk Sundhed Og Psykisk Sygdom 2015*. Psykiatrifonden; 2015. <http://www.psykiatrifonden.dk/media/1417569/tal-til-psyken-september-2016.pdf>.
94. Neumann T, Rasmussen M, Ghith N, Heitmann BL, Tønnesen H. The Gold Standard Programme: smoking cessation interventions for disadvantaged smokers are effective in a real-life setting. *Tob Control*. 2013;22(e9):Epub 2012 Jun 16. doi:10.1136/tobaccocontrol-2011-050194.
95. Neumann T, Rasmussen M, Heitmann BL, Tønnesen H. Gold Standard Program for Heavy Smokers in a Real-Life Setting. *Int J Environ Res Public Health*. 2013;10(9):4186-4199. doi:10.3390/ijerph10094186.

96. Andersen PT, Jensen J-J. Healthcare reform in Denmark. *Scand J Public Health*. 2010;38(3):246-252. doi:10.1177/1403494809350521.
97. Christiansen T. Ten years of structural reforms in Danish healthcare. *Health Policy (New York)*. 2012;106(2):114-119. doi:10.1016/j.healthpol.2012.03.019.
98. Rasmussen M, Ammari ABH, Pedersen B, Tønnesen H. Smoking cessation intervention activities and outcome before, during and after the national Healthcare Reform in Denmark. *Clin Heal Promot*. 2012;2(1):26-35. doi:10.29102/clinhp.12005.
99. Rindel L. *Kortlægning Og Vurdering Af Eksisterende Kliniske Rygestopdatabaser i Europa*. Københavns Universitet; 2009.
100. Grønbaek AS, Rasmussen M, Tønnesen H. *Rygestopbasens Årsrapport - Aktiviteter Afholdt i 2016 Med Opfølgning i 2017*. Rygestopbasens sekretariat; 2017.
101. Leuchs AK, Brandt A, Zinserling J, Benda N. Disentangling estimands and the intention-to-treat principle. *Pharm Stat*. 2017;16(1):12-19. doi:10.1002/pst.1791.
102. Fritz JM, Cleland J. Effectiveness versus efficacy: more than a debate over language. *J Orthop Sports Phys Ther*. 2003;33(4):163-165. doi:10.2519/jospt.2003.33.4.163.
103. Singal AG, Higgins PDR, Waljee AK. A primer on effectiveness and efficacy trials. *Clin Transl Gastroenterol*. 2014;5(2):e45. doi:10.1038/ctg.2013.13.
104. Thorpe KE, Zwarenstein M, Oxman AD, et al. A pragmatic-explanatory continuum indicator summary (PRECIS): a tool to help trial designers. *J Clin Epidemiol*. 2009;62(5):464-475. doi:10.1016/j.jclinepi.2008.12.011.
105. Gartlehner G, Hansen RA, Nissman D, Lohr KN, Carey TS. A simple and valid tool distinguished efficacy from effectiveness studies. *J Clin Epidemiol*. 2006;59(10):1040-1048. doi:10.1016/j.jclinepi.2006.01.011.
106. Merckx P, Paugam-Burtz C, Boudinet S, Bonnet A, Mantz J. Explanatory versus Pragmatic Trials? The Methods Make the Difference. *Anesthesiology*. 2008;108(3):542-543. doi:10.1097/ALN.0b013e3181650ecc.
107. Kim SY. Efficacy versus Effectiveness. *Korean J Fam Med*. 2013;34(4):227. doi:10.4082/kjfm.2013.34.4.227.
108. Ernst E, Pittler M. LETTER TO THE EDITOR Efficacy or effectiveness? *J Intern Med*. 2006;260:488-490. doi:10.1111/j.1365-2796.2006.01707.x.
109. Bak Andersen I, Matzen P, eds. *Evidensbaseret Medicin*. 2. udgave. Gads Forlag; 2007.
110. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ*. 1996;312(7023):71-72. doi:10.1136/BMJ.312.7023.71.
111. Anthonisen NR, Connett JE, Kiley JP, et al. Effects of Smoking Intervention and the Use of an Inhaled Anticholinergic Bronchodilator on the Rate of Decline of FEV1: The Lung Health Study. *JAMA J Am Med Assoc*. 1994;272(19):1497-1505. doi:10.1001/jama.1994.03520190043033.
112. El Hajj M, Kheir N, Al Mulla A, Shami R, Fanous N, Mahfoud Z. Effectiveness of a pharmacist-delivered smoking cessation program in the State of Qatar: a randomized controlled trial. *BMC Public Health*. 2017;17:215. doi:10.1186/s12889-017-4103-4.
113. Baker A, Richmond R, Haile M, et al. A Randomized Controlled Trial of a Smoking Cessation Intervention Among People With a Psychotic Disorder. *Am J Psychiatry*. 2006;163(11):1934-1942. doi:10.1176/ajp.2006.163.11.1934.
114. Asfar T, Weg M V, Maziak W, Hammal F, Eissenberg T, Ward KD. Outcomes and adherence in Syria's first smoking cessation trial. *Am J Heal Behav*. 2008;32(2):146-156. doi:10.5555/ajhb.2008.32.2.146.
115. Alterman AI, Gariti P, Mulvaney F. Short- and long-term smoking cessation for three levels of intensity of behavioral treatment. *Psychol Addict Behav*. 2001;15(3 PG-261-4):261-264.
116. Jorenby DE, Smith S, Fiore M, et al. Varying nicotine patch dose and type of smoking cessation counseling. *JAMA*. 1995;274(17):1347-1352.
117. Tonnesen P, Fryd V, Hansen M, et al. Two and four mg nicotine chewing gum and group counselling in smoking cessation: an open, randomized, controlled trial with a 22 month follow-up. *Addict Behav*. 1988;13(1):17-27.
118. Nohlert E, Tegelberg Å, Tillgren P, Johansson P, Rosenblad A, Helgason ÅR. Comparison of a high and a low intensity smoking cessation intervention in a dentistry setting in Sweden: a randomized trial. *BMC Public Health*. 2009;9:121. doi:10.1186/1471-2458-9-121.
119. Wewers ME, Ferketich AK, Harness J, Paskett ED. Effectiveness of a nurse-managed, lay-led tobacco cessation intervention among ohio appalachian women. *Cancer Epidemiol Biomarkers Prev*. 2009;18(12):3451-3458. doi:10.1158/1055-9965.epi-09-0952.
120. Kehlet M, Heeseman S, Tonnesen H, Schroeder T V. Perioperative smoking cessation in vascular surgery: Challenges with a randomized controlled trial. *Trials*. 2015;16:441. doi:10.1186/s13063-

015-0965-x.

121. Mujika A, Forbes A, Canga N, et al. Motivational interviewing as a smoking cessation strategy with nurses: an exploratory randomised controlled trial. *Int J Nurs Stud.* 2014;51:1074-1082. doi:10.1016/j.ijnurstu.2013.12.001.
122. Grant K, Northrup J, Agrawal S, Olsen D, McIvor C, Romberger D. Smoking cessation in outpatient alcohol treatment. *Addict Disord their Treat.* 2003;2:41-46.
123. Romand R, Gourgou S, Sancho-Garnier H. A randomized trial assessing the Five-Day Plan for smoking cessation. *Addiction.* 2005;100:1546-1554. doi:10.1111/j.1360-0443.2005.01215.x.
124. Williams GC, Niemiec CP, Patrick H, Ryan RM, Deci EL. The importance of supporting autonomy and perceived competence in facilitating long-term tobacco abstinence. *Ann Behav Med.* 2009;37:315-324. doi:10.1007/s12160-009-9090-y.
125. Gifford E, Kohlenberg B, Hayes S, et al. Does acceptance and relationship focused behavior therapy contribute to bupropion outcomes? A randomized controlled trial of functional analytic psychotherapy and acceptance and commitment therapy for smoking cessation. *Behav Ther.* 2011;42:700-715. doi:10.1016/j.beth.2011.03.002.
126. Brunner Frandsen N, Sørensen M, Hyldahl TK, Henriksen RM, Bak S. Smoking cessation intervention after ischemic stroke or transient ischemic attack. A randomized controlled pilot trial. *Nicotine Tob Res.* 2012;14(4):443-447. doi:10.1093/ntr/ntr233.
127. Møller AM, Villebro N, Pedersen T, Tønnesen H. Effect of preoperative smoking intervention on postoperative complications: a randomised clinical trial. *Lancet.* 2002;359(9301):114-117. doi:10.1016/S0140-6736(02)07369-5.
128. Mohiuddin SM, Mooss AN, Hunter CB, Grollmes TL, Cloutier DA, Hilleman DE. Intensive Smoking Cessation Intervention Reduces Mortality in High-Risk Smokers With Cardiovascular Disease. *Chest.* 2007;131(2):446-452. doi:10.1378/chest.06-1587.
129. Buist AS. The US Lung Health Study. *Respirology.* 1997;2:303-307.
130. Connett JE, Kusek JW, Bailey WC, O'Hara P, Wu M. Design of the Lung Health Study: a randomized clinical trial of early intervention for chronic obstructive pulmonary disease. *Control Clin Trials.* 1993;14:3S-19S.
131. Frikart M, Etienne S, Cornuz J, Zellweger JP. Five-day plan for smoking cessation using group behaviour therapy. *Swiss Med Wkly.* 2003;133(3-4):39-43. doi:2003/03/smw-10068.
132. Kanner RE, Connett JE, Williams DE, Buist AS. Effects of randomized assignment to a smoking cessation intervention and changes in smoking habits on respiratory symptoms in smokers with early chronic obstructive pulmonary disease: the Lung Health Study. *Am J Med.* 1999;106:410-416.
133. Nohlert E, Ohrvik J, Tegelberg A, Tillgren P, Helgason AR. Long-term follow-up of a high- and a low-intensity smoking cessation intervention in a dental setting-- a randomized trial. *BMC Public Health.* 2013;13(1):592. doi:10.1186/1471-2458-13-592.
134. Villebro NM, Pedersen T, Møller AM, Tønnesen H. Long-term effects of a preoperative smoking cessation programme. *Clin Respir J.* 2008;2(3):175-182. doi:10.1111/j.1752-699X.2008.00058.x.
135. Williams GC, McGregor H, Sharp D, et al. A self-determination multiple risk intervention trial to improve smokers' health. *J Gen Intern Med.* 2006;21:1288-1294. doi:10.1111/j.1525-1497.2006.00621.x.
136. Higgins JPT, Altman DG SJ (Editors). Chapter 8: Assessing risk of bias in included studies. In: *Cochrane Handbook for Systematic Reviews of Interventions Version 5.2.0 (Updated June 2017)*; 2017. www.training.cochrane.org/handbook.
137. Guyatt GH, Oxman AD, Vist GE, et al. Rating Quality of Evidence and Strength of Recommendations: GRADE: An Emerging Consensus on Rating Quality of Evidence and Strength of Recommendations. *BMJ Br Med J.* 2008;336:924-926. doi:10.2307/20509578.
138. West R, Hajek P, Stead L, Stapleton J. Outcome criteria in smoking cessation trials: proposal for a common standard. *Addiction.* 2005;100(3):299-303. doi:10.1111/j.1360-0443.2004.00995.x.
139. Velicer WF, Prochaska JO, Rossi JS, Snow MG. Assessing outcome in smoking cessation studies. *Psychol Bull.* 1992;111(1):23-41.
140. Benchimol EI, Smeeth L, Guttman A, et al. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) Statement. *PLOS Med.* 2015;12(10):e1001885. doi:10.1371/journal.pmed.1001885.
141. Taylor GMJ, Dalili MN, Semwal M, Civljak M, Sheikh A, Car J. Internet-based interventions for smoking cessation. *Cochrane Database Syst Rev.* 2017. doi:10.1002/14651858.CD007078.pub5.
142. Stead LF, Hartmann-Boyce J, Perera R, Lancaster T. Telephone counselling for smoking cessation. *Cochrane Database Syst Rev.* 2013. doi:10.1002/14651858.CD002850.pub3.
143. Puente D, Cabezas C, Rodriguez-Blanco T, et al. The role of gender in a smoking cessation intervention: a cluster randomized clinical trial. *BMC Public Health.* 2011;11:369.

doi:10.1186/1471-2458-11-369.

144. Dobbie F, Hiscock R, Leonardi-Bee J, et al. Evaluating Long-term Outcomes of NHS Stop Smoking Services (ELONS): a prospective cohort study. *Heal Technol Assess*. 2015;19(95). doi:10.3310/hta19950.
145. Fidler J, Ferguson SG, Brown J, Stapleton J, West R. How does rate of smoking cessation vary by age, gender and social grade? Findings from a population survey in England. *Addiction*. 2013;108:1680-1685. doi:10.1111/add.12241.
146. Bauld L, Bell K, McCullough L, Richardson L, Greaves L. The effectiveness of NHS smoking cessation services: a systematic review. *J Public Health (Oxf)*. 2010;32(1):71-82. doi:10.1093/pubmed/fdp074.
147. Martínez C, Martínez-Sánchez JM, Ballbè M, et al. Smoking cessation in hospital workers: effectiveness of a coordinated program in 33 hospitals in Catalonia (Spain). *Cancer Nurs*. 2012;35(5):327-336. doi:10.1097/NCC.0b013e3182321389.
148. Bohadana A, Nilsson F, Rasmussen T, Martinet Y. Gender differences in quit rates following smoking cessation with combination nicotine therapy: influence of baseline smoking behavior. *Nicotine Tob Res*. 2003;5(1):111-116. doi:10.1080/1462220021000060482.
149. Ghith N, Ammari ABH, Rasmussen M, Frølich A, Cooper K, Tønnesen H. Impact of compliance on quit rates in a smoking cessation intervention: population study in Denmark. *Clin Heal Promot*. 2012;2(3):111-119.
150. Peckham E, Brabyn S, Cook L, Tew G, Gilbody S. Smoking cessation in severe mental ill health: what works? an updated systematic review and meta-analysis. *BMC Psychiatry*. 2017;17(1):252. doi:10.1186/s12888-017-1419-7.
151. Tsoi DT, Porwal M, Webster AC. Interventions for smoking cessation and reduction in individuals with schizophrenia. In: Tsoi DT, ed. *Cochrane Database of Systematic Reviews*. Chichester, UK: John Wiley & Sons, Ltd; 2013. doi:10.1002/14651858.CD007253.pub3.
152. Gierisch JM, Bastian LA, Calhoun PS, McDuffie JR, Williams JW. Smoking Cessation Interventions for Patients with Depression: A Systematic Review and Meta-analysis. *J Gen Intern Med*. 2012;27(3):351-360. doi:10.1007/s11606-011-1915-2.
153. George TP, Wu BS, Weinberger AH. A Review of Smoking Cessation in Bipolar Disorder: Implications for Future Research. *J Dual Diagn*. 2012;8(2):126-130. doi:10.1080/15504263.2012.671717.
154. Prochaska JJ, Hall SE, Delucchi K, Hall SM. Efficacy of initiating tobacco dependence treatment in inpatient psychiatry: a randomized controlled trial. *Am J Public Health*. 2014;104(8):1557-1565. doi:10.2105/AJPH.2013.301403.
155. Stepankova L, Kralikova E, Zvoliska K, et al. Depression and Smoking Cessation: Evidence from a Smoking Cessation Clinic with 1-Year Follow-Up. *Ann Behav Med*. 2017;51(3):454-463. doi:10.1007/s12160-016-9869-6.
156. Castaldelli-Maia JM, Loreto AR, Guimarães-Pereira BBS, et al. Smoking cessation treatment outcomes among people with and without mental and substance use disorders: An observational real-world study. *Eur Psychiatry*. 2018;52:22-28. doi:10.1016/J.EURPSY.2018.02.005.
157. Apollonio D, Philipps R, Bero L. Interventions for tobacco use cessation in people in treatment for or recovery from substance use disorders. *Cochrane Database Syst Rev*. 2016. doi:10.1002/14651858.CD010274.pub2.
158. Ballbè M, Gual A, Nieva G, Saltó E, Fernández E. Deconstructing myths, building alliances: a networking model to enhance tobacco control in hospital mental health settings. *Gac Sanit*. 2016;30(5):389-392. doi:10.1016/j.gaceta.2016.04.017.
159. Rasmussen M, Heitmann BL, Tønnesen H. Effectiveness of the Gold Standard Programmes (GSP) for Smoking Cessation in Pregnant and Non-Pregnant Women. *Int J Environ Res Public Health*. 2013;10(8):3653-3666. doi:10.3390/ijerph10083653.
160. Kehlet M, Schroeder T V, Tønnesen H. The Gold Standard Program for Smoking Cessation is Effective for Participants Over 60 Years of Age. *Int J Environ Res Public Health*. 2015;12(3):2574-2587. doi:10.3390/ijerph120302574.
161. Iyer SSC. *Effectiveness of Smoking Cessation Interventions for Young Smokers in Denmark – the Geographic Region’s Influence on the Successful Quitting among Young People Using Data from the Smoking Cessation Database*. University of Copenhagen; 2018.
162. Borglykke A, Pisinger C, Jørgensen T, Ibsen H. The effectiveness of smoking cessation groups offered to hospitalised patients with symptoms of exacerbations of chronic obstructive pulmonary disease (COPD). *Clin Respir J*. 2008;2(3):158-165. doi:10.1111/j.1752-699X.2008.00055.x.
163. Danmarks Radio. Flere supermarkeder vil skjule cigaretter i butikkerne | Ligetil | DR. <https://www.dr.dk/ligetil/flere-supermarkeder-vil-skjule-cigaretter-i-butikkerne>. Accessed June 29, 2018.

164. Ferguson J, Bauld L, Chesterman J, Judge K. The English smoking treatment services: one-year outcomes. *Addiction*. 2005;100 Suppl:59-69. doi:10.1111/j.1360-0443.2005.01028.x.
165. Judge K, Bauld L, Chesterman J, Ferguson J. The English smoking treatment services: short-term outcomes. *Addiction*. 2005;100(Suppl. 2):46-58. doi:10.1111/j.1360-0443.2005.01027.x.
166. Pisinger C, Vestbo J, Borch-Johnsen K, Thomsen T, Jørgensen T. Acceptance of the smoking cessation intervention in a large population-based study: the Inter99 study. *Scand J Public Health*. 2005;33(2):138-145. doi:10.1080/14034940410028370.
167. Wong SL, Shields M, Leatherdale S, Malaisson E, Hammond D. Assessment of validity of self-reported smoking status. *Statistics Canada, Health Reports*, 2012;23(1):1-7.
168. Schmidt M, Schmidt SAJ, Sandegaard JL, Ehrenstein V, Pedersen L, Sørensen HT. The Danish National patient registry: A review of content, data quality, and research potential. *Clin Epidemiol*. 2015;7. doi:10.2147/CLEP.S91125.

Appendix/Papers

Manuscript I

Systematic review:

Efficacy of intensive face-to-face interventions for smoking cessation – a systematic review and meta-analysis

Rasmussen M, Pedersen B, Lauridsen SV, Backer V, Tønnesen H.

(Submitted)

Paper II

Methodology study:

The Danish Smoking Cessation Database

Rasmussen M, Tønnesen H.

Clinical Health Promotion Journal. 2016;6(2):36-42. doi:10.29102/clinph.16006.

Paper III

Cohort study:

Effectiveness of the Gold Standard Programme compared with other smoking cessation interventions in Denmark: a cohort study

Rasmussen M, Fernández E, Tønnesen H.

BMJ Open 2017;7:e013553. doi: 10.1136/bmjopen-2016-013553.

Paper IV

Cohort study:

Effectiveness of the Gold Standard Programme (GSP) for Smoking Cessation on smokers with and without a severe mental disorder—a Danish cohort study

Rasmussen M, Klinge M, Krogh J, Nordentoft M, Tønnesen H.

BMJ Open 2018;8:e021114. doi: 10.1136/bmjopen-2017-021114.