ISOMERIC TRANSFORMATION OF CERTAIN NEW HETEROCYCLIC COMPOUNDS

D. B. Jadhao

Department of Chemistry, Dharampeth M. P. Deo Memorial Science College, Nagpur M.S. (India)
drdbjadhao3@gmail.com**Abstract:**

The isomeric transformation of 2-tetra-O- acetyl - β -D-glucopyranosylamino-4-phenyl-5-aryl/alkylimino-1,3,4-thiadiazoline have been investigated in 5% aqueous sodium carbonate solution under two conditions. When the transformation was tried with heating duration of 60 min., 2-phenyl-3-aryl/alkylimino-4-tetra-O-acetyl- β -D-glucopyranosyl-5-mercapto-1,2,4-triazolines have been isolated almost exclusively. While increasing the heating time to 6-8 h., a mixture of two products, (i) 2-phenyl-3-aryl/alkylimino-4-tetra-O-acetyl- β -D-glucopyranosyl-1-5-hydroxy(enolic)-1,2,4-triazolines (major) and (ii) 2-phenyl-3-aryl/alkylimino-4-tetra-O-acetyl- β -D-glucopyranosyl-5-mercapto-1,2,4-triazolines (minor) have been isolated. The structural assignments have been made on the basis of certain chemical transformations and UV, IR, NMR, mass spectral analysis.

Introduction:

Nucleosides having β -D-glucopyranosyl group at the ring nitrogen of cyclic system or exo-cyclic nitrogen or exocyclic nitrogen of heterocyclic system or cyclic compounds have varied applications in paper, textile and food industries. Some of these compounds have application in **medicinal chemistry** as a diuretic agent, analgesics, antibiotics, bacteriostatic & antituberculous reagents and antitumor agents.

As a part of wider programme of research of synthesis of 2- β -D-glucopyranosylimino-4-H/Phenyl-5-aryl/alkylimino-1,3,4-thiadiazolidines have been carried out by the interaction of 1-tetra-O-acetyl- β -D-glucopyranosyl-4-H/Phenylthiosemicarbazide and alkyl/aryl isothiocyanates followed by deacetylation with methanolic ammonia solution.

The resultant product seems to be of pharmaceutical importance and during the synthesis of it, all required care was taken to use minimum organic solvents. Simple and one/two step reactions are taken into consideration for all chemical transformation.

Experimental: Typical isomeric transformations at two different heating durations are as stated below.

(a) Reaction for heating duration of 30 min.:

In a 100 ml. round bottom flask 50 ml. of 5% aqueous sodium carbonate solution was taken and to it 3.5 g. of 2-tetra-O-acetyl- β -D-

glucopyranosylamino-4-phenyl-5-phenylimino-1,3,4-thiadiazolidines was added, colour of reaction mixture turned pale yellow, after which the reaction mixture was refluxed over boiling water bath for 30 min. The reaction mixture was cooled and filtered. To the filtrate aqueous dil. Hydrochloric acid solution was added to neutralize the alkaline solution, when a solid was obtained. It was filtered and washed thoroughly with distilled water and then dried (2.8g.) solid was isolated, m. p. 119 °C

(b) Reaction for heating duration of 6-8

hours: When 5 g. of 2-tetra-O-acetyl- β -D-glucopyranosylamino-1,3,4-thiadiazoline was dissolved in 5% aqueous sodium carbonate solution (65 ml). The reaction mixture turned slightly brownish in colour. It was stirred thoroughly and refluxed directly over a gentle flame for about 6-8 h. during defluxion slow elimination of hydrogen sulphide was observed. After heating, the reaction mixture was cooled and filtered. The resultant alkaline solution was neutralized by adding dil. Hydrochloric acid solution, when product was precipitated out. It was filtered and washed thoroughly with distilled water and then dried, a white solid (3.2 g) was isolated m. p. 126-132 °C. This product was found to be a mixture of two products. These two products have been separated by column chromatography (silica gel- aqueous ethanol 1: 5). (i) m. p. 125 °C (major) (ii) m. p. 119 °C

EXPERIMENTAL: SYNTHESIS OF INTERMEDIATES

Cyclo condensation

The overall reaction may be represented as follows:-

Where R = (a) p-tolyl, (b) phenyl, (c) p-Cl-phenyl, (d) tert-butyl, (e) methyl, (f) TAG
 TAG = Tetra-O-acetyl- β -D-glucopyranosyl group.
 Ph = Phenyl group.

Table - 1

Isomeric transformation of 2-tetra-O-acetyl- β -D-glucopyranosylamino-4-phenyl-5-aryl/alkylimino-1,3,4-thiadiazolines : Formation of 2-phenyl-3-aryl/alkylimino-4-tetra-O-acetyl- β -D-glucopyranosyl-5-hydroxy/mercapto-1,2,4-triazolines

S.No.	1,3,4-thiadiazoline (III)	5-hydroxy-1,2,3-triazoline (II)	M.P. °C	5-mercapto 1,2,4-triazoline (I)	M.P. °C
1.	.. phenyl, (IIIa)	...3-phenyl-5-hydroxy (IIa)...	125	..3-p-phenyl-5-mercapto (Ia)	119
2.	.. p-tolyl (IIIb)	...3-p-tolyl-5-hydroxy (IIb)...	145	..3-p-tolyl-5-mercapto (Ib)	138
3.	.. p-Cl-phenyl (IIIc)	...3-p-Cl-phenyl-5-hydroxy(IIc)...	193(d)	..3-p-Cl-phenyl-5-mercapto (Ic)	178(d)
4.	.. Tert butyl (IIId)	tert.butyl-5-hydroxy (IIId)...	141	..3-tert.butyl-5-mercapto (Id)	137
5.	.. Methyl (IIIe)	...3-methyl-5-hydroxy (IIe)...	157	..3-methyl-5-mercapto (Ie)	146
6.	.. Tetra-O-acetyl- β -D-glucopyranosyl (IIIff)	...3-tetra-O-acetyl- β -D-glucopyranosyl-5-hydroxy(IIff)	158	..3-tetra-O-acetyl- β -D-glucopyranosyl-5-hydroxy (If)	149

TABLE 3: SPECTRAL DATA OF ACETYLATED PRODUCT:

Compound no.	IR band at	NMR Signal at	Mass
3-a	3320 (NH) Stretching 1750 (C=O) Stretching 1640 (C=N) Stretching 680 (C-S) Stretching 900 (β -isomeric form of pyranosyl ring)	δ , 2.2 ppm (acetyl) δ , 8.1 ppm (NH) δ , 4.13 to 4.21 ppm β -D-Glucopyranosyl	Parent peak at M/e 522, Other prominent peaks also observed.
3-b	3320 (M) (NH) Stretching 1750 (S) (C=O) Stretching 1520 (M) (C=N) Stretching 800 (S) (C-S) Stretching 899 (M) (β Glucoside anomeric ring)	δ , 2.66 ppm (NH) δ , 2.00 ppm (methyl) δ , 7.00 to 7.16 ppm (β aromatic configuration)	—

TABLE-4:SPECTRAL DATA OF DEACETYLATED PRODUCT

Compound no.	IR band at	NMR Signals at	Mass
1-a	3420(m)O-H Sretching 3330(m)N-H Sretching 1510(w)C=N Sretching 600(s) C-S Sretching 891(w) Glucoside deformation (β -anomer)	δ , 2.23 ppm N- H δ , 4.84 ppm O- H δ , 7.14 to 7.80 ppm Aromatic proton δ , 3.68 to 4.20 ppm β -anomeric configuration	Product Shows Parent Peak at M/e 354 However other important fragment peak also observed.

TABLE-4:SPECTRAL DATA OF DEACETYLATED PRODUCT

Compound no.	IR band at	NMR Signals at	Mass
1-a	3420(m)O-H Stretching	δ,2.23 ppm N- H δ,4.84 ppm O- H	Product Shows Parent Peak at M/e 354 However other important fragment peak also observed.
	3330(m)N-H Stretching		
	1510(w)C=N Stretching	δ,7.14to 7.80ppm Aromatic proton δ, 3.68 to 4.20ppm β-anomeric configuration	
	600(s) C-S Stretching		
	891(w) Glucoside deformation (β-anomer)		

Result and Discussion:

The product with m. p. 119 °C was assigned the structure 2-phenyl-3- phenyl imino-4- tetra - O - acetyl- β-D-glucopyranosyl-5-- β-D-glucopyranosyl ring mercapto-1,2,4-triazoline and the product with m. p. 125 °C did not contain sulphur. It contains four nitrogen, out of which three nitrogen were the part of 1,2,4-triazoline ring and one present as imino nitrogen while oxygen was present as an enolic form. The product charred with conc. Sulphuric acid and showed positive Molisch, sTest, optically active. The UV spectrum of the product was recorded and showed lambda max at 270 nm in alcohol. The IR spectrum of the product clearly indicated the presence of ν-OH, ν- -NH, ν- C=N, ν-C=O bands in addition to usual bands of - β-D-glucopyranosyl ring. The NMR spectrum displayed signals due to aromatic nucleus, OH, and NH. It also showed signals due to protons of - β-D-glucopyranosyl ring, mass spectrum of the product shows molecular ion peak at 523, besides this other important fragment peaks at 463, 331, 229, 169, 109, were clearly located.

On the basis of the above fact the product with m. p 125 °C was assigned the structure 2-phenyl- 3- phenylimino-4- tetra-O-acetyl- β-D-glucopyranosyl -5-hydroxy (enolic)-1,2,4-triazoline (IIa) major.

Conclusion:

One can draw a conclusion From the observed fact that Isomeric transformations of certain glucosylated heterocyclic compounds can be done with 5% aqueous sodium carbonate solution by two heating durations that is 30 min. and 6-8 h. These products then tried for deacetylation. in presence of methanolic ammonia at 10°C. Isomeric transformation and blocking and reopening of desired functional

group was carried out successfully. Whatever position of carbon atom in glucosyl ring was blocked in the form of acetyl group was then made available as hydroxyl group for further reactions, that is Phosphorylation of synthetic Nucleosides to convert it in to corresponding synthetic Nucleotides.

References:

1. **B.N. Berad and M. G. Paranjpe:** Annual convention of chemists, Calcutta, Oct. 15, 1984 Paper no. ORG (4).
2. **S. H. Pine:** "Organic Chemistry" McGraw-Hill, New York, International Ed. 5 th Ed. (1979).
3. **I.B. Bedekar, A.R. Joshi and M. G. Paranjpe:** J. Indian Chemical Soc., 58, 1016 (1981).
4. **N. B. Colthup, L. H. Daly and S. E. Wimerley:** Introduction to Infrared and Raman Spectroscopy" Academic Press New York, (1964).
5. **R.M. Silverstein, G.C. Bassler and T.C. Morrill:** Spectroscopic Identification of Organic Compounds" 4th Ed., John Wiley and Sons., INC, New York (1981).
6. **S. A. Barker, E. J. Bourne, r. Stephens and D.H. Whiffen:** J. Chem. Soc., 3468 (1954).
7. **Margareta Avram and GH. D. Mateescu:** "Infrared Spectroscopy by Application in Organic Chemistry" John Wiley and sons., INC, New York (1970).
8. **J.R. Dyer:** Applications of Absorption Spectroscopy of Organic Compounds", Prentice-Hal
9. **Shenyou Nie Wei Li And Biao Yu:** Product Chemistry of Bio organic & Natural Product Chem. J. Am. Chem. Soc. 2014 Mar 19:136(11); 4157-60 doi: 10.1021/ju501460J. Epub. 2014 Mar 11