STATUS OF WATER QUALITY OF KAPILESHWAR DAM, ASHTI, DIST- WARDHA WITH REFERENCE TO PLANKTON DIVERSITY

P. J. Awate¹ and M. R. Kakpure²

¹Department of Zoology, Department of Botany,

²S.M.D. Bharti Mahavidyalaya, Arni Dist-Yavatmal. 445103 (Maharashtra).

mr.prashantawate@gmail.com

Abstract

The present study objectively conducted to analysis the physico-chemical parameters and planktonic diversity of Kapileshwar dam. The study area was divided into five sampling stations to cover the whole dam area comprehensively during the June 2013 to July 2014. The physico-chemical parameters were observed Temperature, Transparency, pH, Turbidity, Total Dissolved Solid, Total alkalinity DO, Total hardness, Nitrate Sulphate and Phosphate during the study. Water temperature exhibited feeble positive correlation with zooplankton. Transparency played an important role in flourishing population of phytoplankton as it established a positive correlation at 1% level of significance ($r = 0.729$). The dissolved oxygen in water of the dam ranged from 5.00mg/l to 9.10mg/l at station I and II characterized as good level of water body. Nutrients like nitrates and phosphate present in the water body were found in the permissible limits as prescribed by WHO, BIS for drinking water quality which confirm that the water is unpolluted and safe for drinking purposes. At present the Kapileshwar dam is of mesotrophic nature.

Keywords: Physico-chemical parameters, Kapileshwar dam, Plankton and Mesotrophic.

Introduction

The water of Kapileshwar dam is utilized for drinking, irrigation and for fisheries activity. It is necessary that the quality of water should be checked periodically because due to use of contaminated water biotic fauna may suffer from a variety of water born diseases Sunkund and Patil (2004). In recent times limnological studies are often related to fisheries, biodiversity conservation, pollution assessment and rejuvenation of degraded inland freshwater ecosystems (Arlinghaus *et al.*, 2008). Zooplankton diversity is one of the most important ecological parameters in water quality assessment. In many areas, the ecological impacts from human activities will far exceed the impacts from climate change, Scholze *et al.*, (2006) have worked on a climate-change risk analysis for world ecosystems, Islam (2007) in a pond of Rajshahi University, has investigated the effects of abiotic parameters on the variations of zooplankton population. Zooplankton is good indicator of the changes in water quality because they are strongly affected by environmental conditions and respond. Nelson *et al.*, (2009) carried out a detailed study on the combined effects of urbanization and climate change on stream ecosystems: from impacts to management options. Similarly, Durance and Ormerod (2009) have investigated existing trends in water quality and the consequent discharge confound long-term warming effects on river macroinvertebrates. The study of plankton and physico-chemical

property of water was carried out by various people like Mishra and Saksena (1991), Kuashik and Saksena (1995). Patel *et al.*, (2013), Vasanthkumar and Vijay Kumar (2011), Garg *et al.*, (2010). The present study was aimed to evaluate water quality and occurrence of zooplankton and phytoplankton with reference to physico-chemical parameters in a Kapileshwar dam.

Material and Methods

The Kapileshwar dam is a man made reservoir constructed on a local nalla. It is at 78°-17'-00" E longitude and 20°-12'-00"N. The length of dam is 589 mtrs. and 7.31 meters high with a catchment area of 6.68 square km situated near Ashti Tahsil of Wardha District. Gross storage capacity of the reservoir is about 1.720 Mm³. The study area was divided into 5 sampling sites. Water samples were collected regularly on monthly basis in between 8 a.m. to 10 a.m. Water pH, DO, and TDS observed at sampling site using water analysis kit (Systronic make) while other abiotic components were analyzed at laboratory condition using the method prescribed by APHA (1989).

Result and Discussion

It is fact that maintenance of healthy aquatic ecosystem is dependent on the physico-chemical properties of water. Station wise mean values of physico-chemical parameters are summarized in Table 1.

Water Temperature (WT)

Temperature is one of the most important ecological factors, which controls the physiological behavior and distribution of organisms. In present study, lowest value of water temperature found in January 19°C and highest in Jun 31.10°C with the mean values 26.11°C. that shows the optimum ranged for growth of aquatic fauna and flora. This result agrees with the reported by Swaranlatha and Rai (1998) in Banjara Lake. Salve (2006) also reported similar trends in Wan Prkalp reservoir, Nagpur.

pH

pH regulated most of biological processes and bio-chemical reactions. pH value ranged from minimum 6.05 to maximum 7.48 at station-I. The pH of water tends towards alkaline nature. Minimum pH was found in the month of April (summer) and reached to the maximum in the month of December (winter). This is in agreement with the findings of Manjare *et al.*, (2010) who studied the Vadgaon tank of Kolhapur. Mean and range pH value of all the five stations observed was 6.98 ± 0.10 .

Transparency

Fresh water body in the present investigation showed no typical trend. Average mean value of water transparency evaluated is 47.68 cm. The lower values of transparency during rainy season may be attributed due to the rains which might have brought silt and mud from the catchment area making the water turbid. This is in the conformation with the observation made by Zafar (1966); Kaur *et al.*, (2008). Mwara, (2006) also observed low transparency during rainy season in manmade reservoirs in Kenya. It was found higher in the summer season which might be due to suspended particles accumulated during summer months or may be attributed to human activity like washing and bathing. Jaybhaye (2009) also reported similar trends in Parola dam of Hingoli district.

Total dissolved solids (TDS)

TDS is measure of all the dissolved substances, both organic and inorganic in water. Higher TDS attributes to high dissolved and suspended particles in to the water. At station-I the total dissolved solid was found higher 400mg/l with the mean value considering all five station was 315.71 mg/l. It is in the range of permissible limit. Similar results were also reported by Chaturbhuj *et al.*, (2004) in the Jamwa Ramgarh Lake, Jaipur.

Turbidity

The water was less turbid as observed during study period. Monsoon months particularly showed turbid water which is attributed to the surface runoff in the rainy season from the

catchment area; recently Manjare *et al.*, (2010) have reported higher turbidity in summer season. Average mean turbidity values of the entire sampling site calculated to 39.95 NTU.

Total alkalinity

Total alkalinity of water is the quality of water and kinds of components present in water such as bicarbonate, carbonate and hydroxide. Total alkalinity was in the ranged from 106mg/l to 330mg/l at sampling station –II and sampling station-V (Table-1) respectively. The maximum value recorded in the month of January (winter) and minimum in the month of August (monsoon). The alkaline water was found productive. Spence (1967) classified the lake into three categories based on alkalinity. On the basis of this classification, Kapileshwar dam considered as a nutrient rich dam.

Dissolved oxygen (DO)

Dissolved oxygen is an important limnological parameter indicating level of water quality, organic production and reflects the physical, biological processes prevailing in the dam water. It ranged was 5.00mg/l to 9.10 mg/l. The level of DO was found less during the month of May (summer). This is because of the low solubility of gases at high temperature Hynes, (1978); similarly during summer water volume also decreased and became more concentrated with the pollutants, Decreased DO in summer correlates with the higher solubility of oxygen at lower temperature. This is in agreement with the recent findings of Garg *et al.*, (2010) who found that dissolved oxygen was less during summer season. Average DO in the present study also exceeded the limit of 5mg/l as per European Environmental Commission Chapman (1997) and mean value observed in the present investigation was 7.91 mg/l. Therefore, it can be concluded that water is safe for human consumption.

Total hardness

The hardness of water is mainly governed by the content of calcium and magnesium which largely combine with bicarbonates & carbonates (temporary hardness) and with sulphate, chlorides and other anions of minerals (permanent hardness). In the present study, mean value of total hardness of five different stations was observed as 127.17 mg/l. The minimum and maximum evaluated was 86 mg/l and 226 mg/l respectively. It was found in permissible limit.

Nitrates

Nitrate is basic nutrient, which is determined the productivity of lake. In the fresh water nitrate content is meager. In the present study the average values of nitrate of all the sampling stations were observed to be 0.48 mg/l. there

was no seasonal trend in the concentration of nitrates and minimum concentration was found 0.19mg/l at station-I while maximum 0.87 mg/l at station-IV. Nitrates were estimated more in winter season. This is in agreement with the findings of Islam(2007).

Sulphate

Most of the sulphate ions are probably derived from the solution of calcium & magnesium ions. Sulphate is naturally occurring ion found in all types of water and wide ranges in nature. Occurrence of sulphate in water is due to influx of runoff and leaching process. The average mean of water sulphate content was found to be

12.46 mg/l which is in permissible limit. Bhagat (2008) have also observed minimum sulphate content in Ambadi dam, near Akot Dt. Akola.

Phosphate

The lower and higher values fluctuated from 0.02 mg/l to 0.60 mg/l at station- V and station- II respectively. Average mean of all the five stations calculated 0.25 mg/l. Winter months contribute higher concentration as compared to monsoon and summer seasons. Higher values may be due to accumulation of surface agricultural runoff and washing activities that contributed to the inorganic phosphate contents.

Table 1 : Showing station wise mean values of Kapileshwar dam during June 2013 to July 2014

Parameters	Sampling stations				
	I	II	III	IV	V
Water temperature(°c)	25.82	25.36	25.66	26.11	25.76
Transparency(cm)	43.95	70.64	61.29	27.11	25.76
Total Dissolved Solid (mg/l)	251.43	264.29	260.00	315.71	282.86
pH	6.95	6.93	6.82	6.98	7.05
Turbidity(NTU)	41.57	32.40	38.17	44.40	43.20
Total alkalinity(mg/l)	153.00	163.57	158.14	185.14	208.29
DO (mg/l)	7.41	6.47	6.20	7.91	7.12
Total Hardness(mg/l)	128.57	128.57	126.29	117.43	135.00
Nitrate (mg/l)	0.43	0.45	0.49	0.49	0.52
Sulphate (mg/l)	9.68	9.27	13.69	19.89	9.75
Phosphate (mg/l)	0.31	0.29	0.16	0.32	0.18

- S-1:-Sampling station 1
- S-2:-Sampling station 2
- S-3:-Sampling station 3
- S-4:-Sampling station 4
- S-5:-Sampling station 5

Figure:-Map of Study area (Kapileshwar Dam of Ashti)

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure (1-6):Graphical representations of mean of physico-chemical parameters of Kapileshwar dam during June 2013 to July 2014.

Conclusion

The pollution indicators phytoplankton and zooplankton were less in number at all the stations similarly As no industrialization has taken place in the surrounding areas, menace of effluent discharge and subsequent pollution is not evident in the dam. Abiotic components are found in the safe limit which confirms that the water is safe for drinking and also for healthy fish culture. The present status of water body is mesotrophic and unpolluted hence, the water can be utilized for irrigation, drinking and fishery activities.

References

- APHA (1989) Standard methods for Examination of water and waste water, 2nd Edn.
- Arlinghaus, R., Johnson, B.M., and Wolter, C. (2008) The past, present and future role of limnology in freshwater fisheries science.. *Int. Rev. of Hydrobio.*, 93:541-549.
- Bhagat, V.B. (2008) Planktonic biodiversity of Ambadi Dam, near Akot Dt. Akola. (Ph.D thesis)
- Chapman D (1997) *Water quality assessment*, 2nd Edn published by E&FN Spon. Chapman & Hall, 2-6 Boundary, Row London, U.K.
- Chaturbhui, M., Sisodia, R., Kulshreshtha, M., and Bhatia, A.L., (2004) A Case study of the Jamwa ramgarh wetland with special reference to physico-chemical properties of water and its environs. Department of Zoology, University of Rajasthan, Jaipur, India, *J. of Hydrobio*, 16.
- Durance Land Ormerod S.J. (2009) Trends in water quality and discharge confound long-term warming effects on river macro-invertebrates.. *Freshwater Bio.*, 54:388-405.
- Garg, R.K. Rao, R.J. and Uchchariya, D., Shukla, G. and Saksena D.N. (2010) Seasonal variation in water quality and major threats to Ramsagar reservoir *India.. Afr. J. Env. Sci. and Tech.*, Vol.4(2):061-076.
- Hynes H.B. (1978) *The biology of polluted water*, Liverpool Uni. Press, Liverpool, 200-204.
- Islam S. N. (2007) Physico-chemical condition and occurrence of some zooplankton in a pond of Rajshahi Uni. Res. *J. Fish and Hydrobio.*, 2(2):21-25.
- Jaybhaye, V.M. (2009) Studies on Physico-chemical parameters of Parol Dam in Hingoli District (MS), Shodh, Samiksha & Mulyankan *Int. Res. J.*, Vol.2(9-10):44-46.
- Kaur, R., Minhas, P.S., Jain, P.C., Singh, P. and Dubey, D.S. (2008) Geo-spatial analysis of land-water resource degradation in two economically contrasting agricultural regions adjoining national capital territory (Delhi). *Environmental Monitoring & Assessment*, doi:10.1007/s10661-008-0378-3.
- Kaushik, S. and Saksena, D.N. (1995) Trophic status and rotifer fauna of certain water bodies in central India. *J. Env. Bio.* 16:283-291.
- Manjare, S.A., Vhanalkar, S.A. and D.V. Mulev (2010) Water quality assessment of Vadgaon tank of Kolhapur with special reference to zooplankton, *Int. J. Advance Biotech & Res.* Vol.1(2), 91-95.
- Mishra, S.R. and Saksena, D.N. (1991) Pollutional ecology with reference to physicochemical characteristics of Morar (Kalpi) river, Gwalior (M.P.). In: Nalin K. Shastree (Ed.). *Current trends in limnology*. Narendra Publishing.
- Mwara, F. (2006) Some aspect of water quality characteristics in small shallow tropical man made reservoirs in Kenya. *Afr. J. Sci. and Tech.*, 7 (1): 82-96.
- Nelson K.C., Palmer M.A., Pizzuto J.E. (2009) Forecasting the combined effects of urbanization and climate change on stream ecosystems: from management options, *J. Applied Ecol.*, 154-163.
- Patel, V., Shukla, S.N. and Pandey, U. (2013) Studies on primary productivity with special Reference to their physicochemical status of Govindgarh Lake Rewa (M.P.), India. *Int. J. Sci. Res.*, 2(11):508-510.
- Salve, B.S. and Hiware, C.J. (2006) Studies on water quality of Wan prakalp reservoir, Nagpur near Parali Vajnath Dt-Beed *J. Aqua. Biol.*, 21(2):113-117.
- Scholze, M., W. Knorr, N. Amell and I.C. Prentice (2006) A climate- change risk analysis for world ecosystems: PIVAS 103, 13116-13120.
- Spence, D.H.N. (1967) The zonation of plants in fresh water lakes. *Adv. Ecol. Res.*, 12:37-125.
- Sunkad B.N. and Patil H. S. (2004) Water quality assessment of fort of Belgaun (Karnataka) with special reference to zooplankton, *J. Envi. Biol.* 25(1), 99-102.
- Swaranlatha, N., and Narsing Rai, A., (1998) Ecological studies of Baniara Lake with reference to water pollution. *J. Env. Bio.*, 19(2):179-186.
- Vasanth kumar, B. and Vijaykumar, K. (2011) Diurnal Variation of physico-chemical properties and primary productivity of phytoplankton in Bheema River. *Recent Res. in Sci. and Tech.*, 3(4): 39-42.
- Zafar, A.R. (1966) Limnology of Hussian sagar Lake, Hyderabad. *India phykas*, 5; 115-126.