18. DEEP SEA DRILLING PROJECT, LEG 4 FORAMINIFERA FROM SELECTED SAMPLES

W. H. Blow, British Petroleum Research Center, London, England

BIOSTRATIGRAPHIC SUMMARY OF AVAILABLE SAMPLES

Hole 23 (lat 6° 08.75' S., long 31° 02.6' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	0 to 9.14	Sections 1, 2 and 4: Zone N. 22; Pleistocene.
3	62.4 to 71.6	Sections 2 and 3: No determinations.
4	112.0 to 121.0	Section 1: No determination. Sections 3 and 4: Upper part Zone N. 4; Early Miocene.
5	121.0 to 131.0	Section 1: Upper part Zone N. 4; Early Miocene.

Hole 24 (lat 6° 16.74' S., long 30° 53.18' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	198.0 to 207.0	Section 1: Zone N. 4; Early Miocene.
4	225.0 to 234.0	Sections 2, 3 and 4: No determinations.

Hole 24A (lat 6° 16.74' S., long 30° 53.18' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	503.0 to 511.0	Section 1: No determination.
2	511.0 to 520.0	Section 1: No determination.
3	540.0 to 550.0	Section 1: No determination.
4	550.0 to 558.0	Sections 1 and 2: No determinations.

Hole 25 (lat 0° 31.00' S., long 39° 14.40' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	0 to 9.0	Sections 1, 2, 3, 4, 5 and 6: Zone N. 22; Pleistocene.
2	9.0 to 18.0	Sections 1, 2 and 3: Zone N. 22; Pleistocene.
3	18.0 to 27.0	Sections 1 and 2 (part): basal Zone N. 19; Pliocene. Section 2 (part): Zone N. 18; Pliocene or latest Miocene. Sections 3, 4, 5 and 6: Within the interval [Zone N. 18 Zone N. 17 upper part] ; Late Miocene.
4	27.0 to 36.0	Section 1: Heterogenous. Sections 2 and 3: Probably Zone N. 17; Late Miocene. Section 4: Within the interval [Zone N. 17 earlier part Zone N. 16] ; Late Miocene.

Hole 26 (lat 10° 53.55' N., long 44° 02.57' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	96.0 to 105.0	Section 1: Zone N. 23; Holocene. Sections 2 and 3: No determinations.
2	105.0 to 114.0	Sections 1 and 2: No determinations.
3	227.0 to 238.0	Sections 1 and 2: No determinations.
4	401.0 to 410.0	Section 1: No determination.
5	477.0 to 483.0	Sections 1, 2, 3 and 4: No determinations.

Hole 27 (lat 15° 51.39' N., long 56° 52.76' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	84.0 to 93.0	Sections 1, 2 and 6: No determinations.
2	141.0 to 150.0	Sections 1, 2 and 3: No determinations.
3	235.0 to 244.0	Section 2: No determination.
4	244.0 to 249.0	Sections 1, 2 and 3: No determinations.
5	369.0 to 378.0	Sections 1, 2, 3 and 4: No determinations.
6	454.0 to 463.0	Sections 2 and 3: No determinations.

Hole 27A (lat 15° 51.39' N., long 56° 52.76' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	25.0 to 34.0	Sections 1, 4, 5 and 6: No determinations.
2	45.0 to 54.0	Section 3: No determination.
3	54.0 to 63.0	Section 1: No determination.
4	63.0 to 72.0	Sections 1, 2, 3, 4, 5 and 6: No determinations.
5	72.0 to 81.0	Sections 1, 2, 3, 4 and 5: No determinations.

Hole 28 (lat 20° 35.19' N., long 65° 37.33' W.)

Core	Depth Below Sea Bed meters	Age Determination
2	68.0 to 77.0	Sections 2, 3 and 4: No determinations.
3	169.0 to 176.0	Sections 1, 2, 3, 4, 5 and 6: No determinations.
5	236.0 to 245.0	Core catcher sample: No determination.

Hole 29 (lat 14° 47.11' N., long 69° 19.36' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	0 to 9.0	Sections 1, 2 and 3: Zone N. 23; Holocene.
2	9.0 to 18.0	Sections 1, 2 and 3: Zone N. 22; Pleistocene.
3	18.0 to 27.0	Section 1: ?Zone N. 21; Late Pliocene.
4	27.0 to 37.0	Sections 1, 2 and 3: Within $\left[\begin{array}{l} \text{Zone N. 21} \\ \text{Zone N. 20} \end{array} \right]$ interval; Pliocene. Section 4: Within $\left[\begin{array}{l} \text{Zone N. 21} \\ \text{Zone N. 19} \end{array} \right]$ interval; Pliocene.
5	37.0 to 46.0	Section 1: No determination.
6	46.0 to 55.0	Section 1: No determination.
7	117.0 to 126.0	Section 1: No determination.
10	144.0 to 153.0	Sections 2, 4 and 5: No determinations.
12	162.0 to 172.0	Section 4: No determination.
15	193.0 to 202.0	Section 4: No determination.
16	202.0 to 211.0	Section 3: No determination.
17	211.0 to 220.0	Section 4: No determination.

Hole 29A (lat 14° 47.11' N., long 69° 19.36' W.)

Core	Depth Below Sea Bed meters	Age Determination
2	50.0 to 59.0	Section 1: No determination.
4	68.0 to 77.0	Section 1: No determination.

Hole 29B (lat 14° 47.11'N., long 69° 19.36' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	57.0 to 66.0	Sections 1, 4, 5 and 6: No determinations.
2	68.0 to 77.0	Section 2: No determination. Section 3: Zone N. 22; Pleistocene. Section 4: ?Zone N. 19; Pliocene.
3	78.0 to 87.0	Sections 1 and 2: No determinations.
4	87.0 to 97.0	Sections 1, 2, 3 and 4: No determinations.
5	97.0 to 106.0	Section 1: Zone N. 8; basal Middle Miocene. Section 2: Zone N. 7 or Zone N. 8. Section 3: Zone N. 7; top early Miocene.
6	106.0 to 115.0	Section 1: Zone N. 7; top Early Miocene.
8	124.0 to 133.0	Sections 1 and 3: No determinations.
9	218.0 to 227.0	Section 1: No determination.
10	227.0 to 232.0	Section 3: No determination.

Hole 30 (lat 12° 52.92' N., long 63° 23.00' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	50.0 to 59.0	Sections 1, 2, 3 and 5: Zone N. 22; Pleistocene.
2	59.0 to 69.0	Sections 2, 3 and 4: Zone N. 22; Pleistocene.
3	108.0 to 117.0	Sections 1, 2, 3 and 4: Zone N. 22; Pleistocene.
4	117.0 to 126.0	Sections 1 and 2: Within the interval Zone N. 22 early part Zone N. 21 ; Late Pliocene/Early Pleistocene.
5	164.0 to 174.0	Sections 1 and 2: Zone N. 21; Late Pliocene.
6	174.0 to 178.0	Sections 1 and 3: Zone N. 21; Late Pliocene.

Hole 30 (lat 12° 52.92' N., long 63° 23.00' W.) (*Continued*)

Core	Depth Below Sea Bed meters	Age Determination
7	260.0 to 269.0	Section 1: ?basal part Zone N. 21; later Pliocene.
8	318.0 to 327.0	Core catcher sample: Zone N. 19; Pliocene.
9	366.0 to 375.0	Sections 1 and 2: Zone N. 14; Middle Miocene.
10	375.0 to 384.0	Sections 2 and 3: Zone N. 14; Middle Miocene.
11	384.0 to 393.0	Sections 1, 2 and core catcher sample: Zone N. 13; Middle Miocene.
12	393.0 to 402.0	Sections 1, 2, 3 and 4 (part): Zone N. 12; Middle Miocene. Sections 4 (part) and 5: Zone N. 11; Middle Miocene.
13	402.0 to 407.0	Sections 1, 2 and 3: Zone N. 10; Middle Miocene.
14	407.0 to 412.0	Section 1: Zone N. 10; Middle Miocene.
15	412.0 to 421.0	Sections 1, 2 and 3 (part): Zone N. 10; Middle Miocene. Section 3 (part): Zone N. 9; Middle Miocene. Sections 4 and 5: Zone N. 8; lower part Middle Miocene.
16	421.0 to 430.0	Section 1: Zone N. 8; lower part Middle Miocene.

Hole 31 (lat 14° 56.60' N., long 72° 01.63' W.)

Core	Depth Below Sea Bed meters	Age Determination
1	0 to 9.0	Sections 1, 2, 3, 4, 5 and 6: Zone N. 22/N. 23; Pleistocene/Holocene.
3	60.0 to 69.0	Sections 1, 2, 3, 5 and 6: Zone N. 21; Late Pliocene.
4	69.0 to 78.0	Section 1: Within interval [Zone N. 21] [Zone N. 20]; Pliocene.

Core	Depth Below Sea Bed meters	Age Determination
5	78.0 to 88.0	Section 1: Within interval [Zone N. 21] [Zone N. 20]; Pliocene.
6	88.0 to 97.0	Sections 1 and 2: No determinations.
7	97.0 to 106.0	Section 1: Zone N. 20; Pliocene.
8	155.0 to 164.0	Sections 1 and 2: No determinations.
9	212.0 to 221.0	Sections 2 and 3 (part): Zone N.4; Early Miocene. Section 3 (part): Zone N. 4 or ?Zone N. 3 (= P. 22); basal Miocene or ?topmost Oligocene.
10	270.0 to 279.0	Section 1: Zone N. 4 or ?Zone N. 3; basal Miocene or ?topmost Oligocene. Sections 2, 3, 4 and 5: No determinations.

FAUNAS OF SAMPLES EXAMINED

(Determinations by W. H. Blow)

Hole 23

Sample 4-23-1-1, 82-84 cm:

No significant planktonic foraminifera.

Sample 4-23-1-2, 72-74 cm:

Poor planktonic foraminiferal fauna with *Globorotalia* (*G.*) *truncatulinoides pachytheca*, *G. (G.) cultrata* (s.l.), *G. (T.) acostaensis pseudopima*, *Globigerina eggeri eggeri*, *Globigerinoides ruber* (white), *Pulleniatina obliquiloculata obliquiloculata*.

Age: Zone N. 22; Pleistocene.

Sample 4-23-1-4, 94 cm:

Rich planktonic foraminiferal fauna with *Globorotalia* (*G.*) *truncatulinoides truncatulinoides* (phylogenetically very primitive), *G. (T.) tosaensis tenuithec*, *G. (G.) multicamerata*, *G. (G.) miocenica*, *G. (G.) cultrata cultrata*, *G. (G.) cultrata menardii*, *Globoquadrina altispira altispira*, *G. altispira* cf. *globosa*, *Pulleniatina obliquiloculata obliquiloculata*, *Globigerina eggeri eggeri*, *Globorotalia (G.) cultrata exilis*, *Globigerina calida praecalida*, *G. juvenilis*, *Hastigerina pelagica*, *H. siphonifera siphonifera*, *Globigerinoides fistulosus*, *G. conglobatus conglobatus*, *Sphaeroidinella dehiscens dehiscens*.

Age: Basal Zone N. 22; earliest Pleistocene.

Sample 4-23-3-2, 75-77 cm:

No significant planktonic foraminifera.

Sample 4-23-3-3, 75-76 cm:

No planktonic foraminifera.

Sample 4-23-4-1, 12-13 cm:

No significant planktonic foraminifera.

Sample 4-23-4-3, 75-76 cm:

Poor planktonic foraminiferal fauna with *Globigerinita dissimilis ciperoensis*, *G. incrusta*, *G. stainforthi praestainforthi*, *Globorotalia (T.) kugleri*, *G. (T.) pseudokugleri*, *Globigerinoides quadrilobatus primordius*, *G. quadrilobatus immaturus*.

Age: Upper part Zone N. 4; Aquitanian, Early Miocene.

Sample 4-23-4-4, 75-76 cm:

As above plus *Miogypsina*.

Age: Upper part Zone N. 4; Early Miocene.

Sample 4-23-5-1, 25-27 cm:

Rich planktonic foraminiferal fauna with *G. (T.) obesa*, *Globorotalia (T.) kugleri*, *G. (T.) pseudokugleri*, *G. (T.) siakensis*, *G. (T.)* cf. *scitula praescitula*, *G. (T.) peripheroronda forma* α, *Globigerinita dissimilis* (s.s.), *G. dissimilis ciperoensis*, *G. incrusta*, *Globorotaloides* cf. *suteri*, *Globigerinoides quadrilobatus primordius*, *G. quadrilobatus immaturus*, *G. quadrilobatus* cf.

quadrilobatus, *G. quadrilobatus* cf. *altiaperturus*, *Globigerina praebulloides praebulloides*, *G. juvenilis*, *G. bradyi*, also with *Miogypsina* and *Spiroclypeus*.
Age: Upper part Zone N. 4; Early Miocene.

Hole 24

Sample 4-24-1-1, 34-35 cm:
Globorotalia (T.) *siakensis*, *G. (T.) kugleri*, *G. (T.) pseudokugleri*, *Globigerinoides quadrilobatus primordius*, *Globigerinita dissimilis dissimilis*, *Globorotaloides* cf. *suteri*.

Age: Zone N. 4; Aquitanian, Early Miocene.

Sample 4-24-4-2, 75-76 cm:
No planktonic foraminifera.

Sample 4-24-4-3, 75-76 cm:
As above.

Sample 4-24-4-4, 75-76 cm:
No significant planktonic foraminifera.

Hole 24A

Sample marked "Center Bit":
No significant planktonic foraminifera.

Sample 4-24A-1-1, 5 cm:
No planktonic foraminifera.

Sample 4-24A-2-1, 5.5 cm:
No planktonic foraminifera.

Sample 4-24A-3-1, Core catcher (bottom):
No recognizable planktonic foraminifera.

Sample 4-24A-3-1, Core catcher (middle):
No recognizable planktonic foraminifera.

Sample 4-24A-3-1, Core catcher (top):
As above.

Sample 4-24A-3-1, 79-80 cm:
No significant planktonic foraminifera.

Sample 4-24A-4-1 (?), Core catcher (bottom):
No significant planktonic foraminifera.

Sample 4-24A-4-2, 74-76 cm:
No significant planktonic foraminifera.

Hole 25

Sample 4-25-1-1, 110 cm:
Rich planktonic foraminiferal fauna with *Globorotalia* (*G.*) *cultrata menardii*, *G. (G.) truncatulinoides truncatulinoides*, *Pulleniatina obliquiloculata finalis*, *P. obliquiloculata obliquiloculata*, *Globigerinoides conglobatus conglobatus*, *G. ruber*, *G. quadrilobatus* (s.l.),

Globigerina eggeri eggeri, *Hastigerina siphonifera siphonifera*, *Sphaeroidinella dehiscentis dehiscentis*.

Age: Zone N. 22; Pleistocene.

Sample 4-25-1-2, 110 cm:
As above.

Sample 4-25-1-3, 110 cm:
As above.

Sample 4-25-1-4, 110 cm:
As above.

Sample 4-25-1-5, 130 cm:
Globorotalia (*G.*) *truncatulinoides truncatulinoides*, *G. (G.) cultrata menardii*, *G. (G.) crassula viola*, *G. (Turborotalia) crassaformis ronda*, *G. (T.) acostaensis humerosa*, *G. (T.) scitula scitula*, *Pulleniatina obliquiloculata obliquiloculata*, *P. obliquiloculata finalis*, *Candeina nitida nitida*, *Globigerinoides quadrilobatus sacculifer*, *G. quadrilobatus* (s.s.), *G. conglobatus conglobatus*, *G. ruber*, *Hastigerina pelagica*, *H. siphonifera siphonifera*, *Sphaeroidinella dehiscentis dehiscentis*, *Turborotalita* cf. *humilis*.

Age: Zone N. 22; Pleistocene.

Sample 4-25-1-6, 0.2 cm:
As above.

Sample 4-25-2-1, 0-2 cm:
Globorotalia (*G.*) *cultrata menardii*, *G. (G.)* cf. *cultrata exilis*, *G. (G.)* cf. *tumida* (s.s.), *G. (G.) miocenica*, *G. (G.) unguata*, *G. (Turborotalia) acostaensis pseudopima*, *G. (T.) acostaensis humerosa*, *G. (T.)* cf. *ventriosa*, *G. (T.) tosaensis tenuitheca*, *G. (T.) crassaformis* (s.l.), *Globigerina eggeri eggeri*, *G. calida praecalida*, *Globigerinoides ruber*, *G. conglobatus conglobatus*, *Pulleniatina obliquiloculata obliquiloculata*, *Hastigerina* spp., *Sphaeroidinella dehiscentis dehiscentis*. Phylogenetically primitive *Globorotalia* (*G.*) *truncatulinoides* (s.l.) (partially carinate).

Age: Lower Zone N. 22; earliest Pleistocene.

Sample 4-25-2-2, 0-2 cm:
As above.

Sample 4-25-2-3, 0-2 cm:
As above.

Sample 4-25-2-3, 30 cm:
As above.

Sample 4-25-2-3, 135 cm:
As above.

Sample 4-25-3-1, 150 cm:
Globorotalia (*G.*) *cultrata* (s.l.), *G. (G.) cultrata exilis*, *G. (G.) multicamerata*, *G. (G.) miocenica*, *Globoquadrina dehiscentis dehiscentis*, *G. altispira altispira*, *G. altispira globosa*, *Globigerinoides obliquus extremus*,

Globigerinoides conglobatus conglobatus, *G. elongatus*, *G. quadrilobatus*, *Globigerina nepenthes*, *Sphaeroidinellopsis subdehiscens paenedehiscens* (abundant), *Sphaeroidinella dehiscens dehiscens forma immatura*, *Globorotalia (Turborotalia) acostaensis humerosa*, *G. (T.) acostaensis acostaensis*, *Orbulina* spp., *Hastigerina siphonifera* (s.s.).

Age: Early Zone N. 19; Early Pliocene.

Sample 4-25-3-2, 25 cm:

As above, but only very few *S. dehiscens dehiscens forma immatura* with minute dorsal supplementary apertures.

Age: Basal Zone N. 19; Early Pliocene.

Sample 4-25-3-2, 150 cm:

As above, but no specimens of *Sphaeroidinella dehiscens dehiscens* (including *forma immatura*) observed.

Age: Zone N. 18; Late Miocene/Early Pliocene.

Sample 4-25-3-3, 122-124 cm:

Globorotalia (G.) cultrata limbata, *G. (G.) cf. miocenica*, *Globorotalia (Turborotalia) acostaensis humerosa*, *G. (T.) acostaensis acostaensis*, *Globigerina nepenthes*, *G. eggeri multiloba*, *G. cf. decoraperta*, *Globigerinoides bollii*, *G. obliquus extremus*, *G. quadrilobatus* (s.l.), *G. cf. conglobatus canimarensis*, *Globoquadrina dehiscens dehiscens*, *G. altispira altispira*, *G. altispira cf. globosa*, *Sphaeroidinellopsis subdehiscens paenedehiscens*.

Age: Within interval [Zone N. 18
Zone N. 17 upper part]; Late
Miocene.

Sample 4-25-3-3, 150 cm:

As above, but with *G. (G.) cf. multicamerata*.

Age: Within interval [Zone N. 18
Zone N. 17 upper part]; Late
Miocene.

Sample 4-25-3-6, 90 cm:

Globorotalia (G.) cultrata (s.l.), *G. (G.) cultrata limbata*, *G. (Turborotalia) acostaensis humerosa*, *G. (T.) acostaensis acostaensis*, *G. (T.) cf. subscitula*, *Globoquadrina larmei obesa*, *G. altispira altispira*, *G. altispira cf. globosa*, *Globigerina decoraperta*, *G. nepenthes*, *G. venezuelana*, *Sphaeroidinellopsis seminulina seminulina*, *S. subdehiscens subdehiscens*, *S. subdehiscens paenedehiscens*, *G. (G.) multicamerata*, *Orbulina* spp., *Globigerinoides bollii*, *G. obliquus extremus*, *G. quadrilobatus* (s.l.).

Age: As above.

Sample 4-25-3-6, 150 cm:

As above.

Sample 4-25-4-1, 0-2 cm:

Strongly heterogeneous assemblage with faunal elements from Zones N. 22, N. 21, N. 19, N. 18 all present.

Age: None.

Sample 4-25-4-2, 92 cm:

Globorotalia (G.) cultrata (s.l.), *G. (G.) cultrata limbata*, *Globorotalia (Turborotalia) acostaensis acostaensis*, *Globoquadrina dehiscens dehiscens*, *G. larmei obesa*, *G. altispira altispira*, *G. altispira cf. globosa*, *Globigerina nepenthes*, *G. venezuelana*, *Sphaeroidinellopsis subdehiscens cf. paenedehiscens*, *S. subdehiscens subdehiscens*, *S. seminulina seminulina*, *Globorotalia (T.) subscitula*, *Globigerinoides obliquus extremus*, *G. bollii*, *G. quadrilobatus* (s.l.).

Age: Probably Zone N. 17; Late Miocene.

Sample 4-25-3-4, 88 cm:

As above.

Sample 4-25-3-4, 150 cm:

Globorotalia (G.) cultrata (s.l.), *G. (G.) cultrata limbata*, *Globorotalia (Turborotalia) acostaensis humerosa*, *G. (T.) acostaensis acostaensis*, *Globoquadrina altispira altispira*, *G. altispira cf. globosa*, *G. dehiscens dehiscens*, *Globigerinoides bollii*, *G. obliquus extremus*, *G. elongatus*, *G. quadrilobatus* (s.l.), *Globigerina nepenthes*, *G. venezuelana*, *G. decoraperta*, *Sphaeroidinellopsis subdehiscens paenedehiscens*, *S. subdehiscens subdehiscens*.

Age: Within interval [Zone N. 18
Zone N. 17 upper part]; Late
Miocene.

Sample 4-25-3-5, 90 cm:

As above, but also with *Sphaeroidinellopsis seminulina seminulina*.

Age: As above.

Sample 4-25-3-5, 150 cm:

As above.

Sample 4-25-4-3, 0-2 cm:

Globorotalia (G.) cultrata (s.l.), *G. (G.) cultrata limbata*, *G. (Turborotalia) acostaensis acostaensis*, *G. (T.) scitula scitula*, *G. (T.) subscitula*, *Globigerinoides bollii*, *G. obliquus extremus*, *G. quadrilobatus* (s.l.), *Globoquadrina altispira altispira*, *G. dehiscens dehiscens*, *G. larmei obesa*, *Hastigerina siphonifera siphonifera*, *Sphaeroidinellopsis seminulina* (s.l.), *S. subdehiscens subdehiscens*, *S. subdehiscens cf. paenedehiscens*, *Globigerina nepenthes*, *G. juvenilis*, *G. bradyi*, *G. venezuelana*, *Globorotaloides hexagona variabilis*.

Age: Probably Zone N. 17; Late Miocene.

Sample 4-25-4-3, 90 cm:

As above.

Sample 4-25-4-4, 0-2 cm:

As above, but with no specimens referable to *Sphaeroidinellopsis subdehiscens cf. paenedehiscens*. The sample shows a little contamination from Pleistocene/Holocene horizons.

Age: Within interval [Zone N. 17 earlier part]
Zone N. 16]; Late
Miocene.

Sample 4-25-4-4, 90 cm:
As for Sample 4-25-4-4, 0-2 cm.
Age: Within interval $\left[\begin{array}{l} \text{Zone N. 17 earlier part} \\ \text{Zone N. 16} \end{array} \right]$; Late
Miocene.

Hole 26

Sample 4-26-1-1, 148-150 cm:
Globorotalia (G.) tumida tumida, *G. (G.) cultrata cultrata*, *G. (G.) cultrata menardii*, *Pulleniatina obliquiloculata finalis*, *Globigerinoides ruber* (pink), *G. conglobatus conglobatus*, *G. quadrilobatus sacculifer*, *Globigerina eggeri eggeri*, *Sphaeroidinella dehiscent dehiscent*, *S. dehiscent excavata*, etc.
Age: Zone N. 23; Holocene.

Sample 4-26-1-2, 75-76 cm:
No significant fauna.
Age: None.

Sample 4-26-1-3, 75-76 cm:
As above.

Sample 4-26-2, Core catcher:
As above.

Sample 4-26-2-1, 124-125 cm:
As above.

Sample 4-26-2-2, 75-76 cm:
As above.

Sample 4-26-3-1, 75-76 cm:
As above.

Sample 4-26-3-1, 148-150 cm:
No significant fauna.

Sample 4-26-3-2, 75-76 cm:
As above.

Sample 4-26-4, Core catcher:
Quartz sand and much authigenic feldspar residue, but no significant fauna.

Sample 4-26-4-1, 150 cm:
As above.

Sample 4-26-5-1, 75-76 cm:
No significant fauna.

Sample 4-26-5-2, 75-76 cm:
As above.

Sample 4-26-5-3, 75-76 cm:
As above.

Sample 4-26-5-4, 75-77 cm:
As above.

Hole 27

Sample 4-27-1-1, 75-76 cm:
No significant fauna.

Sample 4-27-1-2, 62-63 cm:
As above.

Sample 4-27-1-6, 79-80 cm:
As above.

Sample 4-27-2-1, 75-76 cm:
No recognizable planktonic foraminiferal taxa.

Sample 4-27-2-2, 75-76 cm:
No significant planktonic foraminiferal taxa.

Sample 4-27-2-3, 75-76 cm:
As above.

Sample 4-27-3-2, 75-76 cm:
No planktonic foraminifera.

Sample 4-27-4-1, 75-76 cm:
As above.

Sample 4-27-4-2, 75-76 cm:
As above.

Sample 4-27-4-3, 75-76 cm:
As above.

Sample 4-27-5-1, 75-76 cm:
As above.

Sample 4-27-5-2, 75-76 cm:
As above.

Sample 4-27-5-3, 75-76 cm:
As above.

Sample 4-27-5-4, 75-76 cm:
As above.

Sample 4-27-6-2, 75-76 cm:
No planktonic foraminiferal fauna, *Radiolaria* present.
Age: No planktonic foraminiferal determination.

Sample 4-27-6-3, 72-74 cm:
As above.

Hole 27A

Sample 4-27A-1-1, 75-76 cm:
No recognizable planktonic foraminifera.

Sample 4-27A-1-4, 75-76 cm:
As above.

Sample 4-27A-1-5, 75-76 cm:
As above.

Sample 4-27A-1-6, 75-76 cm:
As above.

Sample 4-27A-2-3, 75-76 cm:
As above.

Sample 4-27A-3-1, 75-76 cm:
As above.

Sample 4-27A-4-1, 75-76 cm:
As above.

Sample 4-27A-4-2, 75-76 cm:
As above (residue contains authigenic feldspar).

Sample 4-27A-4-3, 67-68 cm:
As above.

Sample 4-27A-4-4, 73-74 cm:
As above.

Sample 4-27A-4-5, 75-76 cm:
As above.

Sample 4-27A-4-6, 75-76 cm:
As above.

Sample 4-27A-5-1, 41-42 cm:
No planktonic foraminifera.

Sample 4-27A-5-2, 75-76 cm:
As above.

Sample 4-27A-5-3, 75-77 cm:
As above.

Sample 4-27A-5-4, 75-76 cm:
As above.

Sample 4-27A-5-5, 75-76 cm:
As above.

Hole 28

Sample 4-28-2-2, 75-76 cm:
No planktonic foraminifera.

Sample 4-28-2-3, 75-76 cm:
As above.

Sample 4-28-2-4, 75-76 cm:
As above.

Sample 4-28-3-1, 75-76 cm:
No planktonic foraminifera (*Radiolaria* only).
Age: None based on the Globigerinacea.

Sample 4-28-3-2, 75-77 cm:
As above.
Age: None based on the Globigerinacea.

Sample 4-28-3-3, 74-76 cm:
As above.
Age: None based on the Globigerinacea.

Sample 4-28-3-4, 74-76 cm:
As above.
Age: None based on the Globigerinacea.

Sample 4-28-3-5, 74-76 cm:
As above.
Age: None based on the Globigerinacea.

Sample 4-28-3-6, 71-73 cm:
As above.
Age: None based on the Globigerinacea.

Sample 4-28-5, Core catcher:
No planktonic foraminifera.

Hole 29

Sample 4-29-1-1, 75-77 cm:
Much comminuted planktonic foraminiferal fauna, but with *Sphaeroidinella dehiscentis dehiscentis*, *Globigerina eggeri eggeri*, *Globorotalia (Turborotalia) acostaensis* (s.l.), *Pulleniatina obliquiloculata obliquiloculata*.
Age: Probably Zone N. 22/N. 23; Pleistocene/Holocene.

Sample 4-29-1-2, 80-82 cm:
As above.

Sample 4-29-1-3, 68-70 cm:
Rich planktonic foraminiferal fauna with *Pulleniatina obliquiloculata finalis*, *P. obliquiloculata obliquiloculata*, *Sphaeroidinella dehiscentis dehiscentis*, *S. dehiscentis excavata*, *Globorotalia (G.) tumida tumida*, *G. (G.) cultrata* (s.l.), *G. (G.) unguolata*, *G. (G.) fimbriata*, *G. (T.) acostaensis humerosa*, *Globigerina eggeri eggeri*, *Globigerinoides conglobatus conglobatus*, *G. elongatus*, *G. ruber*, *G. quadrilobatus* (s.l.), etc.
Age: Zone N. 23; Late Pleistocene/Holocene.

Sample 4-29-2-1, 72-74 cm:
Sphaeroidinella dehiscentis dehiscentis, *Pulleniatina obliquiloculata finalis*, *P. obliquiloculata obliquiloculata*, *Globorotalia (G.) crassula viola*, *G. (G.) crassula crassula*, *G. (G.) crassula cf. conomiozea*, *G. (T.) crassaformis crassaformis*, *G. (T.) acostaensis cf. pseudopima*, *Globigerina eggeri eggeri*, *Hastigerina siphonifera siphonifera*, *Globigerinoides conglobatus conglobatus*, *G. elongatus*, *G. ruber*, etc.
Age: Zone N. 22; Pleistocene.

Sample 4-29-2-2, 74-76 cm:
Globorotalia (G.) truncatulinoides truncatulinoides, *G. (G.) crassula crassula*, *G. (G.) crassula viola*, *G. (T.) acostaensis* (s.l.), *Globigerinoides conglobatus conglobatus*, *G. elongatus*, *G. ruber*, *G. quadrilobatus* (s.l.), *Globigerina eggeri eggeri*, *Sphaeroidinella dehiscentis dehiscentis*, *Pulleniatina obliquiloculata cf. finalis*, *P. obliquiloculata obliquiloculata*, *Hastigerina siphonifera siphonifera*, *Globorotalia (T.) tosaensis tenuitheca*.
Age: Zone N. 22; Pleistocene.

Sample 4-29-2-3, 75-77 cm:

Much comminuted planktonic foraminiferal fauna.
Age: Probably Zone N. 22; Pleistocene.

Sample 4-29-3-1, 40-42 cm:

Globorotalia (G.) *miocenica*, G. (G.) *cultrata exilis*, G. (G.) *crassula crassula*, *Globigerinoides fistulosus*, G. *conglobatus conglobatus*, G. *elongatus*, G. *quadrilobatus*, *Globigerina eggeri eggeri*, *Candeina nitida nitida*, *Hastigerina siphonifera siphonifera*, *Globorotalia* (T.) *crassaformis ronda*, G. (T.) *acostaensis pseudopima*, *Sphaeroidinella dehiscens dehiscens*.

Age: Probably Zone N. 21; Late Pliocene.

Sample 4-29-4-1, 93-95 cm:

Fauna much comminuted, but with *Globorotalia* (G.) *miocenica*, G. (T.) *acostaensis pseudopima*, G. (T.) *acostaensis humerosa*, *Globigerinoides obliquus extremus*, G. *elongatus*, G. (G.) *cultrata exilis*, G. (T.) *crassaformis ronda*, *Globigerinoides ruber*, G. *quadrilobatus* (s.l.), *Hastigerina siphonifera siphonifera*, *Sphaeroidinella dehiscens dehiscens*.

Age: Within $\left[\begin{array}{c} \text{Zone N. 21} \\ \text{Zone N. 20} \end{array} \right]$ interval; Pliocene.

Sample 4-29-4-2, 65-67 cm:

Globorotalia (G.) *miocenica*, G. (G.) *crassula viola*, G. (T.) *crassaformis ronda*, G. (T.) *acostaensis acostaensis*, G. (T.) *acostaensis humerosa*, *Globigerinoides quadrilobatus fistulosus*, G. *quadrilobatus sacculifer*, G. *ruber*, G. *obliquus extremus*, G. *conglobatus conglobatus*, *Orbulina* sp., *Sphaeroidinella dehiscens dehiscens*.

Age: Probably within $\left[\begin{array}{c} \text{Zone N. 21} \\ \text{Zone N. 20} \end{array} \right]$ interval; Pliocene.

Sample 4-29-4-3, 74-76 cm:

Globorotalia (G.) *multicamerata*, G. (G.) *miocenica*, G. (G.) *crassula crassula*, G. (G.) *crassula* cf. *viola*, G. (T.) *acostaensis acostaensis*, G. (T.) *acostaensis humerosa*, G. (T.) cf. *acostaensis pseudopima*, *Globigerinoides conglobatus conglobatus*, G. *elongatus*, G. *quadrilobatus* (s.l.), G. *ruber*, G. *bollii*, *Globigerina falconensis*, G. *eggeri eggeri*, *Sphaeroidinella dehiscens dehiscens*.

Age: Within interval $\left[\begin{array}{c} \text{Zone N. 21} \\ \text{Zone N. 20} \end{array} \right]$; Pliocene.

Sample 4-29-4-4, 74-76 cm:

Poor planktonic fauna, *Globigerina decoraperta*, *Globigerinoides bollii*, G. cf. *ruber*, G. *quadrilobatus* (s.l.), *Sphaeroidinella dehiscens dehiscens*, *Globorotalia* (G.) *multicamerata*, G. (T.) *acostaensis* (s.l.).

Age: Within interval $\left[\begin{array}{c} \text{Zone N. 21} \\ \text{Zone N. 19} \end{array} \right]$; Pliocene.

Sample 4-29-5-1, 74-76 cm:

No planktonic foraminiferal fauna.

Sample 4-29-6-1, 27-29 cm:

As above.

Sample 4-29-7-1, 105-107 cm:

No significant planktonic foraminifera.

Sample 4-29-10-2, 73-75 cm:

Rich radiolarian assemblage, but no planktonic foraminifera.

Age: None based on the Globigerinacea.

Sample 4-29-10-4, 76-78 cm:

As above.

Sample 4-29-10-5, 74-76 cm:

As above.

Sample 4-29-12-4, 74-76 cm:

As above.

Sample 4-29-15-4, 110-112 cm:

As above.

Sample 4-29-16-3, 90-92 cm:

As above.

Sample 4-29-17-4, 92-94 cm:

As above.

Hole 29A

Sample 4-29A-2-1, 80-82 cm:

No planktonic foraminifera.

Sample 4-29A-4-1, 83-85 cm:

No planktonic foraminifera.

Hole 29B

Sample 4-29B-1-1, 75-77 cm:

No planktonic foraminifera.

Age: None.

Sample 4-29B-1-4, 75-77 cm:

As above.

Sample 4-29B-1-5, 76-78 cm:

As above.

Sample 4-29B-1-6, 74-76 cm:

As above.

Sample 4-29B-2-2, 75-77 cm:

As above.

Sample 4-29B-2-3, 75-77 cm:

Rich planktonic foraminiferal fauna with *Globorotalia* (G.) *truncatulinoidea truncatulinoidea*, G. (G.) *truncatulinoidea pachytheca*, G. (G.) *cultrata menardii*, G. (G.) *ungulata*, *Globigerinoides conglobatus conglobatus*, G. *ruber*, G. *quadrilobatus* (s.l.), *Globigerina eggeri eggeri*, *Pulleniatina obliquiloculata obliquiloculata*, P. *obliquiloculata* cf. *finalis*, *Sphaeroidinella dehiscens dehiscens*.

Age: Zone N. 22; Pleistocene.

Sample 4-29B-2-4, 74-76 cm:

Fairly rich planktonic foraminiferal fauna, but mainly comminuted and with micronodules of manganese, some of which replace foraminifera. *Sphaeroidinella dehiscens dehiscens*, *S. dehiscens dehiscens forma immatura*, *Globigerina nepenthes*, *Sphaeroidinellopsis seminulina* (s.l.), *Globorotalia* (G.) cf. *miocenica*, *G. (G.) cultrata* (s.l.), *G. (T.) acostaensis acostaensis*.

Age: Probably Zone N. 19; Pliocene.

Sample 4-29B-3-1, 74-76 cm:

No significant planktonic foraminifera.

Sample 4-29B-3-2, 74-76 cm:

No planktonic foraminifera.

Sample 4-29B-4-1, 74-76 cm:

As above.

Sample 4-29B-4-2, 74-76 cm:

As above.

Sample 4-29B-4-3, 74-76 cm:

As above.

Sample 4-29B-4-4, 75-77 cm:

No planktonic foraminifera—some micronodules of manganese.

Sample 4-29B-5-1, 70-72 cm:

Globigerinoides cf. *sicanus* (phylogenetically rather primitive), *G. subquadratus*, *G. quadrilobatus trilobus*, *Globoquadrina altispira altispira*, *G. altispira* cf. *globosa*, *G. cf. baroemouensis*, *Globigerinita incrusta*, *Globorotalia* (T.) *siakensis*, *G. (T.) peripheroronda*, *G. (T.) scitula praescitula*.

Age: Zone N. 8 (probably lower part); basal Middle Miocene.

Sample 4-29B-5-2, 74-76 cm:

As above, but no specimens referable to *G. sicanus*.

Age: Zone N. 7 or early Zone N. 8; latest Early Miocene or basal Middle Miocene.

Sample 4-29B-5-3, 74-76 cm:

Globorotalia (T.) *peripheroronda*, *G. (T.) siakensis*, *G. (T.) scitula praescitula*, *Globoquadrina dehiscens advena*, *G. altispira altispira*, *G. baroemouensis*, *Sphaeroidinellopsis seminulina seminulina* (primitive), *Globigerinita incrusta*, *Globigerina juvenilis*, *G. bradyi*, *G. praebuloides praebuloides*, *Globigerinoides subquadratus*, *G. quadrilobatus* (s.l.).

Age: Zone N. 7; Early Miocene.

Sample 4-29B-6-1, 134-136 cm:

Globigerinatella insueta, *Globorotalia* (T.) *peripheroronda*, *G. (T.) scitula praescitula*, *G. (T.) siakensis*, *Globoquadrina dehiscens dehiscens*, *G. dehiscens advena*,

G. altispira altispira, *G. altispira* cf. *globosa*, *G. baroemouensis*, *Globigerina venezuelana*, *Globigerinoides quadrilobatus* (s.l.), *G. subquadratus*.

Age: Zone N. 7; Early Miocene.

Sample 4-29B-8-1, 75-77 cm:

Rich radiolarian assemblage, no planktonic foraminifera.

Age: None based on the Globigerinacea.

Sample 4-29B-8-3, 75-77 cm:

As above.

Sample 4-29B-9-5, 111-113 cm:

As above.

Sample 4-29B-10-3, 103-105 cm:

As above.

Sample 4-29B-13-4, 75-77 cm:

This core was recorded, in the preliminary report as being referable to the Eocene part of the succession on the presence of Eocene Radiolaria and nannoplankton in the cores above this one. However, this sample yields specimens referable to *Globigerinoides diminutus*, *G. quadrilobatus trilobus*, *Sphaeroidinellopsis seminulina seminulina* (phylogenetically very primitive), *Globigerinita incrusta*, *Globorotalia* (Turborotalia) *siakensis*, *G. (T.) peripheroronda* and *Globigerina juvenilis*. This fauna is determinative for a horizon within the interval [Zone N. 8 (lowermost part)] and is, therefore, of the Zone N. 7

same age as seen in Cores 5 and 6 recorded above. This fauna may therefore represent contamination from higher in the succession. The stratigraphic determination for this sample should not be accepted without further investigation. The sample yielded euhedral magnetite and authigenic plagioclase feldspar (euhedral crystals) which are not in agreement with the chert horizons recorded on shipboard. It would seem likely, therefore, that the sample is contaminated from the drilling-bit.

Hole 30

Sample 4-30-1-1, 104-106 cm:

Globorotalia (G.) *truncatulinoides truncatulinoides*, *G. (G.) truncatulinoides pachytheca*, *G. (G.) cultrata menardii*, *G. (T.) crassaformis ronda*, *G. (T.) inflata*, *G. (T.) pseudopima*, *Globigerina eggeri eggeri*, *Globigerinoides elongatus*, *G. ruber*, *G. quadrilobatus* (s.l.), *Pulleniatina obliquiloculata obliquiloculata*, *P. obliquiloculata* cf. *finalis*, *Globigerinoides conglobatus conglobatus*.

Age: Zone N. 22; Pleistocene.

Sample 4-30-1-2, 30 cm:

As above.

Age: Zone N. 22; Pleistocene.

Sample 4-30-1-3, 51 cm:

As above.

Age: Zone N. 22; Pleistocene.

Sample 4-30-1-5, 74-76 cm:

As above.

Age: Zone N. 22; Pleistocene.

Sample 4-30-2-2, 48-50 cm:

Globorotalia (G.) *truncatulinoides truncatulinoides*, G. (G.) *cultrata menardii*, G. (T.) *tosaensis tenuitheca*, G. (T.) *crassaformis ronda*, G. (T.) *pseudopima*, G. (T.) *inflata*, *Globigerinoides conglobatus conglobatus*, G. *elongatus*, G. *ruber*, G. *quadrilobatus* (s.l.), *Globigerina eggeri eggeri*, G. *falconensis*, G. *decoraperta*, *Pulleniatina obliquiloculata obliquiloculata*, *Orbulina* spp.

Age: Zone N. 22; Pleistocene.

Sample 4-30-2-3, 63-65 cm:

No planktonic foraminifera.

Sample 4-30-2-4, 74-76 cm:

Same as for Sample 4-30-2-2, 48-50 cm.

Age: Zone N. 22; Pleistocene.

Sample 4-30-3-1, 136 cm:

Globorotalia (G.) *truncatulinoides truncatulinoides*, *Globorotalia* (T.) *tosaensis* (s.l.), G. (T.) *crassaformis* (s.l.), *Globigerinoides conglobatus conglobatus*, G. *ruber*, G. *elongatus*, G. *quadrilobatus* (s.l.), *Globigerina eggeri eggeri*, G. *decoraperta*, *Pulleniatina obliquiloculata obliquiloculata*, *P. obliquiloculata* cf. *praecursor*, *Globorotalia* (G.) *cultrata menardii*, *Sphaeroidinella dehiscent dehiscent*.

Age: Zone N. 22; Pleistocene.

Sample 4-30-3-2, 74-76 cm:

As above, plus G. (G.) *cultrata exilis*.

Age: Zone N. 22; Pleistocene.

Sample 4-30-3-3, 74-76 cm:

As above.

Age: Zone N. 22; Pleistocene.

Sample 4-30-3-4, 78-80 cm:

As above.

Age: Zone N. 22; Pleistocene.

Sample 4-30-4-1, 123-125 cm:

Globorotalia (T.) *crassaformis ronda*, G. (T.) *acostaensis acostaensis*, G. (T.) *acostaensis pseudopima*, *Globorotalia* (G.) *cultrata menardii*, *Pulleniatina obliquiloculata obliquiloculata*, *Globigerina eggeri eggeri*, G. *decoraperta*, *Globigerinoides conglobatus conglobatus*, G. *ruber*, G. *quadrilobatus* (s.l.), *Sphaeroidinella dehiscent dehiscent*, *Orbulina* spp.

Age: Within the interval [Zone N. 22]
[Zone N. 21]

Sample 4-30-4-2, 74-76 cm:

Globorotalia (G.) *cultrata menardii*, G. (G.) *cultrata exilis*, G. (T.) *acostaensis pseudopima*, G. (T.) *acostaensis humerosa*, *Pulleniatina obliquiloculata obliquiloculata*, *Globigerina eggeri eggeri*, G. *falconensis*, *Globigerinoides conglobatus conglobatus*, G. *elongatus*, G. *ruber*, G. *quadrilobatus* (s.l.), G. *obliquus extremus*, *Sphaeroidinella dehiscent dehiscent*, *Orbulina* spp., *Globorotalia* (T.) *crassaformis ronda*, G. (G.) *crassula crassula*.

Age: Within the interval [Zone N. 22]
[Zone N. 21]

Sample 4-30-5-1, 74-76 cm:

Globorotalia (T.) *tosaensis tenuitheca*, G. (T.) *crassaformis ronda*, G. (T.) *crassaformis crassaformis*, G. (T.) *acostaensis pseudopima*, G. (T.) *acostaensis humerosa*, G. (T.) *scitula scitula*, G. (G.) *crassula viola*, G. (G.) *crassula crassula*, *Globigerina falconensis*, G. *eggeri eggeri*, G. *bulloides apertura*, *Globigerinoides conglobatus conglobatus*, G. *ruber*, G. *elongatus*, *Pulleniatina obliquiloculata obliquiloculata*, *Sphaeroidinella dehiscent dehiscent*, *Orbulina* sp.

Age: Zone N. 21; Late Pliocene.

Sample 4-30-5-2, 74-76 cm:

As above.

Sample 4-30-6-1, 110-112 cm:

Globorotalia (G.) *cultrata menardii*, G. (G.) *crassula crassula*, G. (T.) *pseudopima*, G. (T.) *acostaensis humerosa*, G. (T.) *scitula scitula*, G. (T.) *crassaformis ronda*, *Globigerinoides conglobatus conglobatus*, G. *elongatus*, G. *ruber*, G. *quadrilobatus*, *Globigerina eggeri eggeri*, G. *falconensis*, G. *bulloides bulloides*, G. *parabulloides*, *Hastigerina siphonifera siphonifera*, *Orbulina* spp., *Sphaeroidinella dehiscent dehiscent*.

Age: Zone N. 21; Late Pliocene (from evidence of next deeper sample).

Sample 4-30-6-3, 120-122 cm:

As above plus *Pulleniatina obliquiloculata obliquiloculata*, *P. obliquiloculata praecursor* and *Globorotalia* (T.) *tosaensis tenuitheca*.

Age: Zone N. 21; Late Pliocene.

Sample 4-30-7-1, 76-78 cm:

Globorotalia (T.) cf. *tosaensis tenuitheca*, G. (T.) *pseudopima*, G. (T.) *crassaformis oceanica*, *Globorotalia* (G.) cf. *miocenica*, *Globigerinoides elongatus*, G. *ruber*, G. *quadrilobatus* (s.l.), *Orbulina* sp., *Sphaeroidinella dehiscent dehiscent*.

Age: ? Basal part Zone N. 21; later Pliocene.

Sample 4-30-8, Core catcher:

Sphaeroidinella dehiscent dehiscent forma immatura, *Sphaeroidinellopsis subdehiscent paenedehiscent*, *S. subdehiscent subdehiscent*, *Orbulina* spp., *Globorotalia*

(*G.*) *cultrata menardii*, *G. (G.) miocenica*, *G. (G.) cf. multicamerata*, *G. (T.) acostaensis acostaensis*, *Globigerina bulloides apertura* (well developed), *Globorotalia (T.) acostaensis humerosa*, *Globigerinoides conglobatus conglobatus*, *G. obliquus extremus*, *G. bollii*, *G. quadrilobatus (s.l.)*, *G. elongatus*, *Globoquadrina altispira altispira*, *G. altispira globosa*.

Age: Zone N.19; Pliocene.

Sample 4-30-9-1, 80 cm:

Globorotalia (G.) cultrata cf. cultrata, *Globorotalia (T.) siakensis*, *Globigerina venezuelana*, *G. praebulloides (s.l.)*, *Globoquadrina dehiscens advena*, *Sphaeroidinellopsis subdehiscens nepenthes*, *Globigerinoides bollii*, *G. quadrilobatus (s.l.)*, *G. obliquus obliquus*.

Age: Zone N.14; Middle Miocene.

Sample 4-30-9-2, 60 cm:

As above.

Age: Zone N.14; Middle Miocene.

Sample 4-30-10-2, 78-80 cm:

Glauconitic sand residue with poor planktonic foraminiferal fauna but with *Globigerina nepenthes*, *Globorotalia (T.) siakensis*, *Globigerinoides bollii*, *G. quadrilobatus (s.l.)*, *Globigerina praebulloides (s.l.)*.

Age: Zone N.14; Middle Miocene.

Sample 4-30-10-2, 130 cm:

As above.

Age: Zone N.14; Middle Miocene.

Sample 4-30-10-3, 74-76 cm:

As above.

Age: Zone N.14; Middle Miocene.

Sample 4-30-10-3, 80 cm:

As above.

Age: Zone N.14; Middle Miocene.

Sample 4-30-11-1, 62-65 cm:

Glauconitic sandy residue with poor planktonic foraminiferal fauna, but with *Globorotalia (T.) siakensis*, *G. (T.) continuosa*, *Globigerina cf. nepenthes*, *G. druryi*, *Sphaeroidinellopsis subdehiscens subdehiscens*, *S. seminulina seminulina*, *Orbulina* sp.

Age: Probably uppermost Zone N.13; Middle Miocene.

Sample 4-30-11-2, 55-57 cm:

Sphaeroidinellopsis subdehiscens subdehiscens, *S. seminulina seminulina*, *Globigerinoides cf. subquadratus*, *G. bollii*, *Globorotalia (T.) siakensis*, *G. (T.) minutissima*, *Globigerina druryi*, *G. cf. woodi (s.l.)*, *Globoquadrina altispira (s.l.)*, *Orbulina* sp.

Age: Zone N.13; Middle Miocene.

Sample 4-30-11-2, 75 cm:

Globorotalia (G.) fohsi forma robusta, *Sphaeroidinellopsis subdehiscens subdehiscens*, *S. seminulina seminulina*, *Globorotalia (T.) siakensis*, *G. (T.) minutissima*,

G. (T.) continuosa, *Globoquadrina altispira (s.l.)*, *G. larmeyi obesa*, *Globigerina druryi*, *G. cf. woodi (s.l.)*, *Orbulina* sp.

Age: Zone N.13; Middle Miocene.

Sample 4-30-11, Core catcher:

Sphaeroidinellopsis subdehiscens subdehiscens, specimens ex interc *S. seminulina seminulina*—*S. subdehiscens*, *Globorotalia (T.) siakensis*, *Globigerina druryi*, *G. woodi*, *Globoquadrina altispira* (group), *Globigerinoides bollii*, *G. quadrilobatus (s.l.)*, *G. subquadratus*.

Age: Zone N.13; Middle Miocene.

Sample 4-30-12-1, 27-30 cm:

Poor planktonic foraminiferal fauna with *Globorotalia (T.) siakensis*, *Sphaeroidinellopsis seminulina seminulina*, *Globigerinoides bollii*, *G. subquadratus*, *G. quadrilobatus (s.l.)*, *Globigerina druryi*, *G. cf. woodi (s.l.)*, *Sphaeroidinellopsis cf. subdehiscens subdehiscens*.

Age: Probably uppermost Zone N.12; Middle Miocene.

Sample 4-30-12-2, 10-13 cm:

Specimens ex interc *Sphaeroidinellopsis seminulina (s.s.)*—*S. subdehiscens (s.s.)*, *S. cf. subdehiscens subdehiscens*, *Globorotalia (G.) fohsi (s.s.)*, *G. (G.) praefohsi*, *Globigerinoides subquadratus*, *G. quadrilobatus (s.l.)*, *G. bollii*, *Globigerina druryi*, *Globoquadrina larmeyi obesa*, *G. altispira (s.l.)*, *Globorotalia (T.) siakensis*, *G. (G.) miozea (s.l.)*.

Age: Uppermost Zone N.12; Middle Miocene.

Sample 4-30-12-3, 23-26 cm:

Sphaeroidinellopsis seminulina (s.l.), *Globorotalia (G.) fohsi (s.s.)*, *G. (G.) praefohsi*, *Globigerinoides subquadratus*, *G. quadrilobatus (s.l.)*, *G. bollii*, *Globigerina druryi*, *Globoquadrina altispira (s.l.)*, *Globorotalia (T.) siakensis*.

Age: Zone N.12; middle Miocene.

Sample 4-30-12-3, 122 cm:

Globorotalia (G.) fohsi (s.s.), *G. (G.) praefohsi*, *G. (T.) peripheroacuta*, *G. (G.) miozea (s.l.)*, *G. (T.) siakensis*, *Sphaeroidinellopsis seminulina seminulina*, *Globigerina druryi*, *G. cf. venezuelana*, *G. woodi woodi*, *G. woodi connecta*, *Globigerinoides bollii*, *G. subquadratus*, *G. quadrilobatus (s.l.)*, *Orbulina* spp., *Globorotaloides hexagona variabilis*.

Age: Zone N.12; Middle Miocene.

Sample 4-30-12-4, 0-3 cm:

As above.

Age: Zone N.12; middle Miocene.

Sample 4-30-12-4, 100 cm:

Globorotalia (G.) praefohsi, *G. (T.) peripheroacuta*, *G. (T.) sp. ex interc peripheroronda-peripheroacuta*, *G. (G.) miozea*, *G. (T.) siakensis*, *Orbulina* spp., *Globigerina druryi*, *G. woodi woodi*, *G. venezuelana*, *Globoquadrina dehiscens advena*, *G. altispira* (group), *Sphaeroidinellopsis seminulina seminulina*, *Globigerinoides bollii*

(rare), *G. obliquus obliquus*, *G. subquadratus*, *G. quadrilobatus*.

Age: Zone N.11; middle Miocene.

Sample 4-30-12-5, 5-8 cm:

As above.

Age: Zone N.11; middle Miocene.

Sample 4-30-12-5, 90 cm:

As above.

Sample 4-30-13-1, 0-3 cm:

Globorotalia (T.) *peripheroacuta*, *G. (T.) peripheroronda*, *G. (T.) scitula scitula*, *G. (G.) praemenardii praemenardii*, *G. (G.) miozea miozea*, *G. (T.) siakensis*, *G. (T.) obesa*, *Hastigerina siphonifera praesiphonifera*, *Globoquadrina dehiscens advena*, *G. altispira* (group), *Sphaeroidinellopsis seminulina seminulina*, *Orbulina* spp., *Globorotaloides hexagona variabilis*, *Globigerina woodi woodi*, *G. praebuloides* (s.l.), *G. juvenilis*, *Globigerinoides obliquus obliquus*, *S. subquadratus*, *G. quadrilobatus* (group), *Globigerinita incrusta*.

Age: Zone N.10; middle Miocene.

Sample 4-30-13-1, 120 cm:

As above.

Sample 4-30-13-2, 110-115 cm:

Globorotalia (T.) *peripheroacuta*, *G. (T.) peripheroronda*, *G. (T.) scitula scitula*, *G. (G.) miozea* (s.l.), *G. (G.) praemenardii* (s.l.), *G. (T.) siakensis*, *Globoquadrina dehiscens advena*, *G. altispira* (group), *Globigerina foliata*, *G. woodi* (s.l.), *G. praebuloides* (s.l.), *G. juvenilis*, *G. venezuelana*, *Globigerinoides quadrilobatus* (group), *G. subquadratus*, *Orbulina suturalis*, *Sphaeroidinellopsis seminulina seminulina*.

Age: Zone N.10; middle Miocene.

Sample 4-30-13-3, 5-7 cm:

As above.

Age: Zone N.10; middle Miocene.

Sample 4-30-13-3, 80 cm:

Very rich planktonic foraminiferal fauna. As above.

Age: Zone N.10; middle Miocene.

Sample 4-30-14-1, 83-85 cm:

As above.

Age: Zone N.10; middle Miocene.

Sample 4-30-15-1, 80-83 cm:

Globorotalia (G.) *praemenardii* (s.l.), *G. (G.) miozea*, *G. (T.) peripheroacuta*, *G. (G.) peripheroronda*, *G. (T.) scitula scitula*, *G. (T.) siakensis*, *G. (T.) obesa*, *Globigerina foliata*, *G. venezuelana*, *G. juvenilis*, *G. bradyi*, *Globigerinita incrusta*, *Globigerinoides quadrilobatus* (s.l.), *G. subquadratus*, *G. obliquus obliquus*, *Sphaeroidinellopsis seminulina seminulina*, *Orbulina suturalis*, *Globoquadrina dehiscens advena*.

Age: Zone N.10; Middle Miocene.

Sample 4-30-15-2, 8-11 cm:

Globorotalia (G.) *praemenardii* (s.l.), *G. (T.) peripheroacuta*, *G. (T.) peripheroronda*, *G. (T.) scitula scitula*, *G. (T.) scitula* cf. *praescitula*, *G. (T.) siakensis*, *G. (T.) obesa*, *Orbulina suturalis*, *Globoquadrina dehiscens dehiscens*, *G. dehiscens advena*, *G. altispira* (group), *G. baroemoenensis*, *Globigerina venezuelana*, *G. bradyi*, *G. foliata*, *Sphaeroidinellopsis seminulina seminulina*, *Globigerinoides subquadratus*, etc.

Age: Zone N.10; Middle Miocene.

Sample 4-30-15-2, 70 cm:

As above.

Age: Zone N.10; Middle Miocene.

Sample 4-30-15-3, 8-11 cm:

As above, but *G. (T.) peripheroacuta* very phylogenetically primitive.

Age: Lower Zone N.10; Middle Miocene.

Sample 4-30-15-3, 130 cm:

Globorotalia (G.) *praemenardii archeomenardii*, *G. (T.) peripheroronda*, *G. (T.) scitula praescitula*, *G. (T.) siakensis*, *Orbulina suturalis*, *Praeorbulina glomerata circularis*, *Globoquadrina altispira* (group), *G. dehiscens advena*, *G. baroemoenensis*, *Globigerina venezuelana*, *Globigerinoides subquadratus*, *G. obliquus obliquus*, *G. quadrilobatus*, *Sphaeroidinellopsis seminulina seminulina*, etc.

Age: Zone N.9; Middle Miocene.

Sample 4-30-15-4, 107 cm:

Globorotalia (*Clavatorella*) *bermudezi* (rare), *Praeorbulina glomerata glomerata* (rare), *P. glomerata circularis* (rare), *Globorotalia* (G.) *praemenardii archeomenardii*, *G. (T.) peripheroronda*, *G. (T.) siakensis*, *G. (T.) scitula praescitula*, *G. (T.) obesa*, *Globigerinitella insueta* (rare), *Sphaeroidinellopsis seminulina seminulina*, *Globigerinoides diminutus*, *G. sicanus* (rare), *G. quadrilobatus* (s.l.), *G. obliquus obliquus*, *G. subquadratus*, *Globigerina falconensis*, *G. juvenilis*, *Globigerinita incrusta*, *Globoquadrina* spp., etc.

Age: Zone N.8; lower part Middle Miocene.

Sample 4-30-15-5, 120 cm:

As above.

Age: Zone N.8; lower part Middle Miocene.

Sample 4-30-16-1, 117 cm:

As above, but no specimens of *P. glomerata circularis* present. However, *P. glomerata glomerata* and *P. glomerata curva* both present in the assemblage.

Age: Zone N.8; lower part Middle Miocene.

Hole 31

Sample 4-31-1-1, 74-76 cm:

Rich planktonic foraminiferal fauna with *Globorotalia* (G.) *truncatulinoides truncatulinoides*, *G. (G.) tumida*

tumida, *G. (G.) cultrata menardii*, *G. (G.) unguolata*, *Globigerinoides conglobatus conglobatus*, *G. ruber* (pink), *G. quadrilobatus* (s.l.), *Pulleniatina obliquiloculata finalis*, *Sphaeroidinella dehiscens dehiscens*, *Globigerina eggeri eggeri*, *Orbulina* sp.

Age: Zone N.22/Zone N.23; Pleistocene/Holocene.

Sample 4-31-1-2, 74-76 cm:

As above.

Sample 4-31-1-3, 74-76 cm:

As above.

Sample 4-31-1-4, 74-76 cm:

As above.

Sample 4-31-1-5, 73-75 cm:

As above.

Sample 4-31-1-6, 130 cm:

As above.

Sample 4-31-3-1, 74-76 cm:

Globorotalia (G.) miocenica, *G. (G.) cultrata cultrata*, *G. (G.) cultrata exilis*, *G. (T.) acostaensis pseudopima*, *G. (T.) acostaensis humerosa*, *G. (T.) inflata*, *G. (T.) crassaformis ronda*, *G. (T.) tosaensis tosaensis*, *Globigerina eggeri eggeri*, *Globigerinoides ruber*, *G. quadrilobatus* (s.l.), *G. conglobatus conglobatus*, *Pulleniatina obliquiloculata obliquiloculata*, *Sphaeroidinella dehiscens dehiscens*, *Orbulina* sp.

Age: Zone N.21; Late Pliocene.

Sample 4-31-3-2, 74-76 cm:

As above, but no specimens of *Pulleniatina*.

Sample 4-31-3-3, 73-74 cm:

As above, but no specimens of *Pulleniatina*.

Sample 4-31-3-5, 74-76 cm:

Much comminuted fauna with only *Sphaeroidinella dehiscens dehiscens*, *G. eggeri eggeri*, *Globigerinoides elongatus*, *G. quadrilobatus*, recognizable.

Age: Within the interval $\left[\begin{array}{l} \text{Zone N.21} \\ \text{Zone N.19 later part} \end{array} \right]$

Pliocene.

Sample 4-31-3-6, 74-76 cm:

Rich planktonic foraminiferal fauna with *Globorotalia (G.) miocenica*, *G. (G.) cultrata exilis*, *G. (G.) cultrata cultrata*, *Pulleniatina obliquiloculata obliquiloculata*, *Globorotalia (T.) acostaensis pseudopima*, *G. (T.) inflata*, *G. (T.) cf. tosaensis* (s.l.), *G. (T.) crassaformis ronda*, *G. (T.) crassaformis crassaformis*, *Sphaeroidinella dehiscens dehiscens*, *Globigerinoides conglobatus conglobatus*, *G. ruber*, *Globigerina eggeri eggeri*, *G. cf. pseudofoliata*, *Orbulina* spp.

Age: Probably lower part Zone N.21; Late Pliocene.

Sample 4-31-4-1, 75-77 cm:

Globorotalia (G.) miocenica, *G. (G.) cultrata* (s.l.), *G. (T.) acostaensis pseudopima*, *G. (T.) crassaformis crassaformis*, *Globigerinoides ruber*, *G. conglobatus conglobatus*, *G. quadrilobatus* (s.l.), *G. eggeri eggeri*, etc.

Age: Within the interval $\left[\begin{array}{l} \text{Zone N.21} \\ \text{Zone N.20} \end{array} \right]$; Pliocene.

Sample 4-31-5-1, 74-76 cm:

As above.

Sample 4-31-6-1, 74-76 cm:

Much comminuted fauna, no recognizable taxa.

Age: None.

Sample 4-31-6-2, 74-76 cm:

As above.

Sample 4-31-7-1, 66-68 cm:

Globorotalia (G.) multicamerata, *G. (G.) miocenica*, *G. (G.) cultrata exilis*, *G. (T.) acostaensis acostaensis*, *G. (T.) acostaensis pseudopima*, *G. (T.) acostaensis humerosa*, *Globigerina eggeri eggeri*, *G. eggeri multi-loba*, *Globigerina decoraperta*, *G. cf. nepenthes*, *Globoquadrina altispira* (group), *Globigerinoides obliquus extremus*, *G. bollii*, *G. conglobatus conglobatus*, *G. quadrilobatus* (s.l.), *Sphaeroidinella dehiscens dehiscens*, *S. dehiscens dehiscens forma immatura*, *Globoquadrina larmei obesa*, *Globorotalia (T.) crassaformis* (s.l.), *Globorotalia (G.) crassula conomiozea*.

Age: Zone N.20; Pliocene.

Sample 4-31-8-1, 74-76 cm:

Much comminuted fauna, no recognizable taxa.

Age: None.

Sample 4-31-8-2, 74-76 cm:

As above.

Sample 4-31-9-2, 1-4 cm:

Globorotalia (T.) pseudokugleri, *G. (T.) siakensis*, *Globigerinoides quadrilobatus* cf. *immaturus*, *Globigerinita dissimilis dissimilis*, *G. dissimilis ciperoensis*, *G. unicava unicava*, *Globoquadrina dehiscens praedehiscens*.

Age: Zone N.4; Early Miocene.

Sample 4-31-9-3, 0-4 cm:

As above, plus *G. (T.) kugleri*.

Age: Zone N.4; Early Miocene.

Sample 4-31-9-3, 140 cm:

As above, but no specimens of *Globigerinoides quadrilobatus* (s.l.) observed.

Age: Zone N.4 or ? uppermost Zone N.3; Early Miocene or ? topmost Oligocene.

Sample 4-31-10-1, 0-4 cm:
As above.

Sample 4-31-10-1, 136 cm:
Globorotalia (T.) kugleri, *G. (T.) pseudokugleri*, *G. (T.) siakensis*, *G. (T.) cf. mendacis*, *G. (T.) permicra*, *Globigerina bradyi*, *G. juvenilis*.

Age: Zone N.4 or ? uppermost Zone N. 3; Early
Miocene or ? topmost Oligocene.

Sample 4-31-10-2, 0-6 cm:
No significant planktonic foraminifera.

Sample 4-31-10-3, 0-3 cm:
As above.

Sample 4-31-10-4, 0-3 cm:
As above.

Sample 4-31-10-5, 0-3 cm:
As above.

Sample 4-31-10-5, 107 cm:
As above.