12. LEG 4 GRAIN SIZE RESULTS

A. C. Pimm, Deep Sea Drilling Project, East Coast Repository
Lamont-Doherty Geological Observatory, Palisades, N. Y.

INTRODUCTION

A complete list of the grain size results is given as Table 1; triangle diagrams for each hole are given as Figure 1; and, the sand/silt and silt/clay boundaries downhole are shown in Figure 2A, B. Methods used for grain size analysis are given in the appendix to this volume.

RESULTS

Site 23

A hole drilled at Site 23 mainly penetrated clays with a clay content of 85 to 90 per cent. The clays are olive-gray in Cores 1 and 4, and are yellowish-brown and zeolitic in Core 3 and part of Core 4. Grain size samples from Core 5 are of silty clay and sand, probably of turbidite origin; the wide separation of the sand/silt and silt/clay boundaries in different parts of the same core is seen in Figure 2.

Site 24

Sediments from Holes 24 and 24A are described elsewhere in this volume as deep water geosynclinal "flysch" deposits. Cores 1 and 2 from Hole 24 are of sandy clay and clayey sand of terrigenous origin. In Core 4 from the same hole the sediments are brown and greenish-gray to blue with a clay content of 90 to 95 per cent. Cores 1 and 2 from Hole 24A contain clays similar to those from Hole 24. Silty clay and sand-silt-clay from Cores 3 and 4 are terrigenous derived gray-wackes.

Site 25

All samples taken from Holes 25 and 25A are of foraminiferal chalk ooze. The grain size, dependent upon the amount of foraminifera present, is mainly clayey sand or sandy clay in Cores 1 and 2, and sand-silt-clay in Cores 3 and 4.

TABLE 1
Grain Size Results, Leg 4

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
23	1	1	73-74	0.5	13.3	86.2	Clay
23	1	2	4	0.9	13.8	85.3	Clay
23	1	5	0	6.6	14.0	79.4	Clay
23	3	2	16	0.2	8.4	91.4	Clay
23	3	3	23	0.2	7.5	92.3	Clay
23	4	1	10-12	5.7	9.5	84.8	Clay
23	4	3	40-42	1.9	7.5	90.6	Clay
23	4	4	65-67	1.9	8.7	89.4	Clay
23	5	1	22	17.8	21.0	61.2	Silty clay
23	5	1	46	83.2	6.8	10.0	Sand
24	1	1	26-28	42.7	12.0	45.3	Sandy clay
24	1	1	34	25.7	10.2	64.1	Sandy clay
24	2	1	15-16	71.7	12.1	16.2	Clayey sand
24	4	2	13	0.1	5.5	94.4	Clay
24	4	3	29	2.3	8.5	89.2	Clay

TABLE 1 – *Continued*

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
24	4	4	6	0.2	4.0	95.8	Clay
24A	1	1	1-24	0.3	17.2	82.5	Clay
24A	2	1	1-27	0.3	20.6	79.1	Clay
24A	3	1	84	0.3	24.9	74.8	Silty clay
24A	4	2	115	32.4	20.7	46.9	Sand-silt-clay
25	1	0	10	27.8	17.0	55.2	Sandy clay
25	1	1	26	50.3	11.7	38.0	Clayey sand
25	1	1	110	31.2	15.5	53.3	Sandy clay
25	1	2	30	43.8	13.6	42.6	Clayey sand
25	1	2	110	31.5	16.0	52.5	Sandy clay
25	1	4	39	52.7	5.3	42.0	Clayey sand
25	1	4	110	55.3	11.1	33.6	Clayey sand
25	1	5	30	54.5	12.1	33.4	Clayey sand
25	1	5	130	33.5	20.1	46.4	Sand-silt-clay
25	1	6	68	62.3	9.3	28.4	Clayey sand
25	2	2	30	60.2	0.4	39.4	Clayey sand
25	2	3	70	50.2	12.6	37.2	Clayey sand
25	3	1	125	18.7	28.0	53.3	Silty clay
25	3	2	38	38.0	22.6	39.4	Sand-silt-clay
25	3	2	114	43.3	23.8	32.9	Sand-silt-clay
25	3	3	75	43.0	21.2	35.8	Sand-silt-clay
25	3	4	75	43.7	19.1	37.2	Clayey sand
25	3	5	75	46.1	22.4	31.5	Sand-silt-clay
25	3	6	75	44.3	20.0	35.7	Sand-silt-clay
25	4	1	33	44.6	17.8	37.6	Clayey sand
25	4	2	75	48.3	24.3	27.4	Sand-silt-clay
25	4	3	75	48.8	25.1	26.1	Sand-silt-clay
25	4	4	76	46.5	25.3	28.2	Sand-silt-clay
25A	1	1	7	74.7	12.0	13.3	Clayey sand
25A	1	1	87	50.0	24.3	25.7	Sand-silt-clay
25A	1	2	75	53.1	25.2	21.7	Sand-silt-clay
26	1	2	10	32.3	18.6	49.1	Sandy-clay
26	1	2	90	0.1	18.5	81.4	Clay
26	1	3	13	0.1	20.2	79.7	Clay
26	1	3	90	1.1	22.0	76.9	Clay
26	1	3	138	9.4	79.1	11.5	Silt
26	2	1	124	0.1	22.2	77.7	Clay
26	2	2	10	0.1	24.4	75.5	Clay

TABLE 1 – *Continued*

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
26	2	2	95	0.0	22.7	77.3	Clay
26	3	2	20	0.0	21.3	78.7	Clay
26	3	2	100	0.0	23.5	76.5	Clay
26	5	1	13	0.0	15.6	84.4	Clay
26	5	1	87	0.1	17.0	82.9	Clay
26	5	2	26	0.0	16.6	83.4	Clay
26	5	2	90	0.0	18.0	82.0	Clay
26	5	3	27	0.0	19.5	80.5	Clay
26	5	3	96	0.0	30.0	70.0	Silty clay
26	5	4	20	0.1	71.1	28.8	Clayey silt
26	5	4	87	0.0	19.2	80.8	Clay
27	1	1	13	0.1	4.3	95.6	Clay
27	1	2	15	0.1	5.6	94.3	Clay
27	1	6	73	0.3	7.4	92.3	Clay
27	2	1	86	0.2	8.4	91.4	Clay
27	2	2	13	0.2	7.6	92.2	Clay
27	2	3	21-23	0.9	7.0	92.1	Clay
27	2	3	80-82	0.1	6.3	93.6	Clay
27	3	1	47	0.1	8.3	91.6	Clay
27	3	2	16-17	0.1	10.4	89.5	Clay
27	4	1	35	0.1	5.2	94.7	Clay
27	4	2	4	0.6	8.9	90.5	Clay
27	4	3	16	0.4	6.5	93.1	Clay
27	5	1	19	0.5	7.4	92.1	Clay
27	5	2	39	0.3	20.8	78.9	Clay
27	5	3	25	4.0	19.4	76.6	Clay
27	5	4	33	0.0	14.7	85.3	Clay
27	6	2	90-100	0.6	8.9	90.5	Clay
27	6	3	20-30	1.3	7.3	91.4	Clay
27A	1	1	29-31	0.1	8.9	91.0	Clay
27A	1	1	77-79	0.1	9.0	90.9	Clay
27A	1	4	15-17	0.0	8.4	91.6	Clay
27A	1	5	31	0.1	8.8	91.1	Clay
27A	1	6	18	0.1	9.7	90.2	Clay
27A	1	6	90	0.1	9.0	90.9	Clay
27A	2	3	25	0.2	14.2	85.6	Clay
27A	2	3	73	0.2	75.6	24.2	Silt
27A	3	1	132	0.2	21.2	78.6	Clay

TABLE 1 – *Continued*

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
27A	3	2	22	0.0	77.8	22.2	Silt
27A	3	3	30	0.0	9.4	90.6	Clay
27A	4	1	34-36	0.0	8.3	91.7	Clay
27A	4	2	49	0.0	11.1	88.9	Clay
27A	4	3	15	0.1	15.1	84.8	Clay
27A	4	3	77	0.3	70.5	29.2	Clayey silt
27A	4	4	33	0.1	75.2	24.7	Silt
27A	4	5	35	0.1	5.1	94.8	Clay
27A	4	6	34	0.1	12.8	87.1	Clay
27A	5	1	32-34	0.0	10.1	89.9	Clay
27A	5	2	46	0.3	14.3	85.4	Clay
27A	5	3	22	0.2	10.6	89.2	Clay
27A	5	4	57	0.1	12.5	87.4	Clay
27A	5	5	84	5.0	1.5	93.5	Clay
28	2	2	10	0.7	21.6	77.7	Clay
28	2	3	16	0.2	28.9	70.9	Silty clay
28	2	4	18	0.1	13.1	86.8	Clay
28	3	1	11	0.1	64.4	35.5	Clayey silt
28	3	2	36	0.2	74.1	25.7	Clayey silt
28	3	3	42-44	0.9	62.9	36.2	Clayey silt
28	3	5	11	0.1	59.4	40.5	Clayey silt
28	3	6	24	0.1	58.6	41.3	Clayey silt
29	1	2	20	12.7	40.8	46.5	Silty clay
29	1	3	30	4.7	30.6	64.7	Silty clay
29	2	1	52	8.6	42.8	48.6	Silty clay
29	2	3	50	2.3	19.7	78.0	Clay
29	3	1	63	2.0	18.3	79.7	Clay
29	4	1	109-111	1.3	16.8	81.9	Clay
29	4	2	53	2.4	15.5	82.1	Clay
29	4	3	48	2.4	12.8	84.8	Clay
29	4	4	16-19	0.3	14.2	85.5	Clay
29	5	1	122	0.0	7.1	92.9	Clay
29	6	1	105-107	0.1	7.9	92.0	Clay
29	7	1	137	0.8	49.4	49.8	Silty clay
29	8	1	118	23.0	61.5	15.5	Sandy silt
29	9	1	47-49	16.3	56.8	26.9	Clayey silt
29	9	2	48	17.2	67.4	15.4	Sandy silt
29	9	3	18	16.7	59.4	23.9	Clayey silt

TABLE 1 – Continued

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
29	9	4	35	10.6	47.7	41.7	Clayey silt
29	9	5	21	8.7	41.0	50.3	Silty clay
29	9	6	33	8.6	56.0	35.4	Clayey silt
29	10	1	61	16.7	42.2	41.1	Clayey silt
29	10	2	39-41	8.6	45.5	45.9	Silty clay
29	10	3	29-31	10.5	39.3	50.2	Silty clay
29	15	6	45	14.4	42.2	43.4	Silty clay
29	16	1	46	10.0	42.1	47.9	Silty clay
29	16	2	33	16.2	41.7	42.1	Silty clay
29	16	3	46	20.4	41.7	37.9	Sand-silt-clay
29	16	4	38	10.5	39.4	50.1	Silty clay
29	16	5	32	19.8	38.8	41.4	Silty clay
29	16	6	33	15.7	41.2	43.1	Silty clay
29	17	1	31	23.2	40.8	36.0	Sand-silt-clay
29	17	2	32	19.3	49.5	31.2	Clayey silt
29	17	3	31	21.1	59.3	19.6	Sandy silt
29	17	4	32	24.4	40.8	34.8	Sand-silt-clay
29	17	5	32	19.9	42.7	37.4	Clayey silt
29	17	6	32	18.8	44.6	36.6	Clayey silt
29	18	2	42	18.2	42.2	39.6	Clayey silt
29A	2	1	40	0.2	9.6	90.2	Clay
29A	4	1	60	0.1	19.0	80.9	Clay
29A	5	1	56-58	0.1	11.4	88.5	Clay
29B	1	1	33	0.0	22.6	77.4	Clay
29B	1	4	131	0.0	8.0	92.0	Clay
29B	1	5	37	0.1	8.8	91.1	Clay
29B	1	6	41	0.1	9.0	90.9	Clay
29B	2	2	41	0.6	11.4	88.0	Clay
29B	2	3	22	0.0	10.2	89.8	Clay
29B	2	4	22	0.1	14.5	85.4	Clay
29B	3	1	42	0.2	24.9	74.9	Silty clay
29B	3	2	28	0.1	11.6	88.3	Clay
29B	4	1	27	0.1	10.7	89.2	Clay
29B	4	2	27	0.2	11.4	88.4	Clay
29B	4	3	32	0.1	7.3	92.6	Clay
29B	4	4	28	0.1	5.7	94.2	Clay
29B	5	1	31	2.1	25.2	72.7	Silty clay
29B	5	2	27	4.5	44.0	51.5	Silty clay

TABLE 1 – *Continued*

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
29B	5	3	31	3.2	43.6	53.2	Silty clay
29B	5	4	27	0.1	23.0	76.9	Clay
29B	6	1	104	15.5	45.3	39.2	Clayey silt
29B	8	1	31	21.5	43.3	35.2	Sand-silt-clay
29B	8	3	26	24.8	47.9	27.3	Sand-silt-clay
29B	8	4	31	24.0	50.9	25.1	Sand-silt-clay
29B	8	5	31	27.7	48.1	24.2	Sand-silt-clay
29B	8	6	26	25.8	47.2	27.0	Sand-silt-clay
29B	9	1	36	20.7	38.4	40.9	Sand-silt-clay
29B	9	2	28	17.1	37.5	45.4	Silty clay
29B	9	3	27	16.9	13.7	69.4	Sandy clay
29B	9	4	26	18.5	33.8	47.7	Silty clay
29B	9	5	20	17.3	34.1	48.6	Silty clay
29B	9	6	22	16.6	41.6	41.8	Silty clay
29B	10	2	27	16.1	37.6	46.3	Silty clay
29B	10	3	32	20.6	37.7	41.7	Sand-silt-clay
29B	10	4	33	18.3	42.9	38.8	Clayey silt
29B	10	5	44	19.1	43.6	37.3	Clayey silt
30	1	1	115	14.4	18.6	67.0	Silty clay
30	1	5	36	8.2	20.9	70.9	Silty clay
30	2	3	31	9.8	23.7	66.5	Silty clay
30	2	4	32	1.1	20.5	78.4	Clay
30	3	2	31	2.2	25.2	72.6	Silty clay
30	3	3	21	6.1	51.2	42.7	Clayey silt
30	3	4	22	0.8	37.1	62.1	Silty clay
30	4	1	110	7.2	36.5	56.3	Silty clay
30	4	2	22	0.8	33.3	65.9	Silty clay
30	5	1	42	2.2	16.2	81.6	Clay
30	5	2	29	3.0	23.2	73.8	Silty clay
30	6	1	31	1.3	23.4	75.3	Clay
30	6	3	21	1.4	21.1	77.5	Clay
30	7	1	68	2.0	19.6	78.4	Clay
30	9	1	82	4.3	50.0	45.7	Clayey silt
30	9	2	38	6.4	54.7	38.9	Clayey silt
30	10	1	30	14.3	51.8	33.9	Clayey silt
30	10	2	30	10.9	52.9	36.2	Clayey silt
30	10	3	30	24.6	40.2	35.2	Sand-silt-clay
30	11	2	25	2.3	54.8	42.9	Clayey silt

TABLE 1 – *Continued*

Hole	Core	Section	Interval Sampled (cm)	% Sand	% Silt	% Clay	Classification
30	13	3	7	5.4	50.3	44.3	Clayey silt
30	14	1	145	36.6	35.7	27.7	Sand-silt-clay
30	15	1	136	32.5	39.4	28.1	Sand-silt-clay
30	15	2	10	44.6	33.2	22.2	Sand-silt-clay
30	15	3	10	41.4	38.1	20.5	Sand-silt-clay
30	15	4	8	37.1	42.2	20.7	Sand-silt-clay
30	15	5	7	27.8	50.5	21.7	Sand-silt-clay
31	1	1	41	11.1	12.2	76.7	Clay
31	1	2	30	9.2	8.8	82.0	Clay
31	1	3	91	9.4	11.0	79.6	Clay
31	1	4	30	10.3	10.9	78.8	Clay
31	1	5	28	9.3	14.2	76.5	Clay
31	3	1	25	10.5	14.9	74.6	Silty clay
31	3	2	28	13.2	14.8	72.0	Silty clay
31	3	3	36	8.7	22.3	69.0	Silty clay
31	3	5	26	6.2	19.7	74.1	Silty clay
31	3	6	31	11.9	5.2	82.9	Clay
31	4	1	78	11.4	19.2	69.4	Silty clay
31	4	2	39	15.0	18.3	66.7	Silty clay
31	5	1	45	6.4	19.9	73.7	Silty clay
31	6	1	32	4.2	18.6	77.2	Clay
31	6	2	31	6.5	19.8	73.7	Silty clay
31	7	1	65	4.7	20.2	75.1	Clay
31	7	1	140	0.2	8.8	91.0	Clay
31	8	1	58	1.3	20.2	78.5	Clay
31	8	2	27	0.7	20.3	79.0	Clay
31	9	2	58	9.0	43.8	47.2	Silty clay
31	10	1	146-147	1.8	51.4	46.8	Clayey silt
31	10	2	143	0.6	50.2	49.2	Clayey silt
31	10	3	146-148	0.9	54.1	45.0	Clayey silt
31	10	4	145	0.3	44.3	55.4	Silty clay
31	10	5	145	0.3	50.0	49.7	Clayey silt

Figure 1. Triangle diagram of grain size from Leg 4.

Figure 2A. Down-hole plots of sand/silt and silt/clay boundaries.

Figure 2B. Down-hole plots of sand/silt and silt/clay boundaries.

Site 26

Sediments penetrated in this site are mainly olive-gray to medium gray clays containing about 75 to 85 per cent clay and 15 to 26 per cent silt. A few silt layers probably of turbidity current origin, occur in places (Core 1, Section 3). A graded bed ranging from clayey silt at the base through silty clay to clay at the top is seen in Sections 3 and 4 of Core 5. The presence of these coarser layers of sediment are particularly noticeable in Figure 2.

The lithological summary describes several fine to medium, and occasionally coarse sand stringers (of turbidite origin); but, these were either too thin for sampling, or just were not selected for grain size analysis.

Site 27

Hole 27 penetrated brown clay, in places greenish or grayish, with a clay content of 90+ per cent.

Hole 27A contained grayish-brown, or rarely greenish-brown, clays with a clay content of 85 to 90 per cent. A few graded beds of silt and clayey silt occur in the clay; their occurrence is well exhibited in Figure 2. It should be noted that no nannoplankton were detected in sediments from this hole, and the silt fraction is made up of quartz, pyrite and manganese. In most other holes from Leg 4, and the other Atlantic sites, calcareous nannoplankton usually make up a sizeable proportion of the silt fraction (provided the sediment was deposited above the carbonate compensation depth).

Site 28

Sediments penetrated in this site were brown clay with thin silty clay layers in Core 2, overlying clayey silt in Core 3. The clayey silt contains 58 to 74 per cent silt, and less than 1 per cent sand, which is made up of nannoplankton and siliceous fossils.

Site 29

Hole 29 penetrated a variety of sediments, mainly of clay and silty clay size in the upper part of the section, but becoming coarser lower down due to a predominance of Radiolaria as the main fossil type, in contrast to nannoplankton in the upper part. Core 1 is a foraminiferal-nannoplankton chalk ooze of silty clay-size; Core 2 is similar but in places contains more or less

nannoplankton and, therefore, the sediment either falls in the silt or clay size range. Cores 3 through 6 contain mainly olive brown and brown clays. Core 7 is a nannoplankton chalk ooze of silty clay size. Core 8 shows the beginning of the influx of Radiolaria and also a volcanic component; the sediment is a radiolarian ash ooze of sandy silt size. Cores 9 and 10 are clayey silt and silty clay radiolarian oozes. Cores 15 through 18 are also radiolarian oozes and fall in the sand-silt-clay, clayey silt, and silty clay size ranges.

Hole 29A penetrated brown clay with a clay content of 80 to 90 per cent. Hole 29B penetrated a sequence of brown clays (Cores 1 through 4), silty clays and clayey silts composed of nannoplankton chalk ooze, (Cores 5, 6 and 7), and then from Core 8 down the sediments contain siliceous fossils in place of the calcareous ones above. Core 8 is a sand-silt-clay composed of an ash radiolarian ooze, and Cores 9 and 10 were silty clay or clayey silt of similar composition; the sand content (larger Radiolaria mainly) varies from 15 to 25 per cent.

Site 30

This site penetrated clay or silty clay in its upper part (Cores 1 through 7) and clayey silt or sand-silt-clay lower down (Cores 9 through 15). The finer-grained sediments in the upper part are greenish-gray, calcareous (nannoplankton) and contain less than 10 per cent sand-sized material. The higher sand and silt content of sediments in the lower part of the section here is due to an abundance of volcanic ash, fossils and glauconite.

Site 31

This site penetrated fine-grained sediments ranging in size from clay to clayey silt, the variation in grain size is related to the calcareous fossils present in the sediments.

Core 1 comprised 75 to 80 per cent clay, with sand and silt in about equal amounts. Core 3 is either silty clay or clay with both nannoplankton and foraminifera (in this chalk ooze); Cores 4 through 6 are similar to Core 3 with silt varying from 15 to 20 per cent of the sediment. Cores 7 and 8 contain marl ooze of clay size. Cores 9 and 10 are silty clay or clayey silt, with silt varying from 43 to 54 per cent in these nannoplankton chalk oozes; the sand usually makes up less than 1 per cent of the sediment.