9. SITE 740¹

Shipboard Scientific Party²

HOLE 740A

Date occupied: 23 January 1988
Date departed: 27 January 1988
Time on hole: 3 days, 20 hr, 45 min
Position: 68°41.22'S, 76°43.25'E
Bottom felt (rig floor; m, drill-pipe measurement): 818.0
Distance between rig floor and sea level (m): 10.5
Water depth (drill-pipe measurement from sea level; m): 807.5
Total depth (rig floor; m): 1043.5
Penetration (m): 225.5
Number of cores: 31
Total length of cored section (m): 225.5
Total core recovered (m): 72.19
Core recovery (%): 32
Oldest sediment cored:
 Depth sub-bottom (m): 225.50
 Nature: sandstone
 Earliest age: ?
 Measured velocity (km/s): 2.064

HOLE 740B

Date occupied: 27 January 1988
Date departed: 27 January 1988
Time on hole: 4 hr, 45 min
Position: 68°41.22'S, 76°43.25'E
Bottom felt (rig floor; m, drill-pipe measurement): 824.9
Distance between rig floor and sea level (m): 10.5
Water depth (drill-pipe measurement from sea level; m): 814.4
Total depth (rig floor; m): 826.5
Penetration (m): 1.6
Number of cores: 1
Total length of cored section (m): 1.6
Total core recovered (m): 1.60
Core recovery (%): 100
Oldest sediment cored:
 Depth sub-bottom (m): 1.6
 Nature: diatom ooze
 Earliest age: Quaternary

Principal results: Site 740 is situated in the inner part of Prydz Bay, eastern Antarctica, at 68°41.22'S, 76°43.25'E, in 807.5 m of water. The objective of the drilling was to date the oldest sequence overlying the basement in eastern Prydz Bay, thus possibly providing a date for the formation of the Lambert Graben or the separation of the Antarctic continent from the Indian plate.

Drilling began on 25 January 1988, and was spudded in a local mud basin of a 800-m-deep erosional trough approximately 30 km from the coast. Based on our experience at Site 739, rotary drilling was used. At 15 m below seafloor (mbsf), a unit containing large, hard boulders was found, and further drilling in this and deeper units caused excessive wear on the bit. In order to change the worn bit, a free-fall funnel or minicone was deployed, and an additional mud-line core (Hole 740B) was taken.

Although the sedimentary succession at Site 740 is incomplete because of poor core recovery, it can be divided into three lithologic units. Unit I (0–23.2 mbsf) is a soft diatomaceous ooze alternating with clayey silt with minor diatoms and silty clay with gravel, coarse sand, and minor diatoms. The diatom ooze contains 10%–20% terrigenous clastic grains, 15%–95% diatoms, and 0%–5% radiolarians and silicoflagellates. The clayey silt contains up to 15% diatoms and 2% radiolarians; major nonbiogenic components are quartz (up to 45%) and feldspar (up to 10%). The unit is late Pliocene to Holocene in age.

Unit II (23.2–56.6 mbsf) is drilling breccia composed of gneiss clasts and acid/intermediate igneous rocks. The unit is Pliocene to Holocene in age.

Unit III (56.6–225.5 mbsf, total depth) is a greenish gray and red sandstone alternating with siltstone and claystone. The ratio of sandstone to siltstone and claystone decreases upward through the succession, which consists of fining-upward and rare coarsening-upward sequences. The sandstone is generally cross-stratified in the lower part and less so in the upper part of the unit. The sandstone is locally bioturbated and contains thin conglomerates and rare mud cracks. The siltstones and claystones show color mottling and veining indicative of pedogenesis.

The sedimentary succession records major changes in the depositional environment. The oldest sediments of Unit III, situated approximately 300 m above the basement, represent a "red bed"-type sequence deposited within the proximal reaches of a fluvial depositional system. Climatic conditions at the time were probably warm and dry. Unit II, which was very poorly recovered, may represent a lodgement till. Unit I probably indicates deposition of glaciomarine sediments. The diatom ooze in the upper part of the unit represents present-day deposition.

The age of Unit III is unknown because paleontological data are lacking. This unit possibly could be correlated with parts of the Amery Group sequence in the Lambert Graben 350 km to the southwest, suggesting a Permian age (Mond, 1972) or be part of the Mesozoic(?)–Cenozoic "molasse and/or riftogenic complex," constituting the preglacial sediments in Prydz Bay according to Fedorov et al. (1982).

The only trace of hydrocarbons detected at the site was 8 ppm ethane and 1140 ppm methane in a headspace sample from Section 119-740A-14R-3.

BACKGROUND AND OBJECTIVES

Site 740 (target Site PB-1) lies on the shelf of eastern Antarctica, around 30 km from land in eastern Prydz Bay (68°41'S, 76°43'E). The site lies in an erosional channel at the boundary between the basement and a sedimentary basin in 810 m of wa-

¹ Barron, J., Larsen, B., et al., 1989. *Proc. ODP, Init. Repts.*, 119: College Station, TX (Ocean Drilling Program).

² Shipboard Scientific Party is as given in the list of Participants preceding the contents.

ter. This is the innermost of a series of sites across the shelf along seismic profile 21 of Stagg (1985). The eastern part of this profile is shown in Figure 1, including Stagg's (1985) interpretation. Northwest of the uneven basement is a sequence of slightly inclined reflectors (seismic Unit PB-4) that are unconformably cut by the seafloor or relatively thin, presumably glacial layers. In the eastern part of the bay (profiles 13, 19, and 21), the unit seems to grade into seismic unit PS-2 and may form the base of the glacial sediments in this part of the bay. PS-2 is resting on a fairly even, strong reflector, which Stagg (1985) interpreted as the top of his seismic unit PS-5, but may represent the top of the basement.

Based on interpretation of aeromagnetic data (Fedorov et al., 1982) and seismic data (Stagg, 1985; Mizukoshi et al., in press), Prydz Bay is underlain by a sedimentary basin, which is probably an extension of the Lambert Graben (see "Introduction" and "Explanatory Notes" chapters, this volume). According to the map of the depth to Moho in Fedorov et al. (1982), a clear thinning of the crust seaward is seen along the east coast of Prydz Bay. Therefore, the deposition and tilting of unit PS-4 may be related to the stretching of the crust, forming the Lambert Graben and/or the rifting between the Antarctic and the Indian continental plate (Stagg, 1985). Dating of the sedimentary sequence base is expected to provide important information on the tectonic development in eastern Antarctica.

The age of the sediment is unknown. It could be correlated possibly with the Permian Amery Group in the Lambert Graben, 350 km to the southwest, described by Mond (1972), or be related to part of the Mesozoic or Cenozoic "molasse and/or riftogenic complex" proposed by Fedorov et al. (1982). A direct relation to the separation of the Antarctic from the Indian plate suggests a Neocomian age (Stagg, 1985). However, Ivanov (1983) suggested that a wide spectrum of pre-rift sediments, including coal, deltaic, and shallow-marine facies may have occurred under the Mesozoic molasse sediments. As very few Phanerozoic deposits are known from East Antarctica, any data on the sedimentary record will add interesting knowledge to the geological history of this part of the world.

The main objectives of Site 740 were: (1) to sample and date the oldest sediments overlying the basement in eastern Prydz Bay, possibly providing a date for the formation of the Lambert Graben and the separation of the Antarctic continent from the Indian plate, (2) to describe the environment and climatic development prior to the glaciation in eastern Antarctica, and (3) to study the overlying sediments as an example of glacial-interglacial sedimentation in the erosional channel in the inner part of the shelf (as part of the overall model of a shelf dominated by

glacial processes). Based on our experience at Site 739, rotary drilling was used.

SITE GEOPHYSICS

Site Survey

A brief geophysical survey (ODP line 119-05) was conducted near Site 740 prior to drilling (Fig. 2) with standard geophysical gear (see "Explanatory Notes" chapter). Transit satellite and dead reckoning navigation were used, as Global Positioning System satellite information was not available. Two site crossings were made to augment the only seismic-reflection line (six-channel Australian Bureau of Mineral Resources line PB-21) previously recorded near Site 740. The survey was conducted in flat seas, with no sea ice and only scattered large icebergs.

The first survey line crossed the area from northwest to southeast parallel and close to line PB-21 to obtain greater resolution of the dipping strata near the site. A tentative site was selected to sample the deepest part of the dipping sequence beneath Prydz Bay, including a poorly reflective unit onlapping basement. At the end of the traverse, the ship returned to the site, a beacon was dropped (shotpoint 1845), and the ship continued on for another mile to verify the site location. The ship proceeded to a point 2.5 km from the beacon to make a strike-line crossing of the site. The cross line, which is 1.7 km north of the site shows flat dips, verifying that the dipping sequence was not, as predicted, part of a closed structure. Sonobuoy 5 was deployed along the cross line 1.5 km from the closest point to the site and was recorded to a total offset of about 6 km. The seismic gear was recovered, and the ship returned to the beacon to begin drilling. The seismic results are discussed in the "Seismic Stratigraphy" section (this chapter).

Figure 1. Six-fold, stacked seismic section from the southeast margin of Prydz Bay with Stagg's (1985) designation of seismic units PS-6, PS-5, and PS-4 (after Stagg, 1985). The position of Site 740 is marked.

Figure 2. Index map of seismic lines recorded during site surveys for Site 740.

Sonobuoy Data

The sonobuoy was deployed to acquire velocity data from wide-angle reflection and refraction arrivals through the sedimentary section near Site 740 and to obtain a refraction velocity for the basement.

The sonobuoy was deployed successfully and appeared to be functioning normally; however, only very weak seismic signals were recorded from the direct and reflected arrivals, even with glassy sea conditions (Fig. 3). The poor response resulted in part from high noise levels in the water column and from poor penetration of seismic energy into the sedimentary section. High

noise levels were caused by seismic reverberations from seafloor multiple reflections, which were generated by each 9-s water gun shot. Also, the seismic energy was seemingly dispersed and attenuated by sedimentary units (alternating high- and low-velocity layers) at shallow subsurface depths.

Five wide-angle reflections were traced to offsets of up to 5 km (Fig. 4). Three reflections (AA, BB, and CC) gave good rms velocities, plotting as nearly straight lines on the time-squared vs. distance-squared graph. Two reflections were weak and could only be traced to short offsets; thus, they have poor rms velocities. The direct-arrival wave (*D*-wave) was difficult to identify because of a sharp velocity inversion within the upper

Figure 3. Vertical-incident seismic and sonobuoy seismic record for sonobuoy station 5. Site 740 lies about 1.5 km east of the point where the sonobuoy was deployed. See Figure 2 for location. **A.** Observed data. **B.** With interpretation.

Figure 4. Vertical-incident seismic profile across Site 740 and sonobuoy station 5 showing interval velocities for the sedimentary section.

15 m of the water column (see "Biology and Oceanography" section, this chapter); however, good rms velocities for the seafloor reflector indicated that the *D*-wave had been correctly identified to offsets of at least 4–5 km (see "Underway Geophysics" chapter, this volume). Refraction velocities were determined within the 4- to 5-km offset range and thus should be correct.

Strong and continuous refraction arrivals were recorded for the upper third of the sedimentary section and weak arrivals for the remainder of the section to basement. A refraction arrival was not recorded from basement possibly because of insufficient ship-to-sonobuoy offset. However, a strong wide-angle reflection was observed from basement. In the sedimentary section, the first-break refraction arrivals are curved indicating the presence of velocity gradients, rather than discrete velocity horizons. The good continuity of refractions in the upper sedimentary section suggests continually increasing velocities with depth. Velocities continue to increase at greater depths, but the attenuation of refraction amplitudes at about 3-km offsets suggests that thin, low-velocity horizons, thinning beds, or seismically-dispersive layers may be present. For computing velocities, the curved refraction arrivals were approximated using four straight lines, which gives the refraction velocities from top to bottom of the sedimentary section. The computed rms, interval, and refraction velocities are given in Table 1.

The sonobuoy data indicate that the sedimentary section has interval velocities of 2.46–3.08 km/s and refraction values of 2.27–2.96 km/s (values that are internally consistent and are high for sediments buried at depths of less than 700 m). The in-

Table 1. Preliminary results for sonobuoy 5, Site 740.

Horizon ^a	Vrms (km/s)	Vint ^b (km/s)	Ti ^b (s)	Zi (km)	Vrfr (km/s)	To (s)
AA	1.457	2.46	1.08	0.814	2.27	0.849
BB	1.705	3.08	1.35	1.149	2.53	0.941
CC	2.033	—	1.66	1.628	2.96	1.127

Note: Vrms = rms velocity for horizon; Vint = interval velocity computed using Vrms, Ti, and Dix equation (e.g., Vint between horizons AA and BB is 2.46 km/s); Ti = vertical-incidence reflection time (two-way) to horizon; Zi = total depth from sea level to horizon (water depth = 814 m); Vrfr = refraction velocities cannot be directly associated with a specific horizon, rather they comprise a gradient from near seafloor to basement (horizon CC); To = intercept time for refractor associated with horizon.

^a Letters of horizons do not correlate between drill sites and do not correspond with prior stratigraphic analyses.

^b Velocities and times used for computing depths Zi.

terval velocities are higher than those measured at Site 739 on the outer shelf for similar burial depths. The higher velocities for the stratigraphically lower and older units at Site 740 probably result from cementation and compaction by burial at greater depths than observed today. Uplift and erosion of about 500 m of sediment would explain the higher velocities. Intermittent loading of the sedimentary section by grounded glaciers is probably not the cause of the generally higher velocities because the compaction effects of ice loading do not extend below 100 to

170 m (see "Logging" sections of "Site 739" and "Site 742" chapters, this volume). The once-overlying sedimentary section probably was eroded by glaciers.

The acoustically dispersive character of the sonobuoy record, which precluded tracing wide-angle reflections, may reflect, in part, the variable lithologies beneath the site. The core data indicate zones of high- and low-velocity rocks (see "Physical Properties" section, this chapter) of continental origin. Although the overall refraction velocities increase with depth, the effects of interbedded low-velocity rocks and of discontinuous units would be the attenuation of the wide-angle reflection data, as seen at Sites 739 and 742, where such features are observed in downhole logs. The seismic dispersion makes determination from the sonobuoy data, of the thickness of the surface-mantling unit (0–60 mbsf; young glacial debris) that overlies the seaward-dipping sequence (old continental strata) impossible.

In summary, the sedimentary section at Site 740 is characterized by velocities that are higher than those observed within the sequences lying seaward and at similar subsurface depths. The higher velocities may result from compaction caused by burial by sediments that were glacially eroded from Site 740.

OPERATIONS

Site 739 to Site 740

Maersk Master had been sent ahead to observe the ice accumulation at the inner Prydz Bay operating area. Reports were encouraging as the drillship covered the 126 km to the outer end of the intended survey line. With seismic gear streamed, a profile was recorded southeastward to a point just 40 km from the continent and about 3 km outside the pack ice. The vessel then reversed course and returned about 8 km to the drill site, which had been selected from the first-pass profile. A positioning beacon was dropped at 1234 UTC. About 0.8 km past the drop point, a wide loop to the southwest was made to enable the ship to approach the beacon on a northeasterly course. A sonobuoy was launched, a refraction profile was made as the site was crossed, and the line was extended about an additional 3.2 km. The vessel then turned back toward the site, the gear was retrieved, and dynamic positioning began at Site 740, which is located about 56 km west of Australia's Antarctic Davis Station.

Site 740—Inner Prydz Bay

Hole 740A

A 90-m positioning offset was made to the northwest to maximize the soft sediment cover as seen on the 3.5-kHz precision depth recorder (PDR). The weight indicator showed contact at 818 m. Soft diatom ooze and clay were cored for 23 m before the hard, glacially-deposited material was again entered. The finer sediment was less overcompacted than at Site 739 but contained even more boulders and cobbles of very hard rock types. After patiently enduring the now-expected low rate of penetration (ROP) and recovery of the glacial till, the drillers broke through into a sequence of claystone, siltstone, and sandstone beds at 56 mbsf.

The sediment was firm but seemed quite drillable. The ROP and core recovery were both disappointingly low, and performance continued to deteriorate with depth (Table 2). When the ROP had dropped below 1 m/hr at 125 mbsf, we recognized that some type of bit failure had occurred. Bit balling was considered to be a possibility because of the low circulation rates and the presence of plastic clay and claystone. Standard deballing techniques were tried without success. The most likely problem was broken inserts in the drive rows of the bit as a result of prolonged drilling on the overlying gneiss and granite boulders.

As the ROP was unacceptable and the researchers were interested in coring deeper, a round trip and reentry were performed.

The free-fall funnel, or minicone, had been partially assembled in the moonpool area for fast deployment in case icebergs caused a temporary site abandonment. The drill string was raised until only the bottom-hole assembly (BHA) remained in the hole, the upper guide horn was lifted above the rig floor, and the two halves of the minicone were welded together around the drill pipe. The minicone was then guided through the moonpool before it was allowed to fall to the seafloor.

A routine pipe trip was made for a new bit. The old bit was balled with stiff clay, and the cones had been skidding on the bottom of the hole. One of the circulating nozzles was plugged. The tungsten carbide inserts showed severe wear, probably both from drilling hard formation and from skidding.

After the BHA had been assembled, the vibration-isolated TV (VIT) frame was deployed, and the drill string and the combination TV/sonar reentry module were lowered concurrently. Below 400 m, the concentration of schooling fish was so heavy that visibility below the camera was blocked. As the camera/sonar approached the bit, the lights were turned off in an effort to disperse the fish. Attempts were made, without success, to make the initial reentry approach with the sonar, using the floating glass-ball reflectors of the funnel as sonar targets. When the lights were turned on, the fish were gone, and the funnel was seen directly beneath the bit. The reentry was then a simple matter of quickly lowering the drill string and observing it enter the minicone. The minicone was noted to be resting fairly deep in the crater of the hole, with the floating reflectors not more than 1 m above the seafloor on their 3-m tethers.

The VIT frame was recovered, and the bit was run the short distance to total depth. When the top drive had been picked up, the driller was unable to break circulation. The drill string was plugged off, and all efforts, including pressuring to 5500 psi with the cementing unit, failed to clear it. Therefore, the round trip/reentry procedure was repeated to clean about 35 m of loose sand from the lower BHA.

The second reentry was routine, with fewer fish. The minicone was again found to be directly beneath the drill string, and no offsetting maneuvers were required. The top drive was picked up as soon as the VIT frame had been recovered, and the bit was drilled to total depth through 34 m of loose fill. The hole was then flushed with mud before the first inner core barrel was pumped into place.

Coring then continued for another 95 m with a much-improved ROP. Core recovery was improved, but continued to be below normal. The sediments continued to be a clastic sequence of apparent terrestrial origin and of little value to the scientific objectives of the voyage. At 225.5 mbsf, coring was terminated, the hole was plugged with cement, and the core bit was pulled above the seafloor.

Hole 740B

Because the seafloor sediments were incompletely recovered in the process of spudding Hole 740A, an additional mud-line core was requested. The bit was lowered to 826.5 m, with an inner barrel in place. The core barrel was recovered with 1.6 m of core and the sediment/water interface intact.

LITHOSTRATIGRAPHY AND SEDIMENTOLOGY

Rotary drilling at this site penetrated to a depth of 225.5 mbsf. However, core recovery averaged only 32%, with less than 12% recovery in the upper part of the succession. The sedimentary sequence at this site has been divided into three lithologic units (Table 3).

Table 2. Coring summary, Site 740.

Core no.	Date (Jan. 1988)	Time	Depth		Length		Recovery (%)
			top (mbsf)	bottom (mbsf)	cored (m)	recovered (m)	
119-740A-							
1R	23	2245	0.0	6.0	6.0	0.99	16.5
2R	23	2305	6.0	13.7	7.7	0.73	9.5
3R	24	0235	13.7	23.2	9.5	5.11	53.8
4R	24	0355	23.2	32.6	9.4	0.12	1.3
5R	24	0545	32.6	42.1	9.5	0.20	2.1
6R	24	0745	42.1	51.6	9.5	0.00	0.0
7R	24	1035	51.6	56.6	5.0	0.00	0.0
8R	24	1400	56.6	61.2	4.6	5.08	110.0
9R	24	1450	61.2	66.2	5.0	0.58	11.6
10R	24	1615	66.2	70.9	4.7	2.87	61.0
11R	24	1705	70.9	75.9	5.0	2.25	45.0
12R	24	1830	75.9	80.6	4.7	1.25	26.6
13R	24	1945	80.6	85.6	5.0	1.09	21.8
14R	24	2200	85.6	90.2	4.6	3.56	77.4
15R	25	0155	90.2	95.2	5.0	1.56	31.2
16R	25	0640	95.2	99.9	4.7	0.00	0.0
17R	25	0750	99.9	109.5	9.6	1.72	17.9
18R	25	0840	109.5	119.2	9.7	0.00	0.0
19R	25	1215	119.2	124.8	5.6	1.37	24.4
20R	25	1840	124.8	126.0	1.2	4.97	414.0
21R	26	1435	126.0	128.8	2.8	0.17	6.1
22R	26	1615	128.8	138.5	9.7	3.97	40.9
23R	26	1715	138.5	148.1	9.6	1.39	14.5
24R	26	2130	148.1	157.8	9.7	1.80	18.5
25R	26	2250	157.8	167.5	9.7	1.89	19.5
26R	27	0155	167.5	177.1	9.6	6.16	64.1
27R	27	0450	177.1	186.8	9.7	7.43	76.6
28R	27	0625	186.8	196.5	9.7	3.72	38.3
29R	27	0755	196.5	206.1	9.6	3.09	32.2
30R	27	0850	206.1	215.8	9.7	1.28	13.2
31R	27	1055	215.8	225.5	9.7	7.84	80.8
					225.5	72.19	32.0
119-740B-							
1R	27	1445	0.0	1.6	1.6	1.60	100.0
					1.6	1.60	100.0

Unit I is a diatomaceous ooze (clayey silt) with 75%–95% diatoms alternating with clayey silt with minor diatoms and silty clay with about 1% gravel and coarse sand. It contains boulder-sized gneiss clasts which caused severe drilling problems at this site, necessitating a change of bit and double reentry. This unit is probably late Pliocene to Holocene age.

Unit II consists of drilling breccia composed of amphibolite gneiss clasts and acid/intermediate igneous rocks. Recovery in this interval was very poor; hence, the thickness and overall composition of this unit is unknown. This unit is Pliocene to Quaternary in age.

Unit III consists of sandstone, siltstone, and claystone arranged in pedogenically influenced fining- and rare coarsening-upward sequences. It has yielded only traces of undatable algae spores, and its age at present is uncertain (see "Biostratigraphy" section, this chapter).

Smear-slide data are summarized on the barrel sheets and in Figure 5. Shipboard carbonate analyses were carried out on samples used for physical properties tests and on selected core samples (see "Inorganic Geochemistry" section, this chapter). A few samples from the upper part of the core were run on the Lantec Lab-Tec 100 particle-size analyzer. The results of these analyses are also included on the barrel sheets.

Changes in physical properties occur within Unit I at about 6.1 and 13.7 mbsf and at the boundary between Units II and III at about 56.6 mbsf. The changes in Unit I probably reflect changes in the amounts of diatom ooze, clayey silt, and gravel-bearing

silty clay (see "Physical Properties" chapter, this chapter). Another less significant change occurs at about 190 mbsf within the lower part of Unit III. This change probably reflects the increased amounts of sandstone and conglomerate in this part of the unit and the coarser grain size of the sandstones. As a result, the sandstones show increased levels of porosity and permeability.

Lithostratigraphy

Unit I

Sections 119-740A-1R-1 through 119-740A-3R-CC; depth, 0–23.2 mbsf.

Age: late Pliocene to Holocene, probably Holocene.

Unit I consists of unlithified diatom ooze alternating with clayey silt with minor diatoms and silty clay with gravel and coarse sand. Because of the poor recovery (16.5%, 9.5%, and 53.8%) of the three cores under discussion, the depth range of this unit is only an estimate. The upper portions of the cores are highly disturbed, and the sediment is soupy owing to its high water content. The fact that these sediments are normally consolidated and show no sign of ice loading suggests a Holocene age for the unit. The diatom ooze is olive (5Y 4/3) in color, with a diatom content ranging from 75% to 95%. The sediment is soft, homogeneous, and structureless. In addition to diatoms, the biogenic components of the sediment include radiolarians

Table 3. Summary of lithologic units, Site 740.

Unit	Lithology	Sedimentary structures	Thickness (m)	Depth (mbsf)	Core interval	Age
I	Diatom ooze with 75%-95% diatoms and minor radiolarians and silicoflagellates, alternating with clayey silt containing minor diatoms (up to 15%) and gravel-bearing silty-clay	Homogeneous, soft and structureless Homogeneous and structureless, with granule- and pebble-size clasts; one boulder of gneiss (dropstone?)	23.2	0-23.2	119-740A-1R-1 to 119-740A-3R-CC	late Pliocene to Holocene
II	Drilling breccia composed of clasts of amphibolite gneiss and acid/intermediate rocks, probably derived from diamicton. Very poor recovery in this interval.		33.4	23.2-56.6	119-740A-4R-1 to 119-740A-7R-CC	Pliocene to Quaternary
Contact not seen—no recovery interval						
III	Sandstone alternating with siltstone and claystone. Siltstone and claystone intraclasts occur throughout the sediments; quartz, quartzite, minor granite, and metamorphic extraclasts of pebble grade	Sandstone, siltstone, and claystone arranged in pedogenically influenced, erosively-based fining- and rare coarsening-upward sequences. Sandstones contain cross stratification. The siltstones and claystones are mottled, locally bioturbated, and ripple cross-laminated	168.9	56.6-225.5	119-740A-8R-1 to 119-740A-31R-CC	Unknown

(3%-5%) and silicoflagellates (2%-20%); nonbiogenic components are quartz (5%-20%) and feldspar (2%-6%). Section 119-740A-3R-1, 0-50 cm, emits a strong smell of hydrogen sulfide. The olive diatom ooze in Section 119-740A-3R-2 appears to be contaminated by clasts of darker gray (5Y 5/1) diatom ooze. Particle-size estimates based on smear slide analyses are as follows: 0%-50% sand, 40%-75% silt, and 10%-30% clay.

The clayey silt is gray (5Y 5/1) in color with minor diatoms. It differs from the diatom ooze in its color and lower diatom content, containing up to 15% diatoms and 2% radiolarians. Major nonbiogenic components are quartz (up to 45%) and feldspar (up to 10%). Particle-size estimates show that the sediment contains 5% sand, 60% silt, and 35% clay. The silty clay in the lower part of this unit contains about 1% gravel and coarse sand. The gravel consists of granules and pebbles (<4 cm in length) of quartz, granite, and gneiss.

Unit II

Sections 119-740A-4R-1 to 119-740A-7R-CC; depth, 23.2-56.6 mbsf.

Age: Pliocene-Quaternary.

Unit II is a drilling breccia composed of clasts of amphibolite gneiss and acid/intermediate igneous rocks. Recovery in this interval is very poor.

Unit III

Sections 119-740A-8R-1 to 119-740A-31R-CC; depth, 56.6-225.5 mbsf.

Age: unknown.

Unit III is a "red bed"-type sequence consisting of sandstone with interbedded siltstone and claystone. The sandstone

makes up 64% of the total recovered succession. However, stratigraphic partitioning of the data shows that sandstone is only 34% of the upper part of the succession (Cores 119-740A-8R to 119-740A-15R) whereas it is 74% of the lower part (Cores 119-740A-16R to 119-740A-31R). Thus, the succession shows a fining-upward trend internally structured by vertically stacked, fining-upward and rare coarsening-upward sequences up to 5 m thick. Unfortunately, sequence definition and stacking patterns are poorly constrained because of poor core recovery.

Individual fining-upward sequences, derived partly from pooled data, consist of sandstone sharply or gradationally overlain by siltstone and claystone. Good examples of these sequences are shown in Figure 6. Within these sequences, the sandstone, siltstone, and claystone may themselves show textural variations and fining-upward trends. The base of the sandstone is defined by an irregular scour surface of low relief (Fig. 7). This is overlain by a thin, matrix-supported polymictic conglomerate (Fig. 8) in some sequences; in others the conglomerate is missing, and the scour surface is directly overlain by coarse sandstone. The conglomerate is composed of elongate claystone and siltstone intraclasts and rounded to subrounded extraformational quartzite clasts and rare granite and metamorphic clasts. The clasts range from granule to cobble size. Intraclasts have a maximum length of 140 cm; extraformational clasts are generally smaller and attain maximum lengths of 5 cm. The conglomerates are typically ungraded, matrix supported, and poorly sorted (Fig. 8), but some elongate intraclasts show a long-axis horizontal fabric or imbrication.

The sandstones range in color from weak red (10R 4/2), red (10R 3/2, 10R 3/4), dark reddish gray (10R 3/1), dark reddish brown (5YR 3/2), and reddish black (10R 2.5/1) to greenish gray (5Y 6/1), grayish green (5Y 5/2), and dark greenish gray

Figure 5. Summary diagram of smear slide compositional data at Site 740.

(5G 4/2). Both red and green colors are often complexly intermixed with zones of greenish gray sandstone within the red sandstone and vice versa. The greenish gray zones within the red sandstone commonly show a dark organic-rich center around which the green color nucleated. The sandstones are coarse- to very coarse-grained with subordinate medium- and fine-grained varieties. The red sandstones are mostly immature and poorly sorted in contrast to the greenish gray sandstones, which are better sorted and more mature. Granules and small pebbles of quartz, red siltstone, and claystone are scattered throughout the sandstone or locally concentrated into thin conglomerate lenses and stringers. In some cases, the parallel alignment of two long axes in the clast imparts a distinct fabric to the rock.

The sandstones are composed predominantly of quartz (16%–94%), feldspar (1%–24%, including both fresh and weathered varieties), and mica (1%–47%, including chlorite), with the red sandstones loosely bound by a clay matrix (13%–69%; Fig. 5).

As a result the red sandstones are rather soft, poorly compacted, and friable. The greenish gray sandstones contain little or no clay matrix (0%–11%) and are better compacted and lithified. Two smear slide samples of particular interest are a garnet-rich (55%) sandstone (Sample 119-740A-29R-1, 65–66 cm) and a calcrite containing 87% micrite (Sample 119-740-26R-CC, 0–1 cm). The sandstones contain mainly planar and less common trough cross-bedding. This is particularly well developed in the lower part of the unit. Planar sets attain thicknesses of up to 1.15 m and are characterized by straight tabular foresets (Fig. 9) in sharp angular contact with the lower bounding surface. Trough sets are generally smaller (5–10 cm thick) and characterized by concave-shaped foresets showing opposing dip directions in the core (Fig. 10). Individual foresets are defined in two ways: by textural differences and by contrasts in color. Most foresets are normally graded, but they commonly show alternating red and green colors (Fig. 9) or lighter and darker shades of green or

Figure 6. Lithologic sections illustrating fining-upward sequences, Hole 740A.

red. Detailed examination of planar foresets in Section 119-740A-14R-1 shows that some have pale red interiors whereas others have a very distinct dark reddish brown zone about 1–2-mm-thick at the top. Ripple cross-lamination is also present in the sandstones, and in some examples it directly overlies cross bedding. Some structureless sandstone occurs in the upper part of the unit, but it is rare in the lower part. Other structures present within the sandstone are wavy and horizontal laminations, bioturbation, rare burrow structures, and mud cracks.

The siltstones and claystones are predominantly red in color with greenish gray spots and mottles. The proportion of silt and clay varies throughout individual sections. In addition, the silt-

stones and claystones contain scattered coarse sand grains and quartz granules, as well as claystone and siltstone intraclasts. Some siltstones and claystones, especially in the upper part of the succession, are very rich in clasts. Some intraclasts show a horizontal fabric or imbrication. The siltstones and claystones generally lack well-defined, primary sedimentary structures other than locally developed ripple cross-lamination (Fig. 11). This may be due in part to local pedogenic disruption and mottling of the sediments, bioturbation, and rootlet structures, which are commonly surrounded by greenish gray reduction haloes. Some of the mottling shows unusual diffuse zones of red, reddish brown, yellowish brown, and black. This type of mottling and

Figure 7. Irregular, angular scoured contact between coarse sandstone and underlying silty claystone. Section 119-740A-26R-3, 14–34 cm.

local carbonate veining is typical of pedogenic processes and has been recorded from many “red bed” sequences, ranging from Tertiary to Devonian and older in age (Turner, 1980).

Coarsening-upward sequences are not very common and generally consist of locally rippled, cross-laminated, fine sandstone or siltstone resting sharply or gradationally on claystone or clayey siltstone. The fine sandstone or siltstone grades upward into more medium- and coarse-grained sandstone, some capped by a very thin claystone drape.

Figure 8. Matrix-supported conglomerate of extraformational rounded quartzite pebbles, small subangular quartz pebbles and granules, and elongate siltstone and claystone intraclasts along the base of the fining-upward sequence. Section 119-740A-28R-3, 45–62 cm.

No fossils have been found in this unit so far, apart from undatable algae spores, and its age is unknown. However, seismic stratigraphy suggests that it may lie conformably beneath the succession cored at Site 741. This is a continental, organic-rich sequence containing abundant carbonized plant material and palynomorphs that provisionally date as Early Cretaceous. The possible presence of nonmarine Lower Cretaceous rocks and marine Upper Cretaceous rocks in eastern Antarctica was first suggested by Kemp (1972), but until now, nonmarine rocks of Cretaceous age or younger have not been found in this area. The sediments cored at Site 740 are the thickest sequence of continental sediments drilled by ODP to date. The absence of

Figure 9. Straight tabular foresets of a planar cross-bedding showing alternating light greenish gray and darker red foreset laminae. Section 119-740A-27R-3, 30–65 cm.

organic-rich sediments and diamictites found at Sites 739 and 741 may reflect later Pliocene-Pleistocene glacial erosion and sediment stripping of the shelf.

Interpretation

The absence of marine fossils and presence of scoured (channeled) fining-upward sequences containing rootlets and indi-

Figure 10. Small-scale (<10-cm-thick) trough cross-bedding showing opposing foreset dip directions. Section 119-740A-11R-2, 0–50 cm.

cations of pedogenesis in Unit III are consistent with deposition in a fluvially-influenced continental environment as documented by Allen (1965) and Miall (1977). Moreover, the coarse grain size of most of the conglomerate-bearing sandstones, the abundant intraclasts indicative of repeated episodes of scouring, the absence of abandoned channel fills, and the evidence of epsilon cross-stratification is more indicative of a braided channel system than a meandering one. This view is consistent with the ratio of sandstone to siltstone and claystone (especially in the lower part of the unit), the dominance of planar cross-bedding, and the sharp upper contacts shown by many sand bodies, implying rapid channel avulsion. The scour surface and overlying

Figure 11. Red clayey siltstone showing local ripple cross-lamination and pale greenish gray reduction spots. Note the dark organic-rich nucleus to one reduction spot. Section 119-740A-15R-1, 60–100 cm.

sand body are attributed to channel scouring and filling. The sandy bed of the channel was periodically molded into a series of transverse bars and/or two- and three-dimensional dune bed-forms, giving rise to the planar and trough cross-bedding. Fining- and coarsening-upward trends, recorded in the vertical variations in texture and sequence of sedimentary structures, are attributed to fluctuations in discharge related to decelerating and accelerating flood flows. Additional evidence for fluctuations in discharge and exposure of the depositional surface is the presence of small *in-situ* desiccated clay clasts.

The siltstone and claystone are interpreted as flood basin and overbank deposits laid down between the active channel tracts.

Overbank flooding accounts for the presence of coarse sandstone interbeds and lenses and intraclasts within the finer-grained siltstones and claystones, most of which were deposited under lower energy conditions. The environment was extensively colonized by vegetation, as evidenced by the rootlets and pedogenically-influenced flood basin deposits. Comparison with Holocene and older pedogenic sediments of a similar nature allows for a tentative reconstruction of the paleoclimatic conditions (Goudie, 1973; Allen, 1974). Carbonate pedogenesis, for example, normally requires a warm dry climate with a seasonably distributed, but not excessively peaked, rainfall. Conditions generally, however, do not appear to have been very favorable for the development of thick, extensive, mature pedogenic intervals as in some of the older Permian-Triassic "red bed" sequences in other parts of Gondwanaland (Turner, 1978). The presence of micrite in smear slides from these pedogenic intervals may be a result of pedogenically-induced carbonate precipitation and replacement within the vadose zone (Leeder, 1982).

The red color of the bulk of the sediments and their large clay content can be accounted for by the infiltration of clays into the porous and permeable sediments, followed by early diagenetic alteration of iron-bearing minerals such as mica, which releases a red ferric iron pigment (Walker, 1976). Thus, "red beds" are typically immature first-cycle sediments. Furthermore, clays may play an essential role in this reaction, which initially takes place under warm climatic conditions in the presence of adequate oxygenated ground water in the vadose zone. The close association of greenish gray reduced sediments shows that in some cases reddening did not occur, probably because of brief exposure to the oxygenated vadose zone. Local greenish gray reduction spots and mottles are a result of local concentrations of organic matter in sufficient quantities to resist oxidation.

The overall fining-upward trend of the cored sequence suggests a gradual reduction of the paleoslope through time and an increase in the amount of fine material supplied to the depositional site. This can be attributed to weathering, the gradual reduction in source area relief, and/or decreased tectonic activity. As a result, the depositional site may have been located progressively further away from the Antarctic source area during deposition of the upper part of the succession. Accordingly, deposition of the upper part may have been influenced, at least partially, by meandering rivers. Poor core recovery and the commonly unconsolidated nature of some of the sediments in the upper part of the unit make environmental interpretations speculative.

The clasts in Unit II probably represent a lodgement till, as evidenced by (1) the occurrence of similar clasts at about the same level at Sites 739 and 742, (2) the preferred orientation fabric recorded at Site 742, (3) the variability in downhole logging, suggesting interbedded sandy and bouldery sediment, and (4) the consolidation of the underlying diamictite below this unit at other sites, indicative of loading by ice. The sediments of Unit I indicate mainly glaciomarine deposition. The diatom ooze at the top of Unit I indicates that conditions at this time were more favorable for increased biogenic surface-water productivity, probably due to the absence of continuous ice cover. Pore-water chemistry shows (see "Inorganic Geochemistry" section, this chapter) that the silica concentrations in Unit I are very high in contrast to those in Unit II. They are also highly alkaline and emit a strong smell of hydrogen sulfide indicative of microbial activity and sulfate reduction.

BIOSTRATIGRAPHY

The paucity of both calcareous and siliceous microfossils in the samples examined from Holes 740A and 740B limits the biostratigraphic and chronostratigraphic determination of the sediments recovered at Site 740. Diatoms were seen in Cores 119-

740A-1R to 119-740A-3R, 119-740A-26R, and 119-740B-1R. Planktonic and benthic foraminifers were seen in Cores 119-740A-1R and 119-740B-1R. Calcareous nannofossils, radiolarians, and organic-walled microfossils were not seen in the samples examined.

The occurrence of abundant and well-preserved diatoms in core-catcher Samples 119-740A-1R and 119-740A-2R allows the interval from 0 to 13.7 mbsf to be assigned to the Holocene *Thalassiosira lentiginosa* Zone, suggesting an age younger than 0.62 Ma for this interval. Although the sparse and poorly preserved diatom assemblage in Sample 119-740A-3R-3, 150–152 cm, lacks zonal diatom taxa, the species present suggest a late Pliocene or younger age.

Rare specimens of *Nitzschia kerguelensis*, *Nitzschia angulata*, *Nitzschia cylindrus*, and *Thalassiosira gravis* were observed in Sample 119-740A-26R-CC. Although the occurrence of this assemblage suggests a Pliocene or younger age, these specimens are considered to result from contamination.

Foraminifers at Site 740 were studied in the mud-line sample and in most core-catcher samples. Benthic and planktonic species occur in the first core of each hole and are absent in the entire sequence below. The mud-line sample of Core 119-740A-1R contains the only relatively diverse foraminiferal assemblage, consisting entirely of arenaceous species that were not found to be preserved downcore. A living population of about 10%–20% indicates the recovery of the sediment/water interface. The most abundant species is *Miliammina arenacea*. The genus *Trochammina* is represented with a number of species. *Adercotryma glomerata*, a special indicator for cold water in the North Atlantic, has a common occurrence. Rare taxa are *Trochamminoides* sp., *Cribratomoides jeffreysi*, *Textularia* sp., *Reophax fusiformis*, *Reophax bilocularis*, *Thurammina papillata*, and *Psammosphaera fusca*. These agglutinated taxa generally show a fine-grained wall texture. Coarse-grained tubular forms and species of *Reophax*, which are representative of the mud-line sample at Site 739, are absent in this assemblage. The sediment is mainly a diatom ooze with a minor mineral component, which is not a substrate favored by arenaceous foraminifers. The glacial sediment from the first core-catcher sample contains only rare specimens of *Trifarina* sp. One specimen of the planktonic species *Neoglobobulimina pachyderma* was found in Sample 119-740A-1R-3, 145–150 cm.

PALEOMAGNETICS

One hundred forty-one discrete oriented paleomagnetic samples were collected from Cores 119-740A-3R and 119-740A-8R through 119-740A-31R. Core 119-740A-3R contains olive green diatom ooze and was normally polarized. The remaining cores consist of red clay, silts, and sandstones. The results of the natural remanent magnetization (NRM) measurements are shown in Figure 12. With the exception of brief intervals at 88, 150–160, 200, and 220 mbsf, the stratigraphic sequence appears to be magnetized with a negative inclination, indicating normal polarity within the Southern Hemisphere, based on the NRM results.

Half-core cryogenic magnetometer measurements were then made on Cores 119-740A-11R through 119-740A-31R after alternating field (AF) demagnetization at 5 mT. The cryogenic results confirmed the results of the discrete sampling and, in addition, located reversed intervals at 92 and 207 mbsf. In each case the zones of positive inclination (reversed polarity) were found to be very brief.

In order to better evaluate the magnetostratigraphy, both AF and thermal demagnetization experiments were conducted on a suite of 13 samples. In general, thermal demagnetization studies are used to study red sediments, because the carrier of magnetization is generally hematite, which is resistant to demagnetiza-

Figure 12. Stratigraphic plot of NRM sample inclination and intensity in Hole 740A. For the most part, the samples have negative inclinations on NRM measurement. The single-point reversals of polarity are marked by arrows. Reliable polarity reversals are characterized by more than one point of positive inclination.

tion by alternating fields. As would be expected, the AF and thermal demagnetization experiments yielded very different results. Examples of the AF demagnetization studies are shown in Figure 13. In general the samples show little change in direction after demagnetization up to 40 mT. Only small decreases in intensity were seen (between 10% and 40% of the normalized intensity). As expected, the samples studied using thermal demagnetization displayed very different magnetic behavior (Fig. 14). Samples 119-740A-19R-1, 126–128 cm, 119-740A-15R-1, 54–56 cm, and 119-740A-24R-1, 11–13 cm, clearly display three components of magnetization. The least stable component was removed at 100°C. The next component was removed in the range of 150° to 475°C, and a stable primary component remained between 475° and 580°C.

Based on the demagnetization experiments, stable magnetic directions apparently can be defined only at relatively high thermal demagnetization levels. Furthermore, these stable directions are characterized by low and commonly positive magnetic inclinations, ranging from -25, -15, +12, -14, -30, +21, +20, to +39.

During drilling, the age of these red units was discussed. The initial observations of dominantly normal polarity suggest that the rocks may be of Late Cretaceous age and, thus, could be re-worked deposits derived from the nearby continental red beds of the Antarctic continent. However, subsequent detailed thermal demagnetization studies have shown conclusively that two polarities are present, despite the limited number of samples of opposed polarity. Furthermore, the primary magnetization appears to be reversed rather than normal polarity. These observa-

Figure 13. Orthogonal plots, stereonet, and thermal demagnetization curves of magnetic behavior for samples after demagnetization from 0 to 40 mT. The samples show very little change in direction upon demagnetization and very little change in intensity. **A.** Sample 119-740A-29R-1, 17-19 cm. **B.** Sample 119-740A-11R-2, 12-14 cm.

tions are not inconsistent with a formation of these red beds in the Permian. Pollen studies will be completed on material from this site to examine this possibility. The multiple components of magnetization found in these rocks indicate that they have experienced a complex geologic history. The secondary components of magnetization could be related to subsequent geologic events resulting from structural tilting, regional tectonics related to the rifting of Gondwana, or the loading of the Antarctic continent by large ice masses in relatively recent time. Thorough, land-based paleomagnetic studies of these sediments will be necessary in order to accurately characterize the age and magnetic history of these rocks.

SEDIMENTATION RATES

The limited occurrence of calcareous and siliceous microfossils from the cores recovered at Site 740 prevents any chronostratigraphic or sediment-accumulation estimates to be completed for the interval below 13.7 mbsf. The interval stratigraphically above this depth (Cores 119-740A-1R and 119-740A-2R) is assigned to the *Thalassiosira lentiginosa* Zone and has an age younger than 0.62 Ma.

INORGANIC GEOCHEMISTRY

Two holes were cored in a water depth of about 800 m at Site 740 on the eastern Antarctic continental shelf in Prydz Bay. Hole 740A was cored to a depth of 226 mbsf, and Hole 740B sampled the mud line to a depth of 1.6 mbsf. The upper 57 mbsf consists of Pliocene(?) to Holocene glaciomarine sediments. The remainder of the section consists of fluvial sediments of probable Mesozoic or older age. A total of eight whole-round minicores 5-10 cm in length were obtained at Site 740 for the purpose of interstitial-water geochemical studies.

Site 740 provides the rare opportunity to study pore-water geochemistry in high-latitude sediments with mixed terrestrial and marine histories. Three lithologic units and four physical-properties units have been defined for Site 740 (see "Lithostratigraphy and Sedimentology" and "Physical Properties" sections). Two interstitial-water samples were taken from lithologic Unit I (0-23.2 mbsf, the same as physical-properties geotechnical unit G1), a diatom ooze with clayey silt, glacially-deposited clasts, and 70%-90% porosity. A 50% reduction in porosity occurs below Unit I. Because of poor core recovery, none of the

Figure 14. Orthogonal plots, demagnetization curves, and equal-area stereographic plots of magnetic behavior upon thermal demagnetization for pilot samples from Hole 740A. **A.** Sample 119-740A-19R-1, 126-128 cm. The orthogonal plots at left show that Samples **(B)** 119-740A-15R-1, 54-56 cm, and **(C)** 119-740A-24R-1, 11-13 cm, have a low-temperature component removed between 0° and 100°C. The demagnetization paths then display a gradual decay between 150° and 475°C indicating a consistent second component. A third component of magnetization present between 475° and 580°C decayed toward the origin of the plot and is interpreted as the primary stable component of magnetization. Large drops in the intensity occur in the area of 580°C, which suggests the magnetic mineral in these rocks is hematite.

glaciomarine sediments in lithologic Unit II (23.2–56.6 mbsf, the same as geotechnical unit G2) were sampled for interstitial waters. Six of the interstitial-water samples were taken from lithologic Unit III (56.6–225.5 mbsf, which coincides with geotechnical units G3 and G4), a Mesozoic(?) fluvial sequence with about 25% porosity. Below 190 mbsf (geotechnical unit G4), anomalous index properties indicate an increase in porosity with increasing depth. This increase in porosity may be due to an increase in grain size noted at the base of lithologic Unit III. The presence of mudstone beds and changes in the physical properties have important implications for the vertical and horizontal diffusion of dissolved chemical species in interstitial waters. All of the sediments analyzed for inorganic carbon contain less than 1% calcium carbonate. The organic carbon is less than 0.1% in all of the interstitial-water sediment samples, except for the two samples from lithologic Unit I, which contain 1.4% organic carbon (see “Organic Geochemistry” section, this chapter).

The eight interstitial-water minicore samples obtained at Site 740 are from cores recovered by the rotary core barrel (RCB) method. The probability of sediment disturbance and pore-fluid contamination is especially high when soft sediments are RCB cored. The sediments comprising lithologic Unit I have low bulk densities (1.22–1.64 g/cm³, see “Physical Properties” section) and experienced some drilling disturbance. Thus, the two interstitial-water samples taken from Unit I may have been contaminated with the surface seawater commonly pumped into the hole during RCB drilling. The remainder of the samples were taken from the denser sediments of lithologic Unit III (bulk density about 2.45 g/cm³) in which drilling contamination was less likely. Methods for recognizing the contamination of interstitial-water samples are briefly discussed for Site 739 (see “Inorganic Geochemistry” section, “Site 739” chapter). Charge-balance calculations were performed on each of the interstitial-water samples to check for gross irregularities in the data. Sodium and potassium concentrations were calculated by multiplying the ratio of each cation to chloride for average surface seawater, using the values given by Stumm and Morgan (1981), by the chloride concentration of each sample. The two interstitial-water samples from lithologic Unit I have charge imbalances of less than 0.4%, implying that contamination levels are extremely low or that the contaminant has a composition similar to these samples. More surprisingly, the six samples from lithologic Unit III have large charge imbalances of 9.8%–17.4%, with the imbalance tending to increase with increasing depth. The chloride concentrations of these samples are much lower than that of most seawater (mean = 508 mmol/L, s.d. = 13 mmol/L vs. 550 mmol/L for average surface seawater), implying that a low-salinity water source is mixing with these interstitial waters. The two most obvious low-salinity sources are the drilling muds used during RCB drilling (see “Inorganic Geochemistry” section, “Site 739” chapter, for drill water composition) and ground waters from the Antarctic continent. A

third possibility, which is not supported by organic geochemistry analyses, is that the samples were diluted by fresh water derived from gas hydrate decomposition (see “Organic Geochemistry” section).

Site 740 interstitial-water samples were analyzed for pH, alkalinity, salinity, chloride, magnesium, calcium, sulfate, phosphate, ammonium, and silica using the methods outlined in the “Explanatory Notes” chapter. Not all of the samples were large enough to allow the full complement of analyses. All of the data obtained at Site 740 are presented in Table 4.

Salinity and Chloride

The salinity and chloride concentrations (Fig. 15) of the Site 740 interstitial-water samples decrease sharply below the diatomaceous oozes of lithologic Unit I. The two interstitial-water samples taken from Unit I have chloride concentrations (mean = 545 mmol/L) similar to those of intermediate ocean waters sampled in the area and probably reflect the composition of seawater near the seafloor. The high alkalinities, high phosphate, ammonium, and silica concentrations of these samples indicate that no significant drilling contamination occurred. The six samples from the terrestrial sediments of Unit III have chloride concentrations (mean = 508 mmol/L, s.d. = 13 mmol/L) 7% lower than those in the Unit I samples. The small increase in sulfate concentrations below 72 mbsf indicates that some drilling contamination may have occurred in these samples. However, the low sulfate concentration (0.9 mmol/L) of Sample 119-740A-11R-1, 140–150 cm, implies that it has not been contaminated, and its chloride concentration of 512 mmol/L suggests that the lower chloride values are characteristic of the *in-situ* interstitial waters. The lack of methane in these sediments indicates that interstitial waters were not diluted by gas hydrate decomposition. The close proximity of Site 740 to the Antarctic continent promotes the possibility of mixing between ground waters and ocean water as the source of these interstitial waters. The abrupt decreases in salinity and chloride concentrations between Units I and III are indicative of the very different densities, porosities, and diffusion coefficients in these lithologic units.

Magnesium and Calcium

Good correlation ($r = -0.974$) occurs between dissolved magnesium and dissolved calcium concentrations in the interstitial-water samples from Site 740 (Figs. 15 and 16). Magnesium and calcium concentrations in the two samples extracted from lithologic Unit I are similar to seawater concentrations. Dissolved magnesium concentrations decrease 75% between Samples 119-740A-3R-2, 145–150 cm, at 17 mbsf (52.0 mmol/L) and 119-740A-31R-6, 0–10 cm, at 223 mbsf (13.1 mmol/L), whereas dissolved calcium concentrations increase by 8.4% between the same two samples. The good correlation between dissolved magnesium and calcium concentrations implies that these cations behave conservatively at Site 740, which means that their concentrations are regulated by diffusion properties between

Table 4. Interstitial-water geochemical data, Site 740.

Core, section, interval (cm)	Depth (mbsf)	Volume (mL)	pH	Alkalinity (mmol/L)	Salinity (g/kg)	Magnesium (mmol/L)	Calcium (mmol/L)	Chloride (mmol/L)	Sulfate (mmol/L)	Phosphate (μmol/L)	Ammonium (mmol/L)	Silica (μmol/L)	Mg ²⁺ /Ca ²⁺
119-740B-													
1R-1, 145–150	1.45	70	7.6	23.19	35.0	50.2	8.9	549	12.8	152	2.04	700	5.6
119-740A-													
3R-2, 145–150	16.65	50	8.0	20.15	35.0	52.0	9.0	541	15.4	130	2.16	653	5.8
8R-3, 140–150	61.00	2	—	—	32.5	41.9	35.2	507	15.0	—	0.34	134	1.2
11R-1, 140–150	72.30	11	7.9	0.69	31.5	27.8	41.6	512	0.9	0	0.78	38	0.7
17R-1, 140–150	101.30	12	7.7	0.94	31.8	24.5	55.8	508	2.6	0	0.72	—	0.4
22R-1, 140–150	130.20	2	—	—	32.2	19.2	64.5	517	1.7	—	0.51	106	0.3
25R-1, 140–150	159.20	8	7.5	0.87	33.8	18.9	75.5	519	8.6	0	0.43	106	0.2
31R-6, 0–10	223.30	7	7.4	1.07	30.5	13.1	76.0	482	5.1	0	0.24	49	0.2

Figure 15. Interstitial-water profiles, Site 740.

Figure 16. Calcium carbonate vs. magnesium ion concentration plot for Site 740 interstitial waters. The regression line passes through data points representing interstitial-water samples in the upper 223 mbsf and has a slope of -1.74 ($r = -0.974$).

sources and sinks (see McDuff and Gieskes, 1976; McDuff, 1978; Gieskes, 1983). The relatively high slope (-1.74) of the regression line plotted through the calcium and magnesium data in Figure 17 indicates that diffusion rates are low within the terrestrial sediments of lithologic Unit III.

The source of magnesium in the interstitial waters is seawater, and a likely sink is the formation of magnesium silicates during the alteration of unstable crustal rocks. Much of the calcium may be produced by means of alteration reactions that consume magnesium. Most of this calcium is presumably lost to the ocean. The good correlation between magnesium and calcium in this data set provides further evidence against significant drilling contamination.

Silica

Similar to chloride concentrations, the silica concentrations (Fig. 15) in the interstitial-water samples from Site 740 decrease sharply below lithologic Unit I. Chloride and silica concentrations are well correlated in this data set ($r = 0.856$). Because chloride ions should behave conservatively in these pore fluids, the sharp drop in silica concentrations reflects the sharp change in the diffusion coefficient between Units I and III rather than a reaction zone in the sediment column. The relatively high silica concentrations in the two interstitial-water samples from Unit I are due to the dissolution of biogenic silica in the sediments. Much of the silica probably is lost to the ocean. The rise in silica concentrations in the samples below 72 mbsf may be due to contamination with drilling muds. Dissolution and reprecipitation reactions involving relatively insoluble silicates may be responsible for the low dissolved silica concentrations below 61 mbsf.

Figure 17. Calcium carbonate and total carbon profiles, Site 740.

pH, Alkalinity, and Sulfate

Alkalinity and sulfate analyses (Fig. 15) of interstitial waters from Site 740 indicate that high levels of microbial catabolism in the upper sediment layers have resulted in anaerobic conditions below the sediment/water interface. Oxidation of abundant reactable organic matter (around 1.4% organic carbon) has exhausted the dissolved oxygen and nitrate, and sulfate reduction has consumed about 50% of the available sulfate in the first 1.5 mbsf. Samples 119-740B-1R-1, 0–10 cm, at 1 mbsf and 119-740A-3R-2, 145–150 cm, at 17 mbsf had a strong hydrogen sulfide odor. Sedimentation rates and the supply of organic matter have been high enough at this site to allow the consumption of nearly all of the sulfate at 72 mbsf (0.9 mmol/L). The alkalinity of the mud-line sample from Hole 740B is about 10 times greater than average seawater. By 72 mbsf, the alkalinity has dropped to very low levels (0.69 mmol/L), indicating that alkalinity is being precipitated at deeper levels and/or that much of it diffuses into the overlying water column. The rapid increase in sediment bulk density below 17 mbsf may present an effective barrier to downward diffusion that allows much of the alkalinity to escape the sediment system.

Ammonium and Phosphate

Both ammonium and phosphate are significantly enriched in interstitial waters near the seafloor at Site 740 (Fig. 15). Most of the ammonium and phosphate in interstitial waters is produced during the oxidation of organic matter. The relatively high ammonium and phosphate concentrations of Samples 119-740A-3R-2, 145–150 cm, at 17 mbsf (ammonium 2.16 mmol/L and phosphate 130 mmol/L) and 119-740B-1R-1, 0–10 cm, at 1.4 mbsf (ammonium 2.04 mmol/L and phosphate 152 mmol/L) indicate that much of the bacterial catabolism occurs near the

sediment/water interface. The gradual depletion of ammonium with increasing depth may be due to its incorporation in clay minerals in deeper levels of the sediment column. The lack of detectable amounts of phosphate at 61 mbsf and below supports the hypothesis that much of the byproducts of organic matter degradation is lost to the water column.

ORGANIC GEOCHEMISTRY

Organic geochemistry was studied on squeeze cakes of interstitial-water samples from Hole 740A (0–225.5 mbsf) as outlined in the “Explanatory Notes” chapter.

Hydrocarbon Gases

The headspace procedure was used approximately every 30 m to determine hydrocarbon gases. Most of the samples had methane levels near or below the mean laboratory air level of 3 ppm. Sample 119-740A-14R-3, 0–1 cm, had 8 ppm ethane and 1140 ppm methane. This sample was an isolated pocket of probable biological origin, as the previous gas samples all had less than 40 ppm methane and no ethane. Samples from below dropped quickly down to less than 40 ppm methane. No other ethane was detected.

Carbon Analysis

Inorganic carbon was measured on most core-catcher samples and all of the interstitial-water squeeze cakes. Total carbon was measured on all of the interstitial-water squeeze cakes and most of the core-catcher samples. Total organic carbon (TOC) was calculated (see “Explanatory Notes” chapter) for samples in which total carbon was determined (Table 5 and Fig. 17).

Site 740 contained little organic matter except from the top of the hole to 17 mbsf, with 1.4% TOC. The levels of organic matter dropped to <0.1% downhole, but no data are available between 17 and 59 mbsf for interstitial-water or TOC determinations because of low core recovery. This lack of information makes assessing the processes responsible for the rapid downhole depletion of organic matter difficult. The uppermost sequence is Holocene in age, and below approximately 60 mbsf it may be older than Cretaceous. The sequence probably was deposited in an arid environment. The downhole depletion is possibly due to this hiatus.

BIOLOGY AND OCEANOGRAPHY

Physical Characteristics of Marine Ecosystem

Moderate east to northeast winds dominated from 24 to 27 January during the occupation of Site 740. The air was clear, with many hours of sunlight and visibility >80 km (see Table 6). Air temperature stayed above freezing, and the sea-surface temperature varied from +0.2° to +1.9°C at the drillship. The surface water was mixed down to 20 m during the first day and became even more shallow the next day. During the second and third days, the wind shifted from the northeast to southeast and then back again, which brought some warmer, lower salinity water into the top 10 m (Fig. 18). On the final day a stepwise thermocline developed, with near-surface waters again appearing homogeneous. The mixed depth was shoaling as the water heated from the long hours of sunlight. The water looked green, with the Secchi disc disappearing at 4 m, giving a euphotic zone of only 10.8 m for three of the days. On the third day when somewhat clearer, lower salinity water intruded, the euphotic zone deepened slightly to 13.5 m.

Phytoplankton

A phytoplankton time series was completed over 3 days of sampling during which net hauls and chlorophyll measurements showed sizeable increases. Chlorophyll showed moderately high

Table 5. Total carbon, inorganic carbon, organic carbon, and carbonate carbon, Site 740.

Core, section, (interval) (cm)	Depth (mbsf)	Total carbon (%)	Inorganic carbon (%)	Organic carbon (%)	CaCO ₃ (%)
119-740A-					
1R-1, 60–63	0.60		0.02		0.2
1R-CC, 0–1	0.86		0.02		0.2
119-740B-					
1R-1, 145–150	1.45	1.41	0.01	1.40	0.1
119-740A-					
2R-1, 22–25	6.22		0.01		0.1
2R-CC, 0–1	6.61		0.06		0.5
3R-1, 120–121	14.90		0.03		0.3
3R-2, 50–51	15.70		0.04		0.3
3R-2, 145–150	16.65	1.39	0.02	1.37	0.2
3R-3, 106–107	17.76		0.04		0.3
3R-4, 4–5	18.24		0.02		0.2
8R-2, 70–73	58.80		0.01		0.1
8R-3, 85–88	60.45		0.01		0.1
8R-3, 140–150	61.00	0.08	0.01	0.07	0.1
9R-1, 10–13	61.30		0.01		0.1
8R-4, 25–27	61.35		0.00		0.0
8R-CC, 0–1	61.46		0.02		0.2
9R-CC, 0–1	61.76		0.02		0.2
10R-1, 46–47	66.66		0.01		0.1
10R-2, 90–93	68.60		0.01		0.1
11R-1, 73–76	71.63		0.00		0.0
11R-1, 140–150	72.30	0.05	0.02	0.03	0.2
11R-2, 65–67	73.05		0.01		0.1
11R-CC, 0–1	73.12		0.01		0.1
12R-1, 10–13	76.00		0.01		0.1
12R-1, 78–81	76.68		0.01		0.1
12R-CC, 0–1	77.13		0.01		0.1
13R-1, 26–29	80.86		0.01		0.1
13R-1, 100–102	81.60		0.00		0.0
14R-1, 37–38	85.97		0.01		0.1
14R-1, 117–118	86.77		0.01		0.1
14R-2, 81–82	87.91		0.00		0.0
14R-2, 140–150	88.50	0.06	0.00	0.06	0.0
14R-3, 30–31	88.90		0.00		0.0
15R-1, 48–49	90.68		0.00		0.0
15R-1, 145–147	91.65		0.00		0.0
17R-1, 63–64	100.53		0.01		0.1
17R-1, 66–67	100.56		0.01		0.1
17R-1, 140–150	101.30	0.04	0.02	0.02	0.2
17R-2, 9–10	101.49		0.02		0.2
17R-CC, 0–1	101.56		0.03		0.3
19R-1, 16–20	119.36		0.10		0.8
19R-1, 107–109	120.27		0.01		0.1
21R-1, 5–7	126.05		0.01		0.1
20R-3, 78–90	128.58		0.01		0.1
22R-1, 48–52	129.28		0.01		0.1
20R-4, 18–21	129.48		0.01		0.1
22R-1, 140–150	130.20	0.04	0.01	0.03	0.1
22R-2, 60–63	130.90		0.01		0.1
22R-3, 52–53	132.32		0.01		0.1
23R-1, 29–32	138.79		0.01		0.1
23R-1, 72–76	139.22		0.02		0.2
24R-1, 71–73	148.81		0.01		0.1
24R-2, 1–3	149.61		0.01		0.1
25R-1, 3–6	157.83		0.01		0.1
25R-1, 140–150	159.20	0.04	0.00	0.04	0.0
25R-2, 17–20	159.47		0.00		0.0
26R-4, 64–65	172.64		0.01		0.1
27R-2, 31–32	178.91		0.00		0.0
27R-3, 12–13	180.22		0.01		0.1
28R-1, 127–128	188.07		0.01		0.1
28R-1, 140–150	188.20	0.07	0.01	0.06	0.1
29R-2, 18–19	198.18		0.03		0.3
30R-1, 113–114	207.23		0.02		0.2
31R-3, 45–48	219.25		0.02		0.2
31R-5, 133–135	223.13		0.01		0.1
31R-6, 0–10	223.30	0.04	0.01	0.03	0.1

Table 6. Wind and weather summary for Site 740 (68.7°S, 76.8°E), January 1988.

24 Jan/00Z-2359Z:	
Wind	: East-northeast 8-12 kt.
Sea-surface temperature	: 0.5°-1.5°C.
Waves	: 1-1.5 m.
Sky	: Clear, gradually becoming mostly cloudy late in the period. Visibility > 80.5 km. Pressure rising slowly.
Air temperature	: Maximum: 5.7°C at 12Z; minimum: 0.1°C at 00Z.
25 Jan/00Z-2359Z:	
Wind	: Northeast 14-18 kt, shifting to southeast 8-12 kt around 09Z then back to northeast 10-15 kt about 19Z.
Sea-surface temperature	: 0.4°-1.6°C.
Waves	: 0.5-1 m.
Sky	: Partly to mostly cloudy, becoming overcast with light snow by 08Z. Clearing by 18Z. Visibility > 80.5 km. Pressure slowly rising to steady.
Air temperature	: Maximum: 4.3°C at 12Z; minimum: 1.3°C at 00Z.
26 Jan/00Z-2359Z:	
Wind	: East-northeast 7-17 kt, shifting to southeast 6-12 kt by 10Z then back to the northeast 10-15 kt by 20Z.
Sea-surface temperature	: 0.2°-1.0°C.
Waves	: 0.5 m.
Sky	: Clear to partly cloudy. Visibility > 80.5 km. Pressure rising.
Air temperature	: Maximum: 3.1°C at 12Z; minimum: 0.5°C at 00Z.
27 Jan/00Z-1500Z:	
Wind	: Northeast 15 kt, decreasing to 5-7 kt late in the period.
Sea-surface temperature	: 0.6°-1.9°C.
Waves	: 0.5-1 m.
Sky	: Clear. Visibility > 80.5 km. Pressure rising then falling late.
Air temperature	: Maximum: 2.5°C at 12Z; minimum: 0.4°C at 06Z.

Note: Due to local effects, each 24-hr day is summarized.

maxima between 25 and 33 m, well below the 1% light level. Opening and closing net hauls down to 200 m were crowded, resembling a thick green soup. A trip by *Maersk Master* to the pack ice near shore allowed the collection of seven brown ice samples, taken from about 1 m above sea level.

Dominant in the vertical net hauls and surface net tows were *Corethron criophilum* and *Corethron inerme*, both in auxospore formation, as noted at Site 739. For the first time these small (5 µm) microspores were seen externally (G. Fryxell, pers. comm., 1988), clustered around the girdle of another cell (net haul, 200-0 m). However, many cells were seen that were empty or did not show autofluorescence of active chlorophyll (Table 7).

Some of the species were reaching peak numbers, with long chains in such genera as *Chaetoceros*, *Thalassiosira*, and *Eucampia*. The onset of resting spore and winter form production was noted especially below 40 m. These have heavy cell walls and can often sink to the sediments, where life information adds to ecological interpretation. Winter forms were observed in *Eucampia antarctica* (doublets with straight margins in broad girdle view), especially in 80- to 40-m net hauls. Similarly, resting spores were commonly seen in *Chaetoceros neglectus* in 200- to 100-m net hauls, and in *Chaetoceros* sp. cf. *wighamii* in 80- to 40-m and 200- to 100-m net hauls. The *Chaetoceros* resting spores were in abundance (and still pigmented) in fecal pellets from all depths, not only the depths where they were found forming.

Species found in the brown ice are listed in Table 8. This is a preliminary listing, with identifications made from water mounts and based on shape, size, and colony type only. Although this is an effective means of identifying *Chaetoceros*, for example, it is not at all effective for *Nitzschia* species, especially those that form ribbon colonies. A more rigorous identification and estimate of relative abundance is planned for a shorebased study. The preliminary table is included here to indicate that no

one species is found only in ice, but that those species adapted to growth in brine pockets flourish. The pieces of ice probably were not formed *in situ*, but were rafted in from diverse sites and further represent several inoculations with surface seawater. Nevertheless, few species in the plankton at Site 740 are absent in the ice (Table 9).

Some general statements can be made concerning the ice communities, despite differences in the history of each sample. Brine pockets and channels are habitats for communities with recycling (and not only assemblages with little interaction), as indicated by the presence of fecal pellets of small grazers. Greater numbers of delicate, small choanoflagellates (bacteriovores) can be expected with careful shorebased study. Resting spores are not commonly formed in the ice, although spores can be frozen into the ice. Although the flora and relative abundance varies from one ice flow to the next, as seen even in this small sampling, the flora is commonly dominated by small needle-shaped cells and ribbon colonies of the genus *Nitzschia*. *Amphiprora* sp., *Chaetoceros neglectus*, and *Tropidoneis* spp. were seen frequently also. Large centric diatoms, such as *Thalassiosira tumida*, formed chains in all of the samples. Although their numbers were not high, they appeared viable.

These samples were collected about 1 m above the water line and were subject to extremes of temperature, light, and salinity. When diatoms are capable of auxospore production, such stresses can trigger sexual reproduction in cultures. Either auxospores or the post-auxospore stages were seen in both *Corethron* species, *E. antarctica*, and *Rhizosolenia alata*.

Other Observations of Productivity

Prydz Bay provided a location for an interdisciplinary project during a productive season. For instance, when *Maersk Master* cleaned the water intake filter screen at Site 740, it was clogged with *Euphausia superba* (krill). The seawater filter system on *JOIDES Resolution* also had a large number of krill.

When the camera descended after the drill bit was replaced, swarms of notothenioid fish followed it down from about 500 to 700 m (on tape). This productivity was supported by phytoplankton growth, and the shipboard chemists reported that the silica in surface water was being depleted. *Phaeocystis*, a prymnesiophyte, was also abundant, although it did not show much autofluorescence below 100 m. The sediment traps deployed (50, 100, 200 m) for three 24-hr periods provided one index of the amount of grazing taking place, with rapid transport of fecal pellets out of the surface layers even during the short-term deployment. Most of these fecal pellets showed very little autofluorescence, but *Chaetoceros neglectus* resting spores showed good color. Some fecal pellets had mostly broken cells (previously noted in krill stringlike fecal pellets), but even in these the *Chaetoceros* resting spores were commonly intact and showed autofluorescence of chlorophyll.

Diatoms in the Surface Sediment

Two mud-line samples, from Holes 740A and 740B, allowed comparison of the flora in the water column and the death assemblage. The sediments are dominated by the heavily silicified species of *Nitzschia* that form ribbon colonies and are associated with ice (Table 10). Quite delicate valves of *Asteromphalus*, with thin hyaline areas, were well represented, although some are fragmented. Heavily silicified forms of *T. tumida* and *Thalassiosira ritscheri* were present. *T. tumida* makes heavier valves in water below 0°C (Fryxell, 1988), and these were occasionally seen in the deeper net hauls. Since heavily silicified *T. ritscheri* cells were seen in the 200- to 100-m haul, and its resting spores are not known, it may have a cold-water or winter form. Comparison of the lightly and more heavily silicified forms may show similar patterns.

Only the straight-sided winter forms of *E. antarctica* were seen in the sediment, suggesting that the asymmetric form seen in the northern sites of the Kerguelen Plateau does not reach this latitude. Since doublet winter forms have been seen frequently and called resting spores (Hoban et al., 1980; Fryxell, in press), one would expect that the ratio of valves with flattened elevations (intercalary valves in chains) to those with pointed elevations (end valves of chains) would be 1-1. Instead, the ratio was close to 2-1 (Table 11). Possible explanations include the following:

1. Instead of only doublet winter stage cells, doublets may continue to divide during the winter as conditions allow growth.
2. The winter stage transition is truncated 50% of the time, perhaps by death of the cells or by grazing, when only the first-formed resting valves (with flattened elevations) are completed.
3. The pointed elevations on terminal valves are more subject to breakage and dissolution.

Testing the validity of these explanations will require study of both the plankton and the sediments.

PHYSICAL PROPERTIES

The main objectives of the physical-properties program at Site 740 were to study downhole variations in geotechnical parameters and relate this to different lithologies and main phases in the glacial history of eastern Antarctica. At Site 739, cyclic changes in the physical properties were found, and Site 740 was considered important for comparison and possible correlation, as this was a much more glacier proximal site, only 30 km from the present-day coast.

Physical properties measured were (1) index properties, (2) undrained shear strength, and (3) compressional-wave velocity. A discussion of the sampling and laboratory techniques used at

Site 740 is in the "Explanatory Notes" chapter. Both Holes 740A and 740B were rotary drilled. Hole 740A was cored to 225 mbsf. Hole 740B, a mud-line core, penetrated to 1.6 mbsf. Physical-properties measurements were taken entirely from Hole 740A, which had only 32% core recovery. The well-consolidated sediments found below 17 mbsf were probably not significantly disturbed by the rotary procedure. However, in view of the rather low recovery, the physical-properties data presented may not be entirely representative of the complete sedimentary section penetrated at Site 740.

Results

As at previous Leg 119 sites, physical-properties profiles enable subdivision of the sedimentary section into distinct geotechnical units. Changes in the range of values, overall character, and trends in the property depth profiles serve to distinguish three geotechnical units, G1 to G3, at Site 740. Because of the poor core recovery, exact unit boundaries were difficult to define, and the boundaries were placed tentatively midway between the two nearest data points. All physical properties measured are presented in Figure 19 and Table 12. Table 13 shows the range and mean values for the different parameters within each geotechnical unit.

Geotechnical unit G1 (0-16.7 mbsf) covers most of lithologic Unit I and is composed of normally consolidated diatom oozes and clayey silt (see "Lithostratigraphy and Sedimentology" section). Although based on few data points because of the poor core recovery, this unit can be characterized as having relatively high, but variable, water content (45%-75%), high porosity (70%-88%), low bulk density (1.22-1.64 g/cm³), low sonic velocity (1495 m/s), and low undrained shear strength (<5 kPa).

Geotechnical unit G2 (16.7-193.5 mbsf) covers lithologic Unit II and most of Unit III, consisting of alternating sandstones, siltstones, and claystones of continental "red-bed" type (see "Lithostratigraphy and Sedimentology" section). It is distinguishable from the overlying sediments by an abrupt and large decrease in water content, combined with associated increases in bulk density, velocity, and shear strength.

Poor core recovery limited the data from the upper part of the unit. From approximately 57 mbsf (Core 119-740A-8R), the unit shows slight, but stable, normal depth trends. Water content varies from 8.5% to 17.3%, averaging 13.1%; corresponding porosity values are 20.0%, 35.2%, and 28.8%, and bulk densities are 2.22, 2.70, and 2.40 g/cm³. GRAPE densities are somewhat lower, but show the same trends, averaging 2.35 g/cm³. Reasons for the discrepancy in the differently measured bulk densities are discussed for Site 739 (see "Physical Properties" section, "Site 739" chapter). Compressional-wave velocity was measured both parallel (C direction) and normal (A direction) to the bedding on a number of samples. Both values show, despite a considerable scatter, an increase with depth, with the greatest increase in the upper half of the unit. Average velocities for the A and C directions are 2401 and 2392 m/s, respectively. No apparent anisotropy was seen. Values of undrained shear strength show a sharp increase at the upper boundary of unit G2. Farther downsection, the values are few and highly scattered, varying between 7.7 and 650 kPa. Because of diagenetic effects and drilling disturbance, these values are considered unreliable.

Geotechnical unit G3 (193.5-225.5 mbsf, total depth) also covers the lower part of lithologic Unit III. While average values and ranges are similar to those of unit G2, there are slightly, but definitely, reversed from normal to anomalous depth trends in index properties and velocity profiles. Bulk density, possibly GRAPE density, and velocity show decreasing trends below this boundary, whereas water content and porosity increase.

Figure 18. Conductivity, temperature, and depth (CTD) profiles down and up in the top 200 m of the water column in the vicinity of the drill site. The top 20 m are removed because of the noise in the signal until the self-contained unit came to equilibrium *in situ*. A. CTD-8: 24 January 1988, operation 279, 0828–0850 hr (local time), 68°41.5'S, 76°39.5'E. B. CTD-9: 25 January 1988, operation 286, 0930–0944 hr (local time), 68°39.0'S, 76°35.0'E. C. CTD-10: 26 January 1988, operation 295, 1618–1628 hr (local time), 68°41.0'S, 76°44.0'E. D. CTD-11: 27 January 1988, operation 305, 1048–1058 hr (local time), 68°42.17'S, 76°39.11'E. The surface mixed layer became more shallow before the introduction of less saline water, and a warming trend in the surface water is apparent.

Table 7. Life stage and condition of *Corethron*: *Maersk Master* operation 298, 26 January 1988, with replicate 100- to 200-m net hauls.

Condition	Depth below sea surface (m)		
	0-50	100-200	100-200
Vegetative cells:			
Empty (%)	25	71.1	63.8
Autofluorescing (%)	72.9	14.4	25.6
Auxospores:			
Empty (%)	1.0	12.4	10.8
Autofluorescing (%)	1.0	2.1	—
Total cells counted	194	194	258

Table 8. Preliminary list of diatoms and other plankton found in separate samples of brown ice, near the eastern shore of Prydz Bay (*Maersk Master* operation 293).

Organism	Sample number						
	1	2	3	4	5	6	7
<i>Actinocyclus</i>	—	R	R	F	R	R	R
<i>A. actinochilus</i>	R	—	—	F	—	R	R
<i>A. spiritus</i>	F	F	F	R	R	C	F
<i>Amphiprora</i> sp.	—	—	—	—	—	—	—
<i>Asteromphalus</i>	—	R	R	—	—	—	R
<i>A. hookeri</i>	—	—	—	R	R	—	R
<i>A. hyalinus</i>	—	—	—	R	R	—	R
<i>A. parvulus</i>	—	—	R	—	—	R	—
<i>Chaetoceros</i>	—	—	—	—	—	—	—
<i>C. bulbosus</i>	R	—	—	—	—	—	—
<i>C. convolutus</i>	—	—	R	—	—	R	—
<i>C. criophilus</i>	—	—	R	—	R	R	—
<i>C. curviretus</i>	—	—	F	F	R	F	—
<i>C. dictyota</i>	R	—	R	R	R	R	R
<i>C. neglectus</i>	F	F	C	C	C	C	C
<i>C. neogracile</i>	—	R	R	R	—	—	—
<i>C. pacificus</i>	—	—	—	—	—	C	—
<i>Corethron</i>	—	—	—	—	—	—	—
<i>C. criophilum</i>	—	R	R	—	R	R	R
<i>C. inerme</i>	—	R	R	R	R	R	R
<i>Coscinodiscus</i>	—	—	—	—	—	—	—
<i>C. oculoides</i>	R	—	R	R	R	—	—
<i>Dactyliosolen</i>	—	—	—	—	—	—	—
<i>D. antarcticus</i>	R	—	—	R	R	R	—
<i>D. tenuijunctus</i>	—	—	—	—	—	—	R
<i>Eucampia</i>	—	—	—	—	—	—	—
<i>E. antarctica</i>	F	F	F	F	F	F	F
<i>Haslea</i> sp.	—	—	R	—	—	—	—
<i>Navicula</i>	—	—	—	—	—	—	—
<i>N. directa</i>	—	R	R	R	—	—	R
<i>Nitzschia</i>	—	—	—	—	—	—	—
<i>N. angulata</i>	R	F	R	—	—	—	R
<i>N. closterium</i>	C	C	C	C	C	C	C
<i>N. curta</i>	C	C	C	A	A	A	C
<i>N. cylindrus</i>	A	C	A	A	A	A	C
<i>N. heimii</i>	—	—	—	—	—	R	—
<i>N. inflatula</i>	F	—	C	C	—	F	—
<i>N. lecontei</i>	F	F	C	—	—	—	C
<i>N. lineola?</i>	R	—	—	—	F	R	—
<i>N. prolongatoides</i>	—	—	—	F	C	C	C
<i>N. ritscheri</i>	R	—	—	—	F	R	F
<i>N. subcurvata</i>	—	—	C	C	C	—	C
<i>N. turgidula</i>	—	—	R	—	F	—	F
<i>N. turgiduloides?</i>	—	—	R	F	—	—	—
<i>N. spp.</i>	—	—	—	—	—	—	—
(ribbon colonies)	C	A	A	A	A	A	A
<i>Odontella</i>	—	—	—	—	—	—	—
<i>O. weissflogii</i>	—	R	—	—	—	—	R

Table 8 (continued).

Organism	Sample number						
	1	2	3	4	5	6	7
<i>Porosira</i>	—	—	—	—	—	R	—
<i>P. glacialis</i>	R	R	R	F	F	F	—
<i>P. pseudodenticulata</i>	—	—	—	—	—	—	—
<i>Rhizosolenia</i>	—	—	—	—	—	—	—
<i>R. alata</i>	F	R	R	—	F	F	R
<i>R. hebetata</i>	—	—	—	—	—	—	—
<i>f. semispina</i>	—	—	—	—	—	—	R
<i>Stellarima</i>	—	—	—	—	—	—	—
<i>S. microtrias</i>	R	—	R	R	R	—	R
<i>Synedra</i>	—	—	—	—	—	—	—
<i>S. reinboldii</i>	R	—	—	—	—	R	R
<i>S. sp. (small)</i>	—	—	R	R	—	R	—
<i>Thalassiosira</i>	—	—	—	—	—	—	—
<i>T. gravida</i>	R	—	—	—	—	—	—
<i>T. lentiginosa</i>	—	—	R	—	—	—	—
<i>T. ritscheri</i>	—	—	R	—	—	—	—
<i>T. tumida</i>	—	—	—	—	—	—	—
(ice-related form)	F	F	F	F	F	F	F
<i>T. sp. (small)</i>	—	R	R	—	—	R	—
<i>Tropidoneis</i>	—	—	—	—	—	—	—
<i>T. belgicae</i>	F	F	R	F	F	F	C
<i>T. glacialis</i>	R	F	R	F	F	F	C
Others:	—	—	—	—	—	—	—
<i>Distephanus speculum</i>	—	—	R	R	R	R	—
<i>Phaeocystis</i>	C	C	A	A	A	A	A
Spiny cysts	—	R	—	A	R	R	R
Choanoflagellates	—	R	—	—	—	—	—
Fecal pellets	F	F	F	F	F	F	F

Table 9. Planktonic species at Site 740 in Prydz Bay but not noted in ice samples (preliminary listing).

Chaetoceros sp. cf. *wighamii*
C. sp. cf. dictyota
C. flexuosus
Nitzschia kerguelensis
N. obliquecostata
N. sublinearis
Rhizosolenia chunii

Discussion

The variability of index properties suggested by the few data points of unit G1 is probably real rather than simply a consequence of the limited sample points. Diatomaceous sediments typically show a large scatter, and a more diatom-poor clayey silt (lithologic Subunit IB) is also present.

The abrupt nature of the contact between units G1 and G2 is indicated on all physical-properties profiles. This is probably as much a consequence of loading history as of change in lithology. Although biostratigraphic control is poor below the upper unit, the site has likely been subject to extensive erosion by grounded glacier ice during several periods of an expanded eastern Antarctic ice sheet. Hence, unit G2 has experienced excess loading and is overconsolidated, while unit G1 probably was deposited post-glacially. The very steep gradient indicated by the continuous GRAPE data confirms the sharp nature of the contact, with a very rapid increase to high values.

Table 10. Species found in mud-line Samples 119-740A-1R-1, 0 cm, and 119-740B-1R-1, 0 cm (preliminary listing).

<i>Actinocyclus actinochilus</i>
<i>A. spiritus</i>
<i>Asteromphalus hyalinus</i>
<i>A. hookeri</i>
<i>A. parvulus</i>
<i>Chaetoceros</i> sp. cf. <i>wighamii</i> , resting spores
<i>C. bulbosus</i> (bulbous, closed setae)
<i>C. convolutus</i>
<i>C. criophilus</i>
<i>C. dichæta</i>
<i>C. neglectus</i> , resting spores
<i>Cocconeis imperatrix</i>
<i>Corethron</i> sp.
<i>Coscinodiscus oculoides</i>
<i>Dactyliosolen antarcticus</i> (heavy band)
<i>Eucampia antarctica</i> (straight winter form, pointed and flattened elevations)
<i>Denticula?</i> sp. (curved valve)
<i>Nitzschia angulata</i>
<i>N. curta</i>
<i>N. cylindrus</i>
<i>N. kerguelensis</i>
<i>N. obliquecostata</i>
<i>N. separanda</i>
<i>N. sublinearis</i>
<i>Odontella weissflogii</i>
<i>Porosira glacialis</i> (resting spore)
<i>P. pseudodenticulata</i>
<i>Rhizosolenia hebetata</i> f. <i>semispina</i> also <i>bidens</i> form
<i>Rouxia?</i> sp.
<i>Stellarima microtrias</i> , resting spore
<i>Thalassiosira gracilis</i>
<i>T. lentiginosa</i>
<i>T. oliverana</i>
<i>T. ritscheri</i> (heavy form—resting spore? cold-water form?)
<i>T. tumida</i> (heavy, linear, cold-water form)
<i>T. sp. cf. grævida</i> (heavy form—resting spore??)

Table 11. Proportion of *Eucampia* winter form valve types in mud-line Sample 119-740B-1R-1, 0 cm.

Slide no.	Elevations flattened	Elevations pointed
1	99	45
2	80	33
3	33	9
4	35	18
5	50	22
6	84	50
Total	381	177
Percent	68	32

Variations in measurements of particularly velocity and bulk density in units G2 and G3 are reflective of the lithologic heterogeneity of these units (see "Lithostratigraphy and Sedimentology" section). Both overall and detailed trends of gravimetric and GRAPE bulk density and velocity are, as a rule, well corre-

lated. In general, zones of lower densities and velocities are associated with intervals of greenish sand. These sands are, in contrast to the more fine-grained intervals, loose and more disturbed by drilling. Figure 19 represents a minimum strength profile for unit G2, because most of the red clay and silt are too hard for determination of shear strength. The soft sands suggest horizontal, as well as vertical, permeability barriers within the depositional basin. The former could result from one or more of several likely causes, including updip pinch-out of the sands, updip ore occlusion from clay matrix, or lap out against impermeable basement material. Vertical barriers are provided by the overconsolidated red clays and silts. Overconsolidation of these intervals may be the result of either, or both, chemical diagenesis or loading by ice or sediment.

Despite the apparently layered nature of at least part of the drilled sequence, no significant trend of velocity anisotropy exists. The average values of units G2 and G3 (Table 13) show that differences are less than 4% and that the A and C directions alternate in having the highest average velocity. This is somewhat surprising, as apparently a permeability anisotropy is present, causing the diagenetic color alterations observed in the sediment (see "Lithostratigraphy and Sedimentology" section). Possible explanations for the lack of velocity anisotropy include the use of too small sample sizes during velocity determinations or removal of anisotropy by diagenetic effects or by drilling disturbance.

The contact between units G2 and G3 and the reverse depth trends of the latter unit, most evident on GRAPE density and velocity profiles, are, at least in part, attributable to a change in lithology. Percent abundance of the major constituents, in particular clay minerals and feldspar, found upsection at Site 740 decreases at or near this boundary (see "Lithostratigraphy and Sedimentology" section). From core descriptions (see barrel sheets), a transition appears into generally more coarse-grained sandstones. Although the reverse trends can be explained by the aforementioned effects, increased drilling disturbance cannot be excluded. This will be enhanced in the less competent coarse-grained sandstones.

In summary, the physical properties in the upper 16.7 mbsf at Site 740 show values typical for relatively soft, post-glacial diatom oozes and diatom-bearing glaciomarine clays. Below this, an abrupt change in all properties marks the transition into a suite of much more competent sandstones, siltstones, and claystones across a boundary probably representing major phases of glacial erosion. A large scatter, particularly in density and velocity, superimposed on a general trend of increased compaction with depth reflects alternating lithology, with the less competent sands having lower velocity and bulk density.

SEISMIC STRATIGRAPHY

Site 740 is on the southeastern side of Prydz Bay, about 40 km from the Antarctic coast and 28 km landward of Site 741. The site lies above a sequence of reflectors that dips seaward beneath the continental shelf. The regional setting of these reflectors and other general information regarding the seismic stratigraphy of Prydz Bay are discussed in the "Seismic Stratigraphy" section of the "Site 739" chapter.

The dipping reflectors beneath Site 740 are part of Stagg's (1985) unit PS-4, which he also identified beneath the most landward parts of other seismic profiles across Prydz Bay. On ODP line 119-05 (Fig. 20), three seismic units are recognized. The uppermost seismic Unit I is thin, with flat-lying reflections. This unit corresponds to the thin layer of acoustically transparent ponded sediment seen on bathymetric profiles beneath Site 740 (Fig. 21). This unit is not well defined on the seismic profile and is confined to a 2- to 3-km-wide trough. Diatomaceous sediments were drilled from this unit.

Figure 19. Physical-properties profiles, Site 740. Small dots around the G1/G2 boundary mark continuous GRAPE measurements, blocked into 0.1-m averages, whereas the other values represent 2-min counts on discrete samples. Compressional-wave velocity was measured normal to bedding (filled circles) and parallel to bedding (open circles); small dots around the G1/G2 boundary are continuous *P*-wave-logger measurements, blocked into 0.1-m averages. The instrument used to measure shear strength is shown in Table 12.

The second seismic unit is also not well delineated, but is characterized by high-amplitude banded reflections that dip landward, opposite to the dip of underlying strata. These reflections, or banded group of interbed multiple-reflections, comprise the flanks of a depression that lies adjacent to the shoreline. The gentle and alternating dips of the high-amplitude reflections within the depression (e.g., 5 km southeast of Site 740) suggest that it is a channel, filled with sands and gravels that are highly reflective. In seismic data at Site 740, the depth to the bottom of the strong reflectors (gravels?) appears to be more than 100 m; however, cores from Hole 740A indicate the bottom is about 56 mbsf. No rocks, other than a few cobbles, were recovered in the interval 17–56 mbsf. At 56 mbsf fine-grained sands and silts were found. Consequently, the greater apparent thickness of high-amplitude reflections in seismic Unit II is probably due to the summation of interbed reflections and to the large impedance contrast (e.g., a broad reflector) between the overlying glacial gravels and underlying sands/silts.

Seismic Unit III is composed of the seaward-dipping reflections extending downward from the strong, low-frequency reflector forming acoustic basement. The amplitude of the dipping reflections is variable, but generally weak. Reflectors are semicontinuous and can be traced for distances up to 7–8 km before fading out and later reappearing. Sonobuoy and laboratory (Hamilton Frame) velocities show uniformly increasing velocities with depth for the upper part of this seismic unit. The uniform increase is characteristic of a compacting sand/shale sequence, such as that recovered in the cores below 60 mbsf.

The velocities of seismic Unit III are higher than those expected for rocks at a 60-m burial depth, indicating that the section was previously compacted and cemented. Prior compaction may have occurred during burial beneath grounded ice or sediment. The abrupt truncation of seismic Unit III indicates erosion of a previous sedimentary section.

Acoustic basement near Site 740 (Fig. 21) is a high-amplitude reflector, indicating a large velocity change; yet the flat geome-

Table 12. Physical-properties measurements, Hole 740A.

Core, section, interval (cm) ^a	Depth (mbsf)	Water content (%)	Porosity (%)	Wet-bulk density (g/cm ³)	Dry-bulk density (g/cm ³)	Grain density (g/cm ³)	GRAPE density (g/cm ³)	Undrained shear strength		Compressional-wave velocity (m/s)	
								(kPa)	Instrument ^b	A direction	C direction
1R-1, 60	0.60	74.90	87.43	1.22	0.31	2.31		0.8	F		1500.0
1R-1, 60	0.60							0.9	W		
2R-1, 22	6.22	45.23	69.53	1.64	0.90	2.79		3.2	W		1490.7
2R-1, 22	6.22							4.2	F		
3R-1, 120	14.90	72.80	87.97	1.32	0.36	2.72		0.2	W		
3R-2, 50	15.70	69.77	87.55	1.27	0.38	3.05					
3R-3, 106	17.76	18.10	37.80	2.22	1.82	2.79		24.0	F		1701.5
3R-3, 116	17.86							14.3	W		
3R-4, 4	18.24	17.58	35.90	2.17	1.79	2.67		53.0	F		1704.7
3R-4, 18	18.38							74.0	F		
8R-1, 40	57.00										2250.0
8R-1, 40	57.00	18.06	37.41	2.22	1.82	2.75		275.0	F		
8R-2, 70	58.80	22.10	44.68	2.36	1.84	2.89		145.0	F		1923.1
8R-3, 85	60.45	14.83	32.50	2.45	2.09	2.81	2.33	370.0	F	2102.9	2274.1
8R-4, 25	61.35	13.39	29.52	2.42	2.09	2.75	2.32			2275.5	1703.3
9R-1, 10	61.30	16.02	33.81	2.26	1.90	2.72					
10R-1, 46	66.66	18.14	38.76	2.25	1.84	2.90	2.17			2097.7	
10R-2, 90	68.60	15.48	33.50	2.31	1.95	2.79	2.28			2027.2	
11R-1, 73	71.63	16.34	35.12	2.29	1.92	2.81	2.19			1889.8	
11R-2, 65	73.05	17.94	38.07	2.28	1.87	2.86					2197.0
12R-1, 10	76.00	15.13	31.98	2.29	1.95	2.68	2.27			2348.3	
12R-1, 62	76.52							650.0	P		
12R-1, 78	76.68	18.23	37.63	2.26	1.85	2.75	2.35			2416.7	
13R-1, 26	80.86	17.60	36.38	2.15	1.77	2.72	2.33			2227.3	
13R-1, 90	81.50							7.7	P		
13R-1, 100	81.60	17.13	35.36	2.22	1.84	2.69					1954.5
14R-1, 37	85.97	15.43	33.60	1.77	1.49	2.82	2.39			2086.6	
14R-1, 117	86.77						2.33			2369.0	
14R-2, 81	87.91	13.91	31.59	2.46	2.12	2.90	2.42			2147.0	
14R-3, 30	88.90	12.18	27.13	2.40	2.11	2.73	2.34			2270.0	
15R-1, 48	90.68	11.97	27.23	2.47	2.18	2.80	2.37			2418.0	2515.0
15R-1, 125	91.45	11.55	26.09	2.46	2.17	2.75	2.41			2288.0	2651.0
17R-1, 63	100.53	17.31	35.24	2.22	1.84	2.64	2.24			2108.4	2098.0
17R-2, 9	101.49	14.44	31.29	2.32	1.99	2.74	2.35			2088.2	2227.3
19R-1, 16	119.36	12.41	28.20	2.48	2.17	2.82	2.37			2541.4	2680.4
19R-1, 107	120.27	11.27	26.55	2.56	2.27	2.89				2337.8	
20R-3, 87	128.67	10.89	25.14	2.59	2.31	2.79	2.29			2407.4	
20R-4, 18	129.48	11.39	25.27	2.54	2.25	2.67	2.37			2659.3	
21R-1, 5	126.05	9.87	22.91	2.70	2.43	2.76	2.48			2489.6	
22R-1, 48	129.28	11.59	25.76	2.45	2.17	2.69	2.39			2313.3	
22R-2, 60	130.90	14.36	31.42	2.34	2.00	2.77	2.28			2277.3	
22R-3, 52	132.32	12.61	27.52	2.34	2.04	2.67	2.29			2291.7	
23R-1, 29	138.79	14.30	31.09	2.32	1.99	2.75	2.23			2701.1	
23R-1, 72	139.22	14.83	39.30	2.35	2.00	2.78	2.24			2362.2	
24R-1, 71	148.81	8.50	19.99	2.65	2.42	2.73	2.39			2259.3	
24R-2, 1	149.61						2.31			2569.4	
25R-1, 3	157.83	10.85	24.42	2.39	2.13	2.70	2.44			2712.7	
25R-2, 17	159.47	12.06	27.28	2.43	2.13	2.78	2.41			2311.8	
26R-1, 26	167.76	8.86	20.82	2.51	2.29	2.75	2.47			2628.9	2527.3
26R-1, 123	168.73	10.35	24.27	2.53	2.27	2.82	2.42			2586.4	2721.6
26R-2, 85	169.85	11.16	24.96	2.45	2.18	2.69	2.34			2532.5	2533.3
26R-3, 38	170.88	9.70	21.56	2.41	2.18	2.60	2.33			2567.6	2650.0
26R-3, 117	171.67	11.26	25.87	2.50	2.22	2.80	2.39			2385.7	2467.9
26R-4, 64	172.64	11.15	24.40	2.44	2.17	2.61	2.35			2464.3	2480.8
27R-1, 13	177.23	11.54	25.42	2.45	2.17	2.65	2.37			2624.3	2542.1
27R-1, 129	178.39	10.21	22.52	2.45	2.20	2.60	2.35			2417.7	2176.5
27R-2, 31	178.91	13.97	29.96	2.30	1.98	2.68					
27R-3, 12	180.22	10.57	22.99	2.35	2.10	2.56				2558.8	2839.5
27R-3, 141	181.51	11.04	25.02	2.50	2.22	2.73	2.41			2720.5	2739.6
27R-4, 30	181.90	10.90	23.61	2.41	2.14	2.57				2641.2	2650.6
27R-5, 54	183.64	10.91	23.84	2.43	2.17	2.60	2.38			2516.1	
28R-1, 15	186.95	9.49	22.07	2.52	2.28	2.74	2.35			2691.0	2555.6
28R-1, 127	188.07	8.90	21.39	2.63	2.40	2.83	2.44			2445.9	2766.7
28R-2, 47	188.77	9.10	20.77	2.49	2.26	2.66	2.45			2732.0	3039.2
28R-3, 34	190.14	8.70	20.05	2.53	2.31	2.67	2.52			2560.0	
29R-1, 50	197.00	12.23	26.06	2.37	2.08	2.57	2.25			2321.4	2261.3
29R-2, 18	198.18	11.78	26.01	2.39	2.11	2.68	2.10			2291.3	2297.3
29R-2, 120	199.20	11.76	27.18	2.51	2.21	2.85	2.36			2420.7	2415.1
30R-1, 30	206.40	10.90	23.93	2.46	2.19	2.61	2.28			2385.7	2358.5
30R-1, 113	207.23	10.01	22.50	2.44	2.20	2.65	2.32			2430.1	2533.3
31R-1, 1	215.81	14.25	30.17	2.32	1.99	2.64	2.74			2064.4	2258.1
31R-2, 44	217.74	9.76	22.31	2.66	2.40	2.70	2.31			2625.5	2722.8
31R-3, 45	219.25	11.04	25.32	2.58	2.30	2.78	3.35			2302.3	2517.2
31R-4, 98	221.28	10.90	26.50	2.76	2.46	3.00	2.46			2555.6	2613.4
31R-5, 133	223.13	12.87	31.33	2.34	2.04	3.14	2.32			2341.7	2383.3

^a Top of measured interval (never exceeding 10 cm).^b F = fall cone apparatus; P = pocket penetrometer; W = Wykeham-Farrance motorized vane.

Table 13. Physical properties minimum, mean, and maximum values for the different geotechnical units in Hole 740A.

Geotechnical unit		Water content (%)	Porosity (%)	Wet-bulk density (g/cm ³)	GRAPE density (g/cm ³)	Compressional-wave velocity (m/s)		Undrained shear strength (kPa)
						A direction	C direction	
G1	min.	45.23	69.53	1.22	No data	No data	1491	0.2
	mean	65.68	83.12	1.36			1495	1.9
	max.	74.90	87.97	1.64			1500	4.2
G2	min.	8.50	19.99	2.22	2.17	1890	1702	7.7
	mean	13.12	28.83	2.40	2.35	2401	2392	179.0
	max.	17.31	35.24	2.70	2.52	2767	3039	650.0
G3	min.	9.76	22.31	2.32	2.10	2064	2258	Too stiff
	mean	11.55	26.13	2.48	2.35	2374	2436	
	max.	14.25	31.33	2.66	2.74	2626	2723	

try and absence of a high-velocity refractor suggest that the reflector may not be metamorphic basement. Stagg (1985) indicated that older sedimentary strata (PS-5) lie beneath the reflector and that metamorphic basement, which lies beneath the landward flank of the channel-like depression, is steeply down-faulted beneath the depression. We see no evidence for the fault or for layering beneath the acoustic basement. If the strong reflector is metamorphic basement, then it has been highly eroded before deposition of the overlying sedimentary section.

The uniform acoustic character of seismic Unit III and the continuous sequence of fluvial sandstones, siltstones, and mudstones from 60 to 225 mbsf suggest that the entire sequence to acoustic basement comprises a nonmarine sequence. The layers do not thicken substantially downdip, although at the bottom of the unit reflectors onlap (updip) the basement. The uniform layer thickness and reflector onlap suggest that seismic Unit III was deposited in a relatively stable environment on a slowly subsiding, eroded (i.e., unconformity) acoustic basement.

High-amplitude reflectors occur near the bottom of Hole 740A (Fig. 20). These and other large-amplitude reflectors in the seismic profiles across Prydz Bay may be caused by low-velocity, rather than high-velocity, rocks. A distinct change in lithology, which is accompanied by a decrease in velocity and density trends, occurs at 190 mbsf at Site 740 (see "Physical Properties" section). The velocity reversal may be the cause of the high-amplitude reflections. Here, as at Sites 739 and 742, where downhole logging was conducted, the lower velocities may be due, in part, to undercompaction or lack of cementation of sandy intervals. Unlithified sands with lower velocities than the surrounding fine-grained siltstones were observed at shallower depths in the Site 740 core.

If the reflector near the base of Site 740 is traced to the northwest and is projected through the seafloor multiple using dips of the overlying strata, then the reflector should lie about 2.0 km below the bottom of Hole 741A (Fig. 22). A subtle unconformity, or disconformity, appears on the seismic profile between Sites 740 and 741. Minor changes in stratal thickness and dip occur at the unconformity, and similar changes may occur beneath the seafloor multiple, as suggested by dips of shallower strata. The unconformity was identified by Stagg (1985) as separating units PS4 and PS2. This unconformity may, as Stagg suggests, separate pre-breakup (Permian to Lower Cretaceous) and post-breakup (Upper Cretaceous–Holocene) rocks. Below the unconformity the age of seismic Unit III at Site 740 is unknown, and above the unconformity the rocks at Site 741 are of Early Cretaceous(?) or Eocene(?) age.

Summary

Three seismic units are identified at Site 740. The upper two units are thin and difficult to resolve because of the interference

of high-amplitude reflections resulting from thin high-impedance layers. Seismic Unit III, which begins at 56 mbsf (from core data), comprises the seaward-dipping sequence beneath the Prydz Bay shelf. The weak and semicontinuous reflections seen throughout seismic Unit III suggest that the entire section is nonmarine, like the sands and silts found at Site 740. Strong reflections within seismic Unit III may be caused by low-velocity layers, possibly unlithified sands or sandy layers such as those postulated from downhole logging at Sites 739 and 742.

A subtle unconformity occurs between Sites 740 and 741, and the unconformity may be the boundary between the pre-rift, nonmarine rocks at Site 740 and the post-rift units at Site 741.

SUMMARY AND CONCLUSIONS

Site 740 is in the inner part of Prydz Bay on the East Antarctic continental shelf at 68°41'S, 76°43'E, in 810 m of water. Geophysical data suggests that Prydz Bay is underlain by a sedimentary basin, which is probably an offshore extension of the Lambert Graben. The main objectives of this site, therefore, were to sample and date the basinal sediments in order to provide information on (1) preglacial depositional environments and climate in East Antarctica and (2) the age of graben formation or the separation of East Antarctica from India. An additional objective was to study the younger sequence of glacial-preglacial sediments as a means of developing a generalized model for sedimentation on glacially-dominated shelves.

Rotary drilling began at this site on 25 January 1988 within an 800-m-deep seismically-defined erosional channel feature some 30 km off the Antarctic coastline. Soft diatom ooze, clayey silt, and silty clay were cored for 23.2 m prior to encountering boulders and cobbles of gneiss and acid/intermediate igneous rocks. These are underlain by a sequence of claystone, siltstone, and sandstone from 56.6 mbsf. Drilling of these resistant rock types resulted in slow penetration, poor core recovery, and excessive wear on the bit. The old bit was replaced using a free-fall reentry cone. Because of incomplete core recovery of the seafloor sediments, an additional mud-line core was taken (Hole 740B).

The sedimentary succession at this site is incomplete because of poor core recovery. It can be divided into three lithologic units. However, the contacts between these units are not seen, as they lie within zones of no recovery.

Lithologic Unit I (0–23.2 mbsf) is a soft diatom ooze alternating with clayey silt, with minor diatoms and gravel-bearing silty clay. The diatom ooze consists of 10%–20% terrigenous clastic grains, 15%–95% diatoms, and 0%–5% radiolarians and silicoflagellates of late Pliocene to Holocene age. The clayey silt contains 15% diatoms and 2% radiolarians; major nonbiogenic components are quartz and feldspar. The diatoms

Figure 20. Analog seismic-reflection profile (ODP line 119-05) across Site 740.

Figure 21. The 3.5-kHz bathymetric record showing acoustically transparent, ponded sediments filling a small depression beneath Site 740.

Figure 22. Analog seismic-reflection profile and interpretation across the inner part of Prydz Bay. The sonic reflector between Sites 740 and 741 may be an unconformity separating Permian-Mesozoic? (Site 740) and Lower Cretaceous? (Site 741) nonmarine rocks.

suggest a late Pliocene to Holocene age. The silty clay contains about 1% gravel and coarse sand in those parts recovered. The gravel component consists predominantly of granules and small pebbles of quartz, granite, and gneiss, with rare boulders of gneiss.

Lithologic Unit II (23.2–56.6 mbsf) consists of boulders and cobbles of gneiss and acid/intermediate rocks.

Lithologic Unit III (56.6–225.5 mbsf) consists of greenish gray and red sandstone interbedded with reddish and mottled siltstone and claystone. The ratio of sandstone to siltstone and claystone decreases toward the top of the unit, thereby imparting a fining-upward trend to the succession. No fossils have been found in this unit except for undatable algae spores.

Internally, the succession contains vertically stacked, pedogenically-influenced fining- and rare coarsening-upward sequences up to 5 m thick. Individual fining-upward sequences, where fully developed, are comprised of a scour surface overlain successively by conglomerate, coarse- to medium-grained, cross-bedded sandstone, and locally rippled and laminated fine sandstone, siltstone, and claystone. In some cases the conglomerate is missing, and the scour surface is directly overlain by sandstone. Rare coarsening-upward sequences typically consist of locally rippled fine sandstone or siltstone, sharply or gradationally underlain by claystone or silty claystone.

The sedimentary succession at this site records major changes in depositional environments. The older sediments of lithologic Unit III, about 300 m above the basement, are a "red bed"-type sequence deposited within a fluvially-influenced continental environment. Thus, the major sandstones are interpreted as channel sandbodies, with the finer siltstones and claystones representing low-energy flood basin and overbank deposits. The fining- and rare coarsening-upward trends recorded in the vertical changes in lithology, texture, and sedimentary structures are attributed to channel shifting (avulsion) and fluctuating discharge related to decelerating and accelerating flood flows. The presence of pedogenic carbonates (calcrete) provides useful evidence of the climatic conditions prevailing at this time. Comparison with Holocene and older equivalent pedogenic deposits shows that carbonate pedogenesis normally requires a warm, dry climate with a seasonal, but not excessively peaked, rainfall. The red and green color of the sediments is determined primarily by their residence time in the oxygenated vadose zone and the degree of diagenetic alteration of iron-bearing minerals such as mica. The overall upward-fining trend to the unit can be ex-

plained by a gradual decrease in paleoslope in response to weathering and the gradual reduction in source area relief or to decreased levels of tectonic activity. Poorly recovered Unit II probably represents a lodgement till; the overlying Unit I probably represents deposition of glaciomarine sediments, with the diatom ooze at the top representing present-day conditions of sedimentation.

The occurrence of well-preserved diatoms dates the interval from 0.0 to 13.7 mbsf to the Quaternary, with a probable age of less than 0.62 Ma. Below this interval, the diatoms are sparse, poorly preserved, and lack zonal taxa. Nevertheless, the species indicate a late Pliocene or younger age for the lower part of Unit I. Unit II is Pliocene to Holocene in age. Because of the lack of fossils in Unit III its age is uncertain, but it may correlate with parts of the Permian Amery Group sediments in the Lambert Graben, some 360 km to the southwest. Alternatively, they may form part of the Mesozoic?-Cenozoic "molasse and/or riftogenic" complex, which is thought to represent the preglacial sediments in Prydz Bay (Fedorov et al., 1982).

Paleomagnetic studies demonstrate that two polarities are present in the "red beds" of lithologic Unit III and that the primary magnetism is reversed rather than normal polarity. These results are incompatible with a Permian age for these sediments and require more detailed land-based paleomagnetic and paleontological (palynology) studies in order to more accurately characterize the age and magnetic history of the sediments. However, the existence of multiple components of magnetism in these rocks is an indication that they must have undergone a complex geological history, possibly related to the development of the Lambert Graben.

Alkalinity and sulfate analyses of interstitial waters from this site indicate high levels of microbial activity in the uppermost sediment layers, which emit a strong hydrogen sulfide smell. Oxidation of organics is responsible for the significant ammonium and phosphate enrichment of the interstitial waters in the uppermost sediments. High values for ammonium and phosphate suggest that most bacterial catabolism occurs near the sediment/water interface. A trace of hydrocarbon gas (8 ppm ethane and 1140 ppm methane) of probable biological origin was found in Sample 119-746A-14R-3, 0–1 cm, at this site. Both organic carbon and carbonate are low, except for the top of the hole down to 17 mbsf, where up to 1.4% TOC was recorded. However, these levels drop off downhole. The reason for this is not clear because of poor core recovery between 17 and 59 mbsf.

Below 59 mbsf the depletion in TOC probably reflects the dry, oxidizing nature of the original depositional environment.

Preliminary study of the marine ecosystem in Prydz Bay shows that few plankton species exist at this site that are not associated with ice. Furthermore, no single species is restricted to the ice, although those species adapted to growth in brine pockets in the ice seem to flourish best.

Abrupt changes in physical properties enable the division of the sediments into three geotechnical units. Geotechnical unit G1 lies within lithologic Unit I. Geotechnical unit G2 also lies within lithologic Unit I, extending down through Unit II into lithologic Unit III. The boundary between geotechnical units G2 and G3 occurs at about 190 mbsf and is indicated by a slight but significant change, which manifests itself in a slight reversal from normal to anomalous depth trends on index-property and velocity profiles. In addition, bulk density and velocity show decreasing trends below this boundary, while water content and porosity increase. This manifests itself in the occurrence of relatively thick, coarser, poorly consolidated, and more porous and permeable sandbodies.

REFERENCES

- Allen, J.R.L., 1965. A review of the origin and characteristics of Recent alluvial sediments. *Sedimentology*, 5:89-191.
- , 1974. Studies in fluvial sedimentation: implications of pedogenic carbonate units, lower Old Red Sandstones, Anglo Welsh outcrop. *J. Geol.*, 9:181-208.
- Fedorov, L. V., Grikurov, G. E., Kurinin, R. G., and Masolov, V. N., 1982. Crustal structure of the Lambert glacial area from geophysical data. *Antarct. Geosci., Symp. Antarct. Geol. Geophys.*, 3rd, 931-936.
- Fryxell, G. A., 1988. Polymorphism in relation to environmental conditions as exemplified by clonal structures of *Thalassiosira tumida* (Janisch) Hasle temperature on valve morphogenesis. In Mann, D., and Stoermer, E. (Eds.), *Proc. 9th International Diatom Symposium, Bristol*, 61-73.
- Fryxell, G. A., in press. Antarctic marine phytoplankton at the Weddell Sea ice edge: seasonal changes at the specific level. *Polar Biol.*
- Gieskes, J. M., 1983. The chemistry of interstitial waters of deep sea sediments: interpretation of deep sea drilling data. In Riley, J. P., and Chester, R. (Eds.), *Chemical Oceanography* (vol. 8): New York (Academic Press), 221-269.
- Goudie, A., 1973. *Duricrusts in Tropical and Subtropical Landscapes*: Oxford (Oxford Univ. Press).
- Hoban, M. A., Fryxell, G. A., and Buck, K. R. 1980. Biddulphoid diatoms: resting spores in Antarctic Eucampia and Odontella. *J. Phycol.*, 16:591-602.
- Ivanov, V. I., 1983. Sedimentary basins of Antarctica and their preliminary structural and morphological classification. In Oliver, R. L., James, P. R., and Jago, J. B. (Eds.), *Antarctic Earth Science*: Cambridge (Cambridge Univ. Press), 539-541.
- Kemp, E. M., 1972. Reworked palynomorphs from the west ice shelf area, East Antarctica, and their possible geological and paleoclimatological significance. *Mar. Geol.*, 13:145-157.
- Leeder, M. R., 1982. *Sedimentology: Processes and Products*: London (Allen and Unwin).
- McDuff, R. E., 1978. Conservative behavior of calcium and magnesium in interstitial waters of marine sediments: identification and interpretation [Ph.D. dissert.]. Univ. of California, San Diego.
- McDuff, R. E., and Gieskes, J. M., 1976. Calcium and magnesium profiles in DSDP interstitial waters: diffusion or reaction? *Earth Planet. Sci. Lett.*, 33: 1-10.
- Miall, A. D., 1977. A review of the braided river depositional environment. *Earth Sci. Rev.*, 13:1-62.
- Mizukoshi, I., Sunouchi, H., Saki, T., Sato, S., and Tanahashi, M., in press. Preliminary report of geological geophysical surveys off Amery Ice Shelf, East Antarctica. *Mem. Nat. Inst. Polar Res., Spec. Issue Jpn.*
- Mond, A., 1972. Permian sediments of the Beaver Lake area, Prince Charles Mountains. In Adie, R. J. (Ed.), *Antarctic Geology and Geophysics*: Oslo (Universitetsforlaget), 585-589.
- Stagg, H.M.J., 1985. The structure and origin of Prydz Bay and MacRobertson Shelf, East Antarctica. *Tectonophysics*, 114:315-340.
- Stumm, W., and Morgan, J. J., 1981. *Aquatic Chemistry*: New York (Wiley).
- Turner, B. R., 1978. Sedimentary patterns of uranium mineralization in the Beaufort Group of the southern Karoo (Gondwana) Basin, South Africa. In Miall, A. D. (Ed.), *Fluvial Sedimentology*: Mem. Can. Soc. Petr. Geol., 5:831-848.
- Turner, P., 1980. *Continental Red Beds*: Amsterdam (Elsevier).
- Walker, T. R., 1976. Diagenetic origin of continental red beds. In Falke, A. H. (Ed.), *The Continental Permian of Central, West and South Europe*: Dordrecht (Reidel), 240-282.

Ms 119A-108