Working Memory Against Neuropsychological And Behavioral Profiles in Children with Neurodevelopmental Disorders

Iyer Kamlam Gopalkrishnan¹, S. Venkatesan²

¹Research Scholar, DOS in Psychology, University of Mysore, Mysuru – 06

²Professor in Clinical Psychology, Department of Clinical Psychology,

Corresponding author: Iyer Kamlam Gopalkrishnan

Email – iyerkamlam@gmail.com

ABSTRACT

Background: Very few studies have attempted to investigate working memory (WM) in neurodevelopmental disorders (NDD). This study seeks to profile WM in children with NDD against their matched, typically developing children (TDC), all belonging to high socioeconomic status (SES) families. Secondary assessment of gender differences, if any, was also attempted. Profile analysis of certain domains in neuropsychological and behavioural functioning for the children is delineated.

Methodology: The study employed a cross-sectional 2-group comparison design by drawing a purposive sample of 84 children (42 in each group of NDD and TDC) in 6 to 8 years from across Bangalore city, India. We recruited the groups by matching them on intelligence, developmental skills, and SES. Assessment of WM was using verbal and visual n-back and spatial span tasks.

Results: The study depicted significant differences between the children of the groups in accuracy scores for the tasks. Errors from the n-back tasks showed significant differences, mostly in omission errors. We reported effect size and the probability of superior (PS) outcomes. We did not observe gender differences in the NDD group. Although, we have highlighted differences in profiles of neuropsychological and behavior functions between the groups.

Discussion: This dwells on the aspects that might have caused WM's observed deficits and explains the differences in these children's neuropsychological and behavior profiles.

Conclusion: WM differs significantly between children with NDD and TDC. However, we have not found differences in WM performance and gender in the NDD group.

Keywords: Cognitive, Adaptive skills, Indian, Children, Intellectual functioning.

(Paper received – 3rd March 2020; Peer review completed – 6th April 2021, Accepted – 8th April 2021)

INTRODUCTION

Executive Functions (EF) refers to the complex cognitive processes of control and coordination of thoughts and behaviours [1] and encompasses WM as one of its components [2]. Working memory (WM) remains an essential investigation topic in basic and clinical research [3]. This cognitive function enables the temporary storage and manipulation of information [4] and is required to execute many complex tasks [5]. It is one of the critical parameters of information processing in human development and cognition maturation [6]. We can reliably assess WM from as young as four years [7]. WM's neuropsychological assessment has shown a long development [8] and remains mostly dissociated from other EF [9]. Therefore, WM development continues into young adulthood, and impairments in WM can appear in isolation. Studies depict deficits in WM in children being associated with dyslexia [10], language disorders, mathematical disability [11], learning difficulties [12], attentional deficit disorders (ADHD) [13], specific language impairment [14], conduct disorders / oppositional defiant behavioural problems [15], autism spectrum

disorder (ASD) [16], academic achievement with teacher-student relationship [17], to name a few. Therefore, WM seems essential theoretically and for practical understanding of neurodevelopmental and educational disorders [11].

Neurodevelopmental Disorders (NDD) are difficult to conceptualize [18]. They are a group of conditions that occur at the onset of the developmental period, before the age of 6 years, marked by a range of deficits in various developmental domains such as personal, social, occupational, or academic functioning as provided in DSM 5 [19]. Impairment may occur in any of these development domains, starting in early childhood and assuming a steady course without remission and continuing through adult life [20]. The conditions are behavioural and cognitive disorders, whose etiology is complex to ascertain (ICD 11) [21]. Nearly 1/3rd of the population of India constitutes children under the age of 15 years [22]. Due to the risk factors [23] accompanying the lack of robust evidence on the burden of NDD in such a populous country, INCLEN Trust International conducted a study. It suggested that one child in eight in the age group of 2-9 years has at least one of the conditions of NDD [24]. This burden estimate appears alarming for a developing nation like India.

When studying children and their cognitive abilities, their *socioeconomic status (SES)* assumes importance, especially in a resource-poor country such as India. This construct is challenging to define singularly due to social and economic factors such as income, education, income-to-needs ratio, occupational, and social prestige. The brain is shaped continually by numerous physical and environmental factors that influence our cognitive, emotional, and behavioural abilities. The explanation of childhood SES is about the environment where they are raised. That is, their parent's SES, thereby inheriting the genetic characteristics of the same level of SES [25]. Observations depict a delay in children's overall development as the SES and gestational age decreases [26].

Abundant studies are reporting WM findings, specifically in NDD. For instance, the meta-analytic evaluation of 235 studies by Demetriou et al. [27] on WM's role in ASD suggested an overall impairment. A similar meta-analytic review in ASD across the lifespan depicted impairments in WM [28]. Kaspar, Alderson and Hudec [29] conducted a meta-analysis on ADHD and typically developing children (TDC). They found a large effect size in impairments in phonological and visuospatial tasks. In the learning difficulty domain, a meta-analytic review of 29 studies in reading, mathematical, and children with both difficulties, depicted impairments in verbal and numerical WM compared to TDC [12]. Gender differences in the presentation, symptomatology, and prevalence of NDD have been reported and discussed [30; 31]. However, few studies have implored the cognitive gender differences in early childhood with NDD as per our review. Gender differences might bring in a possible source of variation in WM abilities, and studies suggest they should be examined [32].

Studies on SES and WM in TDC children are many. In a meta-analysis of studies with children aged 2 to 18 years, a significant association with their executive functions was found [33]. Many other separate studies validate this finding on an association between a child's EF and family SES [34; 35]. A few western studies that examined WM in children with NDD have controlled SES variables. For instance, Rosenthal et al. [36] found EF impairments in 185 children across varying age groups with ASD when matched for SES.

Our Indian scenario is quite different. We conducted a brief review of EF/WM in children with NDD while considering the child's SES. We used search engines -- Google Scholar and PubMed. We used phrases -- "neuropsychological profile in Indian children," "cognitive functions in Indian children," "executive functions in ADHD," "executive functions in Autism Spectrum Disorders," "working memory in neurodevelopmental disorders in India," "executive functions in learning disorders," "socioeconomic status and cognitive abilities in Indian children with NDD," "socioeconomic status and cognitive abilities in Indian children with Autism," "socioeconomic status and cognitive abilities in Indian children with ADHD," "socioeconomic status and cognitive abilities in Indian children with ID," "socioeconomic status and cognitive abilities in Indian children with LD." We found a few articles on WM in children with NDD while incorporating SES (Table 1).

Table 1 shows one study on WM in NDD by considering SES. There are no studies on gender differences. Our review offered very few studies on functional domain differences in the developmental and intellectual functioning of NDD. For instance, Malhi and Singhi [52] charted children's developmental profile with ASD

and developmental delay on four-year-old's. Another study on children with Down's syndrome examined profiles of the Binet-Kamat Intelligence Scale (BKIS) of intelligence, with and without mild ID [53].

Table-1 Studies on EF in Indian children with NDD incorporating SES and gender differences

NDD	Sample Size	Age group (in years)	SES assessed Y/N	EF assessed	Gender differences assessed Y/N	Significant results in EF comparison to other / control group Y/N	Citations
LD	35	7 to 14	N	EF	N	Y	[37]
SLD	59	4 to 13	N	Attention, Language	N	N	[38]
LD	46	7 to 14	N	Attention, Immediate recall	N	Y	[39]
ADHD	30	6 to 14	N	WM	N	Y	[40]
LD	18	8 to 12	N	Processing Skills	N	N	[41]
LD & ADHD	32	11 to 15	N	EF	N	Y	[42]
ADHD	60	6 to 13	N	Verbal WM	N	N	[43]
LD	4	11 to 14	N	EF	N	Y	[44]
LD	51	10 to 12	N	The verbal short-term memory	N	Y	[45]
ADHD	20	6 to 14	N	EF	N	Y	[46]
SLD	104	8 to 10	Y	EF and Processing skills	N	Y	[47]
LD	30	8 to 15	N	WM	N	Y	[48]
LD	5	8 to 11	N	Processing skills	N	Y	[49]
ID	7	8 to 15	N	WM	N	Y	[50]
ADHD	30	8 to 12	N	WM	N	Y	[51]

[SES = Socioeconomic Status; ADHD = Attention-Deficit Hyperactivity Disorder; LD=Learning Disorder; SLD=Specific Learning Disorder; ID=Intellectual Disability; Y=Yes; N=No]

Similar profiling of developmental and intellectual functioning in children with NDD might provide essential information on its functional domains. Notably, more research and subsequent interventions can help reduce the NDD burden [54]. Therefore, there is a felt need to examine WM, its gender differences, and the profiling of developmental and intellectual functioning in NDD and TDC, with a backdrop of SES.

The Present Study:

This study attempts to capture WM performance in NDD and TDC. Using a unitary definition of NDD that denotes delays/impairment in a child's identified developmental domains is justified for many reasons.

- The classification of heterogeneous conditions of NDD based on diagnostic manuals such as the Diagnostic and Statistical Manual of Mental Disorders (5th ed.; DSM-5) [19] and ICD-11 [21] take into account particular signs or groups of symptoms observed in patients in a standardized manner [55]. Diagnostic distinctions are essential [18]. However, such differences can change depending on the given context. They should best reflect the purpose at hand [55].
- NDD could be looked at as a dimensional trait, thereby offering more flexibility, both for research and clinical practice [55].
- Problems of co-morbidity and co-occurrence of many symptom patterns in NDD may exist due to shared genetic risks and biological characteristics. This problem might make it challenging to seek single patterns of symptoms of a disorder [56]. The same conditions can give mixed patterns of symptoms in different children who certainly go against clarity and understanding, especially in research [20].
- Cognitive deficits such as EF have considerable overlap across various conditions of NDD [57].

Therefore, this study has taken NDD to mean neurodevelopmental impairment as an essential aspect of liability rather than categorical diagnostic grouping. With this framework of NDD and in the background of a high SES, it could be that WM might play an essential role in cognitive development. As per Baddeley's multi-component model, we assessed WM and its sub-systems of verbal, visual, and visuospatial components in this study [5]. The sample is in the age group of 6-8 years, as findings on improvement in neuropsychological functions from early childhood onwards in Indian children is noted [58]. We assessed the background measures of development (for the classification of groups NDD and TDC) and intellectual functioning.

The research questions that emerge from the preceding are:

- Are there deficits in WM performance in children with NDD and the TDC?
- If deficits are observed between NDD and TDC, what are the similarities and differences between them?
- Are there gender differences in WM performance within the NDD group?
- What might be the profiles of neuropsychological and behavioral functioning in children with NDD and the TDC?

Operational Definitions

The key terms used in this study: NDD, WM, SES, and neuropsychological functioning are operationally defined as under:

- NDD is any impairment in the developing brain or the CNS originating during the developmental period. It is characterized by a delay of three or more months or disturbance in acquiring skills in at least two domains: motor, sensory, speech, language, social, cognition, play, and academics. It is measured on valid tools.
- WM is the ability to store and manipulate information for brief periods. It covers verbal and visuospatial facets as measured by N-back and spatial span tasks.
- SES is the child's parental level of education, occupation, annual family income, and total net worth as assessed on a standardized contemporary scale.
- Neuropsychological functioning refers to mental aspects such as general cognitive ability, verbal, non-verbal, numerical reasoning, language ability, memory skills, and conceptual thinking. It is measured on a valid normative age scale.
- Behavioural functioning is the specific asset or deficit observable, or measurable actions assessed on a standardized behavior assessment tool for the targeted population.

METHODOLOGY

We employed a two-group cross-sectional comparative study design and recruited one group of children with NDD and the other TDC group.

Participants: We used a non-probability purposive technique to recruit the participants. A total of 84 children with each group having 42 children were selected. The age group was 6 to 8 years, with both boys and girls. There were 32 boys (76.2 %) and 10 girls (23.8 %) in the NDD group, with 23 boys (64.3 %) and 15 girls (35.7 %) among the TDC. The following inclusion criteria were used (Table 2).

Table-2 Inclusion criteria for the groups

Inclusion Criteria for NDD	Inclusion Criteria for TD
In the age – a group of 6 to 8 years	In the age – a group of 6 to 8 years
Children from high SES families	Children from high SES families
Found to be having a deficit in at least two domains and delay of three months or more in development	Found to be having no deficit/delay in development
IQ above or equal to 70	IQ above or equal to 70
Children not on medication for NDD or other disorders	Children not on medication for any disorders
No other medical conditions associated with NDD	No history of physical/neurological/medical conditions

[NDD=Neurodevelopmental Disorders, TD=Typically Developing, SES=Socioeconomic Status]

Measures:

The measures used in the study are divided into two sub-headings, as given below:

Background (Control) Measures: Parent responded to the child and family demographic information. We used the *NIMH-SES* scale to assess the child's SES [59]. This tool gives an index based on the family's highest levels of education, pooled income, type of occupation, and net assets. We included a family SES score of 16 and above. Along with DSM-5 [19], We used *Activity Checklist for Preschool Children with Developmental Disabilities* (ACPC-DD) [60], a standardized behavior assessment device to assess each child's developmental delay. This checklist consists of 400 items distributed evenly across eight behavioural domains relevant to children's daily activities affected by NDDs and between 0 to 72 months. There are 50 items each under its eight demarcated domains like sensory, gross motor, fine motor, communication, play, pre-academics, self-help, and cognitive. The examiner gives each item a score on a five-point scale, depending on the assistance given to perform that item. The maximum score possible for a child on this tool is 2000. We included children with delays in domains of three or more months and at least two domains in the NDD group while children with nil developmental delay in the TDC. We chose this test to view the developmental delay observed in this age group of children with NDD. We used the same test to assess TDC to compare the NDD and TDC groups. *Binet Kamat Intelligence Scale* (BKIS) [61] is a modified version of the Stanford Binet Scale measuring the intelligence of Indian children [62]. This age-scale is for 3 to 22 years. Non-verbal,

verbal, power, and speed tests are included in BKIS. The 'basal age' is the highest age level below which all test items are passed, and 'ceiling age' is where all items are failed. The examiner calculates mental age by adding appropriate credits to every test's basal age beyond the basal level. The Intelligence Quotient (IQ) is computed as a percentage product of MA over CA. A profile analysis of BKIS test performance helps to understand the pattern for each subject. Profiles depict various neuropsychological functional domains like language, conceptual thinking, numerical and non-numerical reasoning, visuomotor ability, non-verbal reasoning, and social intelligence [63, 64]. We included all children above an IQ score of 70.

WM measures: WM measures were administered on both the groups in random order as provided from the table of random sets generated using computer software [65]. N-back and Spatial span tasks were incorporated to measure verbal and visuospatial components of WM. The N-back task is commonly used as an assessment in cognitive neuroscience to measure a part of working memory and working memory capacity. The N-back task captures the active part of working memory [66]. The spatial span task has many variations. It is a non-verbal counterpart to the digit-span task. Its roots are traced to the Knox cube imitation test [67]. When used with many blocks, this task is called the Corsi Block tapping test [68]. Some intelligence tests, such as the revised Stanford Binet (SB-5), call it the Block-span task [69].

Verbal WM: The 1-back and 2-back verbal [70] tasks consist of 30 consonants common to Indian languages. They would be presented locally at the rate of one consonant per second. Out of 30, 9 of them are repeated in the list randomly interspersed. In the 1-back task, the participant taps his/her dominant hand on the table if the examiner repeats the consonant consecutively. In the 2-back task, the participant responds by tapping the table if the examiner repeats the consonant after an intervening consonant.

Visuospatial WM: We assessed visuospatial WM using Visual n-back tasks and Spatial Span tasks. These tasks are from the NIMHANS Child Neuropsychological Battery [71]. In 1 and 2 back tasks, there are 36 cards of the same dimensions, with a black dot placed randomly in the card, again of the same dimension throughout. In the 1-back task, the participant responds by tapping the table if the dot repeats itself in the same location, consecutively. In the 2-back task, the participant responds if he/she sees the dot at the same location after one intervening random card. The number of correct responses (accuracy) and errors in both tests (verbal and visual) form a score. In the Spatial Span task, four cubes are arranged in a row with 1 inch in between, with the examiner tapping using the fifth cube for different sequences. The child is required to repeat the sequence as tapped by the examiner. The sequences are both forward and reverse, and the accuracy scores are the number of correct sequences tapped for both the conditions.

Procedure: All children were recruited with the parents' written consent and complied with the ethical rules [72]. The period of collection of the data was from May 2019 to November 2019. We recruited children for the NDD group from special education therapy centres, child development clinics, and multispecialty clinics of Bangalore city, Karnataka, India. Figure 1 depicts the flow diagram of the final sample included in the NDD group. For the recruitment of TDC, we contacted the families using the snowball technique of sampling. We visited housing communities and schools in Bangalore city, Karnataka, India. The authors used a computer coded and amenable data intake and record sheet for every child to facilitate ease of scoring and administration of the measures. The children were all assessed in well-lit rooms of either the clinics or their homes in one to two sessions of 50 minutes each by the author, who has a Rehabilitation Council of India (RCI) approved pre-doctoral qualification in clinical psychology.

STATISTICAL ANALYSIS

The results were analysed using the Statistical Package for Social Sciences (SPSS version 23.0) [74]. We screened the data for normality using Shapiro Wilk's test, and depending on the obtained results, parametric (normal distribution) and non-parametric (skewed distribution) tests were conducted to infer appropriately.

RESULTS

We present the results of the study under the following four distinct but interrelated headings about WM in TDC and NDD groups: (a) Sample characteristics; (b) Distribution of WM scores; (c) Gender differences in WM scores in NDD group; and (d) Domain analysis of neuropsychological and behavioural functions.

Figure 1: Flow Diagram of the sample included in the NDD group Adapted [73] **Sample Characteristics**

A perusal of demographic characteristics of the sample (Table 3) depicts the demographic and clinical features of the children from the NDD and TD groups. The children are similar in their chronological age range ($p > .05$). However, they appear to vary by education, IQ, and developmental scores ($p < .001$). The children of the NDD group (N:42), to a more considerable extent, were in upper kindergarten (35.7%) and first grade (31%), while the children of the TDC group (N:42) were in their Grade II (50%) in their current level of education. Many of the NDD group children were studying a grade less than their peers in the TDC group. The groups' probability results for the IQ and developmental scores depict significant differences ($p < .001$). NDD was, on average, in the below-average category of intellectual functioning with a significant delay in development.

Table 3: Demographic characteristics, IQ and developmental scores for the groups (N = 84)

Demographic Characteristics	NDD (n=42)			TDC (n=42)			p*
	f%	Mean	SD	f%	Mean	SD	
Age		6.72	.83		6.70	.51	.94
Education Level							
Lower Kindergarten	7.1			0			
Upper Kindergarten	35.7			4.8			
First Grade	31			35.7			
Second Grade	16.7			50			
Third Grade	9.5			9.5			
Intellectual Functioning		82.50	9.98		113.07	14.99	.000**
Developmental Scores		1314.05	273.21		1967.55	25.89	.000**
SES Scores		19.88	.33		19.98	.15	.09

[*Probability obtained using t-test statistic; **Significant at $p < .001$ level]

Distribution of WM scores

Each N-back task has the accuracy scores (number of correct responses), omission, and commission errors. Omission errors take place when the child fails to give an accurate response to a correct stimulus. Commission errors occur when a child gives an incorrect response to a wrong stimulus. The *distribution for WM scores* presented across span test forward and backward, 1-back as well as 2-back task conditions for both these types of errors shows consistently at least twice or even thrice as much better performance by children with NDD to TDC (Table 4). Thus, we answer the first research question about evident deficits in WM performance of children with NDD vis-a-vis age and SES matched TDC ($p < .001$).

The next level analysis of *accuracy scores* depicts higher median scores for verbal 2 tasks with the lowest scores on the TDC's visual 1 task. Inter-quartile Range (IQR) on all three tasks, viz., verbal, visual, and spatial span, indicates substantial TDC variations. The children with NDD portray a different picture. The IQR's for all the tasks are smaller than the TDC. The highest median is noticed for the verbal 1 and span forward task and the lowest for the visual 2 tasks. In short, the results indicate that the TDC outperformed the NDD group in all the WM tasks. The *analysis of errors*, both omission, and commission, is deemed necessary when using N-back tasks [75]. Once again, we see that the two groups differ significantly for all N-back tasks ($p < .01$). Omission errors, more than commission errors, are observed in the NDD group almost by three to four times as compared to TDC. The mean ranks for commission errors are almost identical for both groups. The analysis of *effect size* usually restricted to studies on meta-analysis and intervention outcomes [76], was attempted in this study as a descriptive value. It was calculated by hand using the formula given by Siegel and Castellan in 1988 [77]. We observed a noticeable impact on WM in the NDD group for the verbal 1 task, its omission errors, and the span task. A moderate effect was observed in visual N-back tasks while a little impact was observed in the verbal 2 task, visual 1 task, and their errors, respectively, from the NDD group. Sometimes, studies suggest estimating the *probability of superior (PS)* outcome for intervention-based

studies [78] where they compare pre and post data. They are attempted in this study purely for its descriptive value, and they doubly confirm the same results as from the effect size.

Table 4: Distribution of WM scores for NDD and TDC (N=84)

WM Scores	NDD (n=42)			TDC (n=42)			p*	η^2	PS
	Mean Ranks	IQR	Median	Mean Ranks	IQR	Median			
Verbal 1-back Accuracy Score	22.64	2.00	2.00	62.36	2.00	8.00	.000***	.68	.69
Verbal 1-back Omission Errors	62.61	2.00	7.00	22.39	2.00	1.00	.000***	.68	.69
Verbal 1-back Commission Errors	46.46	2.00	1.00	38.54	2.00	1.00	.185	-	-
Verbal 2-back Accuracy Score	22.95	2.00	1.00	62.05	5.00	10.00	.000***	.65	.68
Verbal 2-back Omission Errors	62.61	2.00	17.00	22.39	2.00	1.00	.000***	.65	.68
Verbal 2-back Commission Errors	50.71	3.00	1.00	34.29	4.25	5.00	.001**	.12	.53
Visual 1-back Accuracy Score	29.32	1.25	1.00	55.68	4.25	4.00	.000***	.30	.58
Visual 1-back Omission Errors	55.86	1.25	8.00	29.14	3.00	2.00	.000***	.30	.58
Visual 1-back Commission Errors	42.75	3.25	2.00	42.25	2.00	2.00	.924	-	-
Visual 2-back Accuracy Score	28.08	1.00	0.00	56.92	3.00	7.00	.000***	.35	.60
Visual 2-back Omission Errors	57.35	1.00	9.00	27.65	2.00	2.00	.000***	.38	.61
Visual 2-back Commission Errors	38.21	3.00	1.00	46.79	1.25	4.00	.101	-	-
Span Test Forward Accuracy	25.86	1.00	2.00	59.14	2.00	2.00	.000***	.48	.63
Span Test Backward Accuracy	28.06	.25	.00	56.94	3.00	5.00	.000***	.40	.61
Span Test Total Accuracy	25.18	2.00	2.00	59.82	5.00	8.00	.000***	.52	.64

[p* = probability obtained using non-parametric test (Mann-Whitney U Test statistic) and η^2 = Eta square depicting effect size obtained for non-parametric test; ** Significant at $p < .01$ level; ***Significant at $p < .001$ level; WM = Working Memory; IQR = Interquartile range; PS = Probability of Superior Outcome]

Gender Differences in WM scores in NDD group

The analysis of gender differences in WM for children with NDD did not yield significant differences ($p > .05$; Table 5). The mean ranks on girls' accuracy scores are better than boys, indicating their slightly better performance on all WM tasks. The variability of their accuracy scores is greater than boys. We observed the mean ranks of omission and commission errors to be fluctuating in both groups. We also note that the sub-sample size of boys and girls are unequal. Thus, our answer to the second research question is that there are no gender differences in WM performance of children with NDD, matched for SES ($p > .05$).

Table 5: Gender distribution of WM scores for NDD group (n=42)

WM Scores	Boys (n=32)			Girls (n=10)			p*
	Mean Ranks	IQR	Median	Mean Ranks	IQR	Median	
Verbal 1-back Accuracy Score	20.31	2.00	2.00	25.30	1.75	2.00	.25
Verbal 1-back Omission Errors	22.59	2.00	7.00	18.00	1.75	7.00	.29
Verbal 1-back Commission Errors	20.22	2.00	1.00	25.60	2.50	2.00	.21
Verbal 2-back Accuracy Score	20.08	2.00	1.00	26.05	1.50	2.00	.16
Verbal 2-back Omission Errors	23.02	2.00	17.00	16.65	1.50	16.00	.13
Verbal 2-back Commission Errors	20.14	2.00	1.00	25.85	4.25	3.00	.18
Visual 1-back Accuracy Score	19.73	1.00	1.00	27.15	3.50	2.00	.08
Visual 1-back Omission Errors	23.27	1.00	8.00	15.85	3.50	7.00	.08
Visual 1-back Commission Errors	19.28	3.00	2.00	28.60	3.50	3.50	.30
Visual 2-back Accuracy Score	20.81	1.00	0.00	23.70	2.00	0.00	.43
Visual 2-back Omission Errors	22.19	1.00	9.00	19.30	2.00	9.00	.43
Visual 2-back Commission Errors	19.83	3.00	1.00	26.85	9.25	2.00	.10
Span Test Forward Accuracy	20.83	1.00	2.00	23.65	2.25	1.50	.34
Span Test Backward Accuracy	20.55	0.75	0.00	24.55	0.25	0.00	.86
Span Test Total Accuracy	21.64	1.00	2.00	21.05	3.25	1.50	.51

[*p = probability obtained using non-parametric test (Mann-Whitney U test statistic) and η^2 = obtained for non-parametric test; WM = Working Memory; IQR=Interquartile range]

Domain analysis of neuropsychological and behavior functions

The next research question was about uncovering the profiles of neuropsychological and behavior functions for children with NDD and the TDC. On the whole, the overall, as well as individual domain-wise performance of TDC, is superior to children with NDD across all the neuropsychological functional domains as assessed on by profile analysis on BKIS ($p < .01$; Table 6; Figure 2) as well as the behavioral domains of ACPC-DD ($p < .01$; Table 7; Figure 3).

Table 6: Comparative mean and SD mental age scores of samples on functional domains of BKIS (n=42)

Functional Domains	TDC		NDD		p*
	Mean	SD	Mean	SD	
Language	7.29	1.75	5.74	1.25	.000***
Meaningful Memory	7.25	1.86	4.36	1.85	.000***
Non-Meaningful Memory	7.45	1.58	4.29	1.50	.000***
Conceptual Thinking	7.26	1.99	5.46	1.83	.000***
Non-verbal Reasoning	5.30	3.09	3.19	2.80	.002**
Numerical Reasoning	7.61	1.21	5.83	1.84	.000***
Visual-Motor Ability	7.02	0.81	4.69	2.52	.000***
Social Intelligence	7.57	1.08	5.19	1.18	.000***
Overall	7.09	1.67	4.84	1.85	.000***

[* probability using Mann-Whitney test; ** $p < .01$; *** $p < .001$]

Table 7: Comparative mean and SD scores of samples on behavioral domains of ACPC-DD (n=42)

Behavioral Domains	TDC		NDD		p*
	Mean	SD	Mean	SD	
Sensory	217.41	17.90	248.52	1.89	.000**
Gross Motor	193.41	31.69	246.57	3.14	.000**
Fine Motor	163.67	40.91	241.57	6.62	.000**
Communication	173.07	40.96	245.55	4.53	.000**
Play	167.17	41.87	247.29	3.20	.000**
Cognitive	113.62	47.44	247.29	6.53	.000**
Pre-academics	121.64	61.73	246.52	3.58	.000**
Self-Help	157.05	50.34	246.57	3.23	.000**
Overall	163.38	46.61	246.24	4.09	.000**

[* probability using Mann-Whitney test; ** $p < .001$]

Figure 2: Distribution of domains for each of the groups

[NDD=Neurodevelopmental Disorders; TD=Typically Developing; Developmental Domains are Sensory, Gross Motor, Fine Motor, Communication, Play, Cognitive, Pre-academic, and Self-help skills]

Figure 3: Distribution of intellectual functioning domains between the groups

[NDD=Neurodevelopmental Disorders; TD=Typically Developing; Lang = Language; M.M = Meaningful Memory; Non-M.M = Non-Meaningful Memory; Con.Thn = Conceptual Thinking; Non-V.R = Non-Verbal Reasoning; V.R = Verbal Reasoning; N.R = Numerical Reasoning; V.M.A = Visuo-Motor Ability; S.I = Social Intelligence]

On the BKIS, for instance, the profile shows consistently high scores across language, non-verbal reasoning, meaningful and non-meaningful memory, conceptual thinking, visual-motor ability, and social intelligence. We have omitted verbal reasoning since it does not developmentally fall within the age range of children recruited for this study. Likewise, the NDD group of children appears to be affected more in the domains of cognition and pre-academic, less in communication, play, self-help skills, and least affected in the sensory and gross motor domains, as evidenced by ACPC-DD. The mean scores of cognitive and pre-academic domains are twice as high for the TDC than the NDD group. A comparative profile analysis of performance on BKIS (Figure 3: Table 7) as a measure of neuropsychological functions consistently indicates superior performance, higher mental age, and IQs by TDC against children with NDD across all areas including language, meaningful and non-meaningful memory, conceptual thinking, non-verbal reasoning, numerical reasoning, visuomotor ability, and social intelligence respectively. These findings confirm two distinctive levels of profiles in children's neuropsychological functions with NDD and the TDC.

DISCUSSION

The first research question was to see for significant differences in WM between the two groups of children. The present data provided evidence for the differences in WM between children with NDD against their age, and SES matched peers in TDC. A profile analysis of neuropsychological and behavioural functioning shows a consistently high and better TDC performance compared to children with NDD. In a related study on 9-10-year-olds from three groups: one with above-average levels of IQ with Autism, ADHD, and TDC were compared for their EF using the Behavior Rating Inventory of Executive Function (BRIEF). There was clear evidence on the deficits in verbal WM in the children with ASD and ADHD compared to the TDC. However, this study did not find differences in the visuospatial WM [79, 80]. Further, they found no significant impairments in their clinical groups.

The patterns of impairment overlapped that it lent support to the unitary approach towards NDD, as postulated in our present study. Henry and Botting [81] found WM deficits in children with developmental language impairment. Similar support for our study is put by Englund, Decker, Allen, and Roberts [82]. They examined a mixed clinical group of ADHD, ASD, and TDC in 4-18 years for specific cognitive abilities. Their results depicted that the TDC separated both clinical groups on WM deficits. They found verbal WM and visuomotor integration scores to be the best discriminating measure between these children. Specific domain deficits such as language, learning, self-help skills, and motor impairments provide us with several study outcomes to discuss WM results obtained in our study. For example, many studies have reported WM impairment in children with learning difficulties [9, 83, 84]. Studies have found deficits in visuospatial WM in children with arithmetic and learning disabilities [85]. The cognitive and pre-academic domains of the NDD group (Figure 3) are similarly impaired in our study, thereby pointing to learning problems in these children.

Findings report IQ as a dissociable cognitive skill from WM, especially in learning outcomes [86], such as academic attainment [87]. Interestingly, self-help skills such as dressing, eating, and bathing could be impaired due to WM deficits hampering their execution [88]. Spatial WM deficits are observed in ASD [89, 90], while verbal deficits have been observed in SLI [91] and learning difficulties at school [9]. Many studies have surfaced in the last decade on children with varied developmental impairments due to lower gestational ages' etiology. Sparse learning, occupational outcomes, higher motor impairments, and reduced cognitive functioning have been found in children born very preterm [92]. They were susceptible to attentional and poor verbal WM [93]. A missing link in the present study to explain the WM-IQ deficit in the NDD group could be the children's processing speed. Speed and intelligence at different ages of childhood are correlated [94]. Slow processing speed impacts a wide range of cognitive functions, such as EF and attention [87]. Children with WM deficits would also be slower to respond to several tasks [95].

Hence this could be one explanation for the current findings and needs to be explored. The type of assessment used to measure WM also needs evaluation. N-back assessments, which have repeatedly been used in various formats, rely on motor speed and information processing speed, rather than only WM [96]. This study has observed that the number of commission and omission errors is more for the NDD group

than the TDC, especially in the 2-back tasks. An increase in memory load as each difficulty level increases could be a reason. This increase demands more mental strategies (for 1-back = maintenance, 2-back = maintenance + monitoring, and so on) from the participant to complete the task at hand [97]. Although we found no significant differences in commission errors for verbal 1 and visual 1 and 2 tasks, indicating both groups being similar in their response to inhibit unnecessary items.

This study's second research question was to check any *gender differences* in children with NDD on WM measures. This question is supported negatively. A reason for such a finding could be proper gender matching not done in this study. The boys are three times more than the girls in this sample. Such unequal and small sub-sample sizes could probably be the reason [98]. Brain activation studies report gender differences in WM with varying typical populations. Studies show mixed results in adults and adolescents [98, 99, 100]. A few studies on TDC depict lower accuracy scores and more commission errors in boys' N-back task than girls from the age group of 6-8 years [101]. By contrast, Leon, Cimadevilla and Tascon [102] found boys outperforming girls in the age group of 6 to 8 years on computerized and classic psychometric assessments of spatial WM. A few studies found no gender differences in verbal and visuospatial WM tasks between 4 to 11 years [7]. Gender differences in the presentation and prevalence of a few NDD are reported [31, 24], but cognitive functions are far and few. With few studies on clinical samples of children evaluating gender differences, the available findings are inconsistent. For example, in a study on ADHD children and TDC by Munoz-Suazo, Navarro-Munoz, Diaz-Roman, Porcel-Galvez and Gil-Garcia [103], many cognitive functions separated the two groups of children. Nevertheless, they found no significant difference between boys and girls in WM and processing speed. In contrast, Nyden, Hjelmquist, and Gillbery [104] found that in 17 boys and girls with ASD, boys did better on measures of EF, especially visuospatial skills, than girls. Another study depicts similarities in boys' and girls' verbal profile, while no significant differences emerged in the non-verbal domain of neuropsychological assessment [105]. One reason for different conclusions in EF could be that the strategies employed by both genders are supported by different neurobiological mechanisms, though the ability is the same [106].

The third research question was to *delineate the differences in neuropsychological and behavioral functioning domains* in children's two groups. Many studies attempting to profile the functional skills in NDD [107, 108] have elicited the same patterns as derived in this study (Figure 3). For example, Mouga, Almeida, Cafe, Duque and Oliveira [109] profiled the adaptive skills of ASD versus other NDD group children between 6 to 18 years to find lower scores on all the domains assessed-communication, socialization, daily living skills and a composite measure of all the adaptive behaviors. Similar lowered ratings were reported in children between 9 to 10 years with ASD and ASD + ADHD as compared to their matched TDC [88]. Another prospective study on a cohort of 3-year-old children diagnosed with NDD who were later assessed again at age 7 on the same scales reported lower scores on all the domains such as personal-social, adaptive, fine, and gross motor skills, motor, and communication skills [110]. Researchers suggest interpreting these results with caution as a slow rate of progress vis-a-vis TDC rather than a stagnant development stage in children with NDD [107]. Profile analysis of neuropsychological and behavioral functions provides a better understanding of which group differs in that area rather than looking at a gross IQ score alone.

Mouga et al. [111] found results along similar lines. They compared the ASD group and other children with NDDs on an intellectual profile and discussed their influences. They found distinct profiles for all the clinical groups. Such profiles would assist a clinician in remediating the child and psychoeducation of the family [110]. The finding in this study on lowered IQ scores in the NDD group compared to the TDC is corroborated by several others [112-113]. Frith and Happe [114] argued that low IQ could be a developmental consequence of brain damage.

The present study has selected children above an IQ of 70 for two reasons. Firstly, to have reasonably homogenous groups using the cut-off of 70 instead of directly trying to match/co-vary IQ or other IQ covariates. There are reports that otherwise, a distortion of difference may appear between the experimental and control groups [115]. Second, we wanted to avoid the possibility of "double vulnerability" [116] of NDD with comorbid or co-occurring disorders of intellectual development [117].

This study has also intentionally attempted to control the socio-demographic characteristics associated with IQ in this sample, such as parental occupation, parental education, and income. These variables are highly correlated with IQ [25] and may indirectly account for differences in the outcome variable. Considering the

association between WM and IQ in NDD, one study with a clinical sample of ADHD suggests that group differences found in the IQ scores could be due to differences in cognitive abilities like WM, rather than just intellectual functioning [82]. Another study on 5-year-olds and 8-year-olds with epilepsy observed a significant correlation between WM and IQ [118]. WM for children with NDD has a researcher thinking about different diagnostic categories, segregated development domains with varying underlying mechanisms. All this makes the formulation of one conclusion challenging if not difficult. We could describe our study's outcomes at many levels, like the domain impairments, IQ, and functions, other cognitive abilities such as processing speed and choosing a valid neuropsychological task for measurement.

CONCLUSION

NDDs can impact the child, their family, academic achievements, education, play, peer or social relationships, and cognitive functioning. WM happens to be the most apparent neuropsychological function that is affected by these children. There appear to be no gender differences in this predicament, as seen in children from matched high-SES in India's metropolitan city.

REFERENCES

1. Wilson DM, Gross D. Parents' executive functioning and involvement in their child's education: An integrated literature review. *J Sch Health* 2018;88(4):322–29.
2. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A, Wager TD. The unity and diversity of executive functions and their contributions to complex "frontal lobe" tasks: A latent variable analysis. *Cogn Psychol* 2000;41(1):49–100.
3. Zilles D, Lewandowski M, Vieker H, Henseler I, Diekhof E, Melcher T, et al. Gender differences in verbal and visuospatial working memory performance and networks. *Neuropsychobiology* 2016;73(1):52–63.
4. Ma L, Chang L, Chen X, Zhou R. Working memory test battery for young adults: Computerized working memory assessment. *PLoS One* 2017;12(3): e0175047.
5. Baddeley A. Working memory. *Curr Biol* 2010; 20(4): R136–40.
6. Cowan N. Working Memory Maturation: Can we get at the essence of cognitive growth? *Perspect Psychol Sci* 2016; 11(2): 239–64.
7. Alloway TP, Gathercole SE, Pickering SJ. Verbal and Visuospatial Short-Term and Working Memory in Children: Are They Separable? *Child Dev* 2006;77 :1698–716.
8. Huizinga M, Dolan CV, Van der Molen MW. Age-related change in executive function: Developmental trends and a latent variable analysis. *Neuropsychologia* 2006;44(11):2017–36.
9. Gathercole SE, Alloway TP. Practitioner Review: Short-term and working memory impairments in neurodevelopmental disorders: diagnosis and remedial support. *J Child Psychol Psychiatry* 2006;47(1):4–15.
10. Reiter A, Tucha O, Lange KW. Executive functions in children with dyslexia. *Dyslexia* 2005;11(2):116–31.
11. Cowan N, Alloway T. Development of working memory in childhood. In: Courage M L, Cowan N, editors. *The Development of Memory in Infancy and Childhood*. 1st ed. New York, NY, US: Psychology Press; 2009. pp 303–42.
12. Peng P, Fuchs D. A meta-analysis of working memory deficits in children with learning difficulties: Is there a difference between verbal domain and numerical domain? *J Learn Disabil* 2016;49(1):3–20.
13. Maehler C, Schuchardt K. Working memory in children with specific learning disorders and/or attention deficits. *Learn Individual Differ* 2016;49: 341–47.
14. Conti-Ramsden G, Ullman MT, Lum JA. The relation between receptive grammar and procedural, declarative, and working memory in specific language impairment. *Front Psychol* 2015; 6:1090.
15. Saarinen S, Fontell T, Vuontela V, Carlson S, Aronen ET. Visuospatial working memory in 7-to 12-year-old children with disruptive behavior disorders. *Child Psychiatry Hum Dev* 2015;46(1):34–43.
16. de Vries M, Prins PJM, Schmand BA, Geurts HM. Working memory and cognitive flexibility-training for children with an autism spectrum disorder: a randomized controlled trial. *J Child Psychol Psychiatry* 2015; 56(5): 566–76.
17. Vandenbroucke L, Verschueren K, Desoete A, Aunio P, Ghesquière P, Baeyens D. Crossing the bridge to elementary school: The development of children's working memory components in relation to teacher-student relationships and academic achievement. *Early Child Res Q* 2018;42: 1–10.
18. Thapar A, Cooper M, Rutter M. Neurodevelopmental disorders. *Lancet Psychiatry* 2017;4(4):339–346.
19. American Psychiatric Association. Neurodevelopmental Disorders. In: *Diagnostic and statistical manual of mental disorders* [Internet]. 5th ed. Arlington: VA; 2013. pp 31–86. Available from: <https://doi.org/10.1176/appi.books.9780890425596.dsm01>
20. Thapar A, Rutter M. Neurodevelopmental disorders. In: Thapar A, Pine D, Leckman J F, Scott S, Snowling MJ, Taylor E A, editors. *Rutter's child adolescent psychiatry*. 6th ed. West Sussex, UK: John Wiley and Sons Ltd.; 2015. pp 31–40.

21. World Health Organization. International classification of diseases for mortality and morbidity statistics [Internet]. 11th ed. 2018. Available from: Retrieved from <https://icd.who.int/browse11/l-m/en>
22. Chauhan A, Sahu JK, Jaiswal N, Kumar K, Agarwal A, Kaur J, et al. Prevalence of autism spectrum disorder in Indian children: a systematic review and meta-analysis. *Neurol India* 2019; 67(1):100-4.
23. Rathi SK, Francis MR. Neurodevelopmental disorders among Indian children needs attention. *Electronic Physician* 2012;4(2):493-6.
24. Arora NK, Nair MKC, Gulati S, Deshmukh V, Mohapatra A, Mishra D, et al. Neurodevelopmental disorders in children aged 2-9 years: Population-based burden estimates across five regions in India. *PLoS Med.* 2018; 15(7): e1002615.
25. Farah MJ. The Neuroscience of Socioeconomic Status: Correlates, Causes, and Consequences. *Neuron* 2017;96(1):56-71.
26. Potijk MR, Kerstjens JM, Bos AF, Reijneveld SA, de Winter AF. Developmental delay in moderately preterm-born children with low socioeconomic status: risks multiply. *J Pediatr* 2013;163(5):1289-95.
27. Demetriou EA, Lampit A, Quintana DS, Naismith SL, Song YJC, Pye JE, et al. Autism spectrum disorders: a meta-analysis of executive function. *Mol Psychiatry* 2018;23(5):1198-204.
28. Habib A, Harris L, Pollick F, Melville C. A meta-analysis of working memory in individuals with autism spectrum disorders. Zhang J, editor. *PLoS ONE* 2019;14(4): e0216198.
29. Kasper LJ, Alderson RM, Hudec KL. Moderators of working memory deficits in children with attention-deficit/hyperactivity disorder (ADHD): A meta-analytic review. *Clin Psychol Rev* 2012;32(7):605-17.
30. Rinehart NJ, Cornish KM, Tonge BJ. Gender differences in neurodevelopmental disorders: Autism and fragile x syndrome. In: Biological basis of sex differences in psychopharmacology. Springer; 2010. pp 209-229.
31. Bölte S, Westerwald E, Holtmann M, Freitag C, Poustka F. Autistic Traits and Autism Spectrum Disorders: The Clinical Validity of Two Measures Presuming a Continuum of Social Communication Skills. *J Autism Dev Disord* 2011; 41(1):66-72.
32. Voyer D, Voyer SD, Saint-Aubin J. Sex differences in visual-spatial working memory: A meta-analysis. *Psychon Bull Rev* 2017;24(2): 307-34.
33. Lawson GM, Hook CJ, Farah MJ. A meta-analysis of the relationship between socioeconomic status and executive function performance among children. *Dev Sci* 2018;21(2): e12529.
34. Sarsour K, Sheridan M, Jutte D, Nuru-Jeter A, Hinshaw S, Boyce WT. Family socioeconomic status and child executive functions: The roles of language, home environment, and single parenthood. *J Int Neuropsychol Soc* 2011;17(1):120-32.
35. Ursache A, Noble KG, the Pediatric Imaging, Neurocognition and Genetics Study. Socioeconomic status, white matter, and executive function in children. *Brain Behav* 2016 ;6(10):1-13.
36. Rosenthal M, Wallace GL, Lawson R, Wills MC, Dixon E, Yerys BE, et al. Impairments in real-world executive function increase from childhood to adolescence in autism spectrum disorders. *Neuropsychol* 2013;27(1):13-8.
37. Kohli A, Malhotra S, Mohanty M, Khehra N, Kaur M. Specific learning disabilities in children: deficits and neuropsychological profile. *Int J Rehabil Res* 2005;28(2):165-9.
38. Vinayan KP, Biji V, Thomas SV. Educational problems with underlying neuropsychological impairment are common in children with benign epilepsy of childhood with centrotemporal spikes (BECTS). *Seizure* 2005;14(3):207-12.
39. Kohli A, Kaur M, Mohanty M, Malhotra S. Neuropsychological functioning in specific learning disorders—reading, writing and mixed groups. *J Indian Assoc Child Adolesc Ment Health.* 2006;2(4):112-115.
40. Sinha P, Sagar R, Mehta M. Executive Function in Attention Deficit/Hyperactivity Disorder. *Journal of Indian Association for Child and Adolescent Mental Health.* 2008;4(2):44-49.
41. Kar BR, Tripathi N. Cognitive profiles of children with dyslexia. *Adv Cogn Sci* 2008; 1:341-354.
42. Visalakshi N, Thenmozhi S. Executive functions of children with learning problems. *Indian J Community Psychol* 2011;7(1):51-8.
43. VS S, DIM D, Varghese PK, Vinayan KP. Cognitive profile of children with attention deficit-hyperactivity disorder. *Amrita J Med* 2013;10 (1): 28-32.
44. Gupta SK, Venkatesan S. Efficacy of Training Program on Executive Functions in Children with Learning Disability. *J Behav Soc Sci* 2014;2(2):238-291.
45. Khan A. An investigation into prospective memory in children with developmental dyslexia. *Front Psychol* 2014;5: 1308.
46. Kohli A, Malhotra S, Khehra N. Neuropsychological profile of children with hyperkinetic disorder. *J Indian Assoc Child Adolesc Ment Health.* 2014;10(3):167-93.
47. Nishanimut S P, Padakannaya P. Neurocognitive profile of children with reading disability in Kannada. *Avicenna J Neuro Psycho Physiol* 2014;1(2):46-50.
48. Gupta PK, Sharma V, Sinha U K. Working Memory Functioning in children with Specific Learning Disability [Internet]. [Delhi]: Institute of Human Behavior and Allied Sciences, Delhi; 2015 [cited 2020 Aug 28]. Available from: <http://rgdoi.net/10.13140/RG.2.2.18871.32169>
49. Kumar S, Gore GB. Comparison of phonological processing abilities in children with learning disability in Hindi and English language: A pilot study. *Int J Health Sci Res* 2016;6(12):219-26.
50. Kumar H, Singh P, Sharma P. Working memory and academic attainment in children with mild intellectual disability. *Int J Multidisciplinary Res* [Internet] 2016; Available from: Retrieved from http://eprajournals.com/jpanel/upload/1242am_10.Harish%20Kumar-13.10.16.pdf

51. Kotnala S, Halder S. Working memory, Verbal comprehension, Perceptual reasoning and processing speed in ADHD and normal children: A comparative study. *J Indian Assoc Child Adolesc Ment Health* 2018;14(1):60–79.
52. Malhi P, Singhi P. Adaptive Behavior functioning in children with Autism. *Indian J Pediatr* 2015;82(8):677–81.
53. Devi S. A comparative study on profile analysis of the Binet-Kamat Test of intelligence of children having a mild intellectual disability with and without Down syndrome. *Eur Acad Res* 2014; II (7):9056-72. Available from: <https://pdfs.semanticscholar.org/3132/67626e5fea7c2529cfl6857c6648a4396b2.pdf>
54. Boivin MJ, Kakooza AM, Warf BC, Davidson LL, Grigorenko EL. Reducing neurodevelopmental disorders and disability through research and interventions. *Nat* 2015; 527(7578): S155–60.
55. Rutter M, Pine DS. Diagnosis, diagnostic formulations, and classification. In: Thapar A, Pine DS, Leckman J F, Scott S, Snowling M J, Taylor E A, editors. *Rutter's child and adolescent psychiatry*. 6th ed. West Sussex, UK: John Wiley and Sons Ltd; 2015. pp 17–30.
56. Rutter M, Thapar A. Genetics of autism spectrum disorders. In: Volkmar F R, Rogers SJ, Paul R, Pelphrey K A, editors. *Handbook of Autism and pervasive developmental disorders*. 4th ed. New Jersey: USA: John Wiley and Sons Inc; 2014. pp 411–23.
57. Rommelse NNJ, Geurts HM, Franke B, Buitelaar JK, Hartman CA. A review on cognitive and brain endophenotypes that may be common in autism spectrum disorder and attention-deficit/hyperactivity disorder and facilitate the search for pleiotropic genes. *Neurosci Biobehav Rev* 2011;35(6):1363–96.
58. Kar B, Rao S, Chandramouli BA, Thennarasu K. Growth patterns of neuropsychological functions in Indian children. *Front Psychol* 2011; 2:240.
59. Venkatesan S. Readapted version for 2016: NIMH socioeconomic status scale. Secunderabad: National Institute Mentally Handicapped currently 'National Institute for Empowerment of Persons with Intellectual and Developmental Disabilities'; 2009a.
60. Venkatesan S. Children with developmental disabilities: A training guide for parents, teachers and caregivers. SAGE; 2004.
61. Venkatesan S. Reappraisal of Bombay-Karnatak version of Binet-Simon intelligence scales (1964). *Indian J Clin Psychol* 2002;29(1):72–78.
62. Kamat VV. Measuring intelligence of Indian children. Bombay: Oxford University Press; 1967.
63. Lezak MD, H owieson DB, Bigler E, Trane D. Neuropsychological assessment. 5th ed. Oxford: London: Oxford University Press; 2012.
64. Venkatesan S. Profile analysis on the Binet Kamat test: A comparative study on groups of mentally retarded individuals. [An unpublished pre-doctoral thesis submitted to Bangalore University, Bangalore, as part fulfillment for the award of the degree of Master's in Philosophy (Clinical Psychology) in Department of Clinical Psychology]. Bangalore, India: National Institute of Mental Health and Neurosciences; 1987.
65. Urbaniak GC, Plous S. Research Randomizer (Version 4.0) [Internet]. 2013. Available from: <http://www.randomizer.org/>
66. Gazzaniga MS, Ivry RB, Mangun GR. Cognitive control. In: *Cognitive Neuroscience: The Biology of the Mind* 4th ed. New York: W.W. Norton and Company, Inc.; 2014. pp. 506–557.
67. Richardson JTE. Knox's cube imitation test: A historical review and an experimental analysis. *Brain Cogn* 2005;59(2):183–213.
68. Milner B. Interhemispheric differences in the localization of psychological processes in man. *Br Med Bull* 1971;27(3):272–77.
69. Kanne S, Huber DH. Stanford Binet Intelligence Scales and Revised Versions. In: Kreutzer JS, DeLuca J, Caplan B, editors. *Encyclopedia of Clinical Neuropsychology* [Internet]. SpringerLink (Online service); 2011. pp 2371–7.
70. Smith EE, Jonides J. Storage and executive processes in the frontal lobes. *Sci* 1999;283(5408):1657–61.
71. Kar BR, Rao S L, Chandramouli BA, Thennarasau K. NIMHANS neuropsychological battery for children-manual. Bangalore: NIMHANS Publication Division; 2004.
72. Venkatesan S. Ethical guidelines for bio behavioral research. Mysore: All India Institute of Speech and Hearing; 2009b.
73. Hay AD, Wilson A, Fahey T, Peters TJ. The duration of acute cough in pre-school children presenting to primary care: a prospective cohort study. *Fam Pract* 2003;20(6):696–705.
74. IBM C. IBM SPSS Statistics for Windows. Armonk, NY: IBM Corp; 2015.
75. Meule A. Reporting and interpreting working memory performance in n-back tasks. *Front Psychol* 2017; 8:352.
76. Pek J, Flora DB. Reporting effect sizes in original psychological research: A discussion and tutorial. *Psychol Methods* 2018;23(2):208–25.
77. Fritz CO, Morris PE, Richler JJ. Effect size estimates: current use, calculations, and interpretation. *J Exp Psychol: Gen.* 2012 Feb;141(1):2.
78. Grissom RJ. Probability of the superior outcome of one treatment over another. *J Appl Psychol* 1994;79(2):314–6.
79. Corbett BA, Constantine LJ, Hendren R, Rocke D, Ozonoff S. Examining executive functioning in children with autism spectrum disorder, attention deficit hyperactivity disorder and typical development. *Psychiatry Res* 2009; 166(2–3): 210–22.
80. Delehanty AD, Stronach S, Guthrie W, Slate E, Wetherby AM. Verbal and nonverbal outcomes of toddlers with and without autism spectrum disorder, language delay, and global developmental delay. *Autism Dev Lang Impair* 2018; 3:239694151876476.

81. Henry LA, Botting N. Working memory and developmental language impairments. *Child Lang Teach Ther* 2017;33(1):19–32.
82. Englund JA, Decker SL, Allen RA, Roberts AM. Common cognitive deficits in children with attention-deficit/hyperactivity disorder and autism: Working memory and visual-motor integration. *J Psychoeducational Assess* 2014;32(2):95–106.
83. Hwang Y, Hosokawa T. Working memory in Japanese children with learning disabilities: Assessment using the Japanese short form of the Swanson cognitive processing test. *Jpn J Spec Educ* 2007;44(6):463–72.
84. Schuchardt K, Maehler C, Hasselhorn M. Working memory deficits in children with specific learning disorders. *J Learn Disabil* 2008; 41(6), 514–23.
85. Mammarella IC, Lucangeli D, Cornoldi C. Spatial working memory and arithmetic deficits in children with nonverbal learning difficulties. *J Learn Disabil* 2010;43(5):455–68.
86. Alloway TP, Alloway R. Working memory: Is it the new IQ? *Nat Prec* 2008; 1-1.
87. Mulder H, Pitchford NJ, Marlow N. Processing speed and working memory underlie academic attainment in very preterm children. *Arch Dis Child Fetal Neonatal Ed* 2010;95(4): F267–72.
88. Yerys BE, Wallace GL, Sokoloff JL, Shook DA, James JD, Kenworthy L. Attention deficit/hyperactivity disorder symptoms moderate cognition and behavior in children with autism spectrum disorders. *Autism Res* 2009;2(6):322–33.
89. Williams DL, Goldstein G, Carpenter PA, Minshew NJ. Verbal and Spatial Working Memory in Autism. *J Autism Dev Disord*. 2005;35(6):747–56.
90. Williams DL, Goldstein G, Minshew NJ. The profile of memory function in children with Autism. *Neuropsychol* 2006;20(1):21–9.
91. Montgomery JW. Verbal working memory and sentence comprehension in children with specific language impairment. *J Speech Lang Hear Res* 2000;43(2):293–308.
92. Hille ET, Weisglas-Kuperus N, Van Goudoever JB, Jacobusse GW, Ens-Dokkum MH, de Groot L et al. Functional outcomes and participation in young adulthood for very preterm and very low birth weight infants: the Dutch project on preterm and small for gestational age infants at 19 years of age. *Pediatr* 2007;120(3):587-95.
93. Mutch L, Leyland A, McGee A. Patterns of neuropsychological function in a low-birth weight population. *Dev Med Child Neurol* 2008;35(11):943–56.
94. Fry AF, Hale S. Relationships among processing speed, working memory, and fluid intelligence in children. *Biol Psychol* 2000;54(1–3):1–34.
95. Holmes J, Hilton KA, Place M, Alloway TP, Elliott JG, Gathercole SE. Children with low working memory and children with ADHD: same or different? *Front Hum Neurosci* 2014;8:976.
96. Miller KM, Price CC, Okun MS, Montijo H, Bowers D. Is the n-back task a valid neuropsychological measure for assessing working memory? *Arch Clin Neuropsychol* 2009;24(7):711–17.
97. Vogan VM, Morgan BR, Powell TL, Smith ML, Taylor MJ. The neurodevelopmental differences of increasing verbal working memory demand in children and adults. *Dev Cogn Neurosci* 2016;17: 19–27.
98. Schmidt H, Jogia J, Fast K, Christodoulou T, Haldane M, Kumari V, et al. No gender differences in brain activation during the N-back task: An fMRI study in healthy individuals. *Hum Brain Mapp* 2009;30(11):3609–15.
99. Speck O, Ernst T, Braun J, Koch C, Miller E, Chang L. Gender differences in the functional organization of the brain for working memory. *Neuroreport* 2000;11(11):2581–5.
100. Schweinsburg AD, Nagel BJ, Tapert SF. fMRI reveals alteration of spatial working memory networks across adolescence. *J Int Neuropsychol Soc* 2005;11(5):631–44.
101. Vuontela V, Carlson S, Troberg A-M, Fontell T, Simola P, Saarinen S, et al. Working memory, attention, inhibition, and their relation to adaptive functioning and behavioral/emotional symptoms in school-aged children. *Child Psychiatry Hum Dev* 2013;44(1):105–22.
102. León I, Cimadevilla JM, Tascón L. Developmental gender differences in children in a virtual spatial memory task. *Neuropsychol* 2014;28(4):485–95.
103. Muñoz-Suazo MD, Navarro-Muñoz J, Díaz-Román A, Porcel-Gálvez AM, Gil-García E. Sex differences in neuropsychological functioning among children with attention-deficit/hyperactivity disorder. *Psychiatry Res* 2019;278: 289–93.
104. Nydén A, Hjelmquist E, Gillberg C. Autism spectrum and attention – deficit disorders in girls: Some neuropsychological aspects. *Eur Child Adolesc Psychiatry* 2000;9(3):180–5.
105. Koyama T, Kamio Y, Inada N, Kurita H. Sex differences in WISC-III profiles of children with high-functioning pervasive developmental disorders. *J Autism Dev Disord* 2008; 39(1):135-141.
106. Grissom NM, Reyes TM. Let's call the whole thing off: evaluating gender and sex differences in executive function. *Neuropsychopharmacol* 2019;44(1):86–96.
107. Gabriels RL, Ivers BJ, Hill DE, Agnew JA, McNeill J. Stability of adaptive behaviors in middle-school children with autism spectrum disorders. *Res Autism Spectr Disord* 2007;1(4):291–303.
108. Griffith GM, Hastings RP, Nash S, Hill C. Using Matched Groups to Explore Child Behavior Problems and Maternal Well-Being in Children with Down Syndrome and Autism. *J Autism Dev Disord* 2010;40(5):610–9.
109. Mouga S, Almeida J, Café C, Duque F, Oliveira G. Adaptive profiles in Autism and other neurodevelopmental disorders. *J Autism Dev Disord* 2015;45(4):1001–12.
110. Shevell M, Majnemer A, Platt RW, Webster R, Birnbaum R. Developmental and functional outcomes at school age of preschool children with global developmental delay. *J Child Neurol* 2005;20(8):648–53.

111. Mougá S, Café C, Almeida J, Marques C, Duque F, Oliveira G. Intellectual profiles in the Autism spectrum and other neurodevelopmental disorders. *J Autism Dev Disord* 2016;46(9):2940–55.
112. Weyandt LL, Gudmundsdottir BG. Developmental and neuropsychological deficits in children with ADHD. In: Barkley RA, editor. *Attention-Deficit Hyperactivity Disorder: A handbook for diagnosis and treatment*. The Guilford Press; 2015. pp 116–39.
113. Dennis M, Francis DJ, Cirino PT, Schachar R, Barnes MA, Fletcher JM. Why IQ is not a covariate in cognitive studies of neurodevelopmental disorders. *J Int Neuropsychol Soc* 2009;15(3):331–43.
114. Frith U, Happe F. Why specific developmental disorders are not specific: On-line and developmental effects in Autism and dyslexia. *Developmental Sci* 1998;1(2):267–72.
115. Diaz-Asper CM, Schretlen DJ, Pearlson GD. How well does IQ predict neuropsychological test performance in normal adults? *J Inter Neuropsych Soc* 2004; 10(01):82-90.
116. Rose E, Bramham J, Young S, Paliokostas E, Xenitidis K. Neuropsychological characteristics of adults with comorbid ADHD and borderline/mild intellectual disability. *Res Dev Disabil* 2009;30(3):496–502.
117. Neece CL, Baker BL, Lee SS. ADHD among adolescents with intellectual disabilities: Pre-pathway influences. *Res Dev Disabil* 2013;34(7):2268–79.
118. van Iterson L, de Jong PF. Development of verbal short-term memory and working memory in children with epilepsy: Developmental delay and impact of time-related variables. A cross-sectional study. *Epilepsy Behav* 2018; 78:166–74.

Acknowledgements – Nil

Conflict of Interest – Nil

Funding – Nil