**International Journal of Biology, Pharmacy
and Allied Sciences (IJBPAS)**
'A Bridge Between Laboratory and Reader'

www.ijbpas.com

**HAEMOGLOBIN VARIANTS AND HEMOGLOBINOPATHIES IN HAIL
POPULATION – KSA: RETROSPECTIVE STUDY IN A CLINICAL DIAGNOSTIC
LABORATORY**

FAWAZ D. ALSHAMMARI

Department of Clinical Laboratory Sciences, College of Applied Medical Sciences, University of
Hail, KSA

***Corresponding Author: Fawaz D. Alshammari: E Mail: fawazalshammari2016@gmail.com**

Received 5th Sept. 2018; Revised 4th Oct. 2018; Accepted 7th Nov. 2018; Available online 1st Feb. 2019

<https://doi.org/10.31032/IJBPAS/2019/8.2.4645>

ABSTRACT

Hemoglobins have a basic job in the body, as it is the protein in red platelets (RBCs) that conveys oxygen from the lungs to the tissues of the body. In that capacity, anomalous dimensions of hemoglobins or strange kinds of hemoglobins can result in genuine infection. We should take a gander at the ordinary scopes of hemoglobins in grown-ups in youngsters, conditions that reason either a low hemoglobins or a high hemoglobins level, and the tests for and hugeness of genetic strange hemoglobins. Press lack is the most well-known micronutrient insufficiency and can diminish the creation of hemoglobins and subsequently cause press inadequacy pallor (which is a correctable and remediable condition). Frailty caused from Iron inadequacy is one of most wide-spread ailments on the planet. Kids in school age and extraordinarily in the period of immaturity will come to encounter the risk of frailty dependent on iron insufficiency. In this age paleness will have a quick growth. Youths' have to various sustenances, especially press, and will increment however the need is a lot higher among females. Understudies with iron insufficiency will get tainted with microorganisms because of decline of body quality. Insight Q will be lower Iron-deficiency anemia can likewise be caused by a less than stellar eating routine or by certain intestinal infections that influence how the body retains press. The condition is ordinarily treated with iron enhancements. In females of childbearing age, the most successive reason for iron-inadequacy pallor is loss of iron in blood because of critical feminine cycle or pregnancy. In Saudi Arabia the general nation predominance of iron inadequacy frailty was 30– 56%. The examination was driven in Hail organized in

the Saudi district. It has a zone of 103,887 km² and a people of 527,033. All blood got from people were screened by High Performance Liquid Chromatography (HPLC). Current investigation included male and female (1085) members. Among them 72 (6.64%) were sickly and 1013 (93.36%) were non-frail. The present examination, we reason that larger part of the subjects screened were non-sickly.

Keywords: Anemia, HPLC, infection, Saudi Arabia, serum

1. INTRODUCTION

Hemoglobin is made out of four polypeptide chains, which in grown-ups comprise of two alpha (α) globin chains and two beta (β) globin chains (for example α₂β₂). Every polypeptide has a haem prosthetic gathering connected, where every haem can tie one oxygen atom - so there are four haem bunches for each hemoglobin particle those together tie four oxygen particles. There are seven sorts of hemoglobin particles all through a human's life. Four when you are an incipient organism, one once you form into a hatchling and afterward as a grown-up you have two. Which are Embryonic (Hemoglobin Gower II (α₂ε₂), Hemoglobin Portland I (ζ₂γ₂), Hemoglobin Portland II (ζ₂β₂)), Fetal Hemoglobin (Once a fetus forms into a baby and the four sorts of embryonic hemoglobin atoms vanish they are supplanted by Hemoglobin F (α₂γ₂)), Adult (Hemoglobin A (α₂β₂), Hemoglobin A₂ (α₂δ₂)). Anemia caused from Iron lack is one of most wide-spread sicknesses on the planet. Youngsters in school age and extraordinarily in the period of youth will

come to encounter the risk of frailty dependent on iron insufficiency. In this age weakness will have a quick growth. Young peoples have to various nourishments, especially press, and will increment however the need is a lot higher among females.

Wholesome frailty happens because of deficient admission of supplements by cells. Among the most vital supplements whose insufficiency can prompt wholesome paleness are press, folic corrosive, nutrient B12, nutrient B6, nutrient C and protein. Press insufficiency paleness is a standout amongst the most continuous medical issues on the planet (Soleimani, N. what's more, Abbaszadeh, N., 2011). The World Health Organization as of late announced that 1.62 billion of the total populace is iron deficient. The rate among understudies is 25.4% and in preschool age youngsters pallor achieves its most astounding level of 47.4 (Benoist, B., et. al., 2008). Press inadequacy paleness represents 75% of a wide range of sickliness in the third world, influencing 30% of populace (Soleimani, N., 2011). In females

of childbearing age, the most continuous reason for iron-insufficiency pallor is loss of iron in blood because of critical period or pregnancy. Press insufficiency frailty can likewise be caused by a horrible eating routine or by certain intestinal maladies that influence how the body ingests press. The condition is ordinarily treated with iron enhancements.

Writing about iron deficiency in young people and youth is rare, when contrasted with that concentrating on ladies and youngsters. In Saudi Arabia the general nation predominance of iron insufficiency sickliness was 30– 56% (Verster, A. also, van der Pols, J.C., 1995). A cross-sectional investigation led in Riyadh City among students demonstrated that the predominance was 40.5% among female young people (16– 18) years of age (Musaiger, A.O., 2002).

In addition, Anemia is a general helpful issue affecting around 1.62 billion sufferers or roughly a fourth of the total masses in both made and making states (WHO, 2008; Peyrin-Biroulet, L., et al., 2015). A broad assortment of sorts of whiteness exist, and among these, crush deficiency squeeze insufficiency (IDA) is the most expansive (Anand, T., et al., 2014; Shander, An., et al., 2014). Frankly, IDA has achieved pandemic estimations in various making nations

(Sirdah, M.M., et al., 2014) and is beginning at now the most predominant micronutrient inadequacy on earth (Chandyo, R.K., et al., 2016; Abu-Ouf, N.M., et al., 2015). Press insufficiency pallor has to a remarkable degree opposite ramifications for influenced people, who are in risk of disturbed headway and insightful improvement, chop down mental and engine work, poorer work limit, and a for the most part chopped down near and dear satisfaction (Abbaspour, N., et al., 2014). Press deficiency is caused by a put off ominous imbalance between an individual's dietary affirmation of iron and their body's physiological interest (Abbaspour, N., et al., 2014; Hurrell, R. moreover, Egli, I., 2010). Particular non-modifiable and modifiable parts apply an effect on a person's iron uniformity, either in blend or alone (Alquaiz, A.M., et al., 2013), going from socio-estimation characteristics (checking the person's age, sex, marital status, estimation of getting ready, pay, and ethnicity) to the aggregate and nature of the sustenance and refreshments they eat up, their psychological and physical success, the solution they take, any peculiarities they have, and their inborn cosmetics (Alquaiz, A.M., et al., 2013; Al-Quaiz, J.M., 2001; Alquaiz, A.J., et al., 2015).

It is especially considerable for ladies that if IDA isn't expediently perceived and treated, it has significant built up impacts, including negative maternal and neonatal results, and a brought risk of disappointment up in later life (Cao, C. besides, O'Brien, K.O., 2013). Data on IDA rates and the hazard factors related with the condition is unprecedented for female Saudi Arabian school understudies. In the essential basic past examination which could be found, (Al Hassan, 2015) revealed a typical unavoidability of IDA of 64% for an instance of female school understudies of Saudi nationality, yet their examination dismissed any thought of the potential hazard factors related with IDA. On account of this opening in the earlier research, the target of this examination was done to pick and outline the measure of viral pollutions information revealed in Saudi Arabia

2. MATERIALS AND METHOD

2.1 Materials and sources

Preliminary materials were procured from different sources and used in the present work. The fine manufactured inventions, units and mixes are referenced close by their sources as and where required in the substance. The examination was driven in Hail arranged in the Saudi region. It has an area of 103,887 km² and a people of 527,033. All serum got from individuals were

screened by High Performance Liquid Chromatography.

2.2 Method

2.2.1 Data collection

The present research was led on the grounds of the hail region of the Saudi Arabia. The information was gathered at different areas which had been explicitly decided for their accommodation to understudies over all the medical college's universities. A group containing a restorative specialist, a dietician, medical attendants, and research facility professionals played out the dietary and anthropometric information accumulation and blood test extraction.

2.2.2 Blood sample collection

Around the same time when the dietary and anthropometric information were gathered, an attendant took a 10-ml test of venous blood from every member for haematological also, biochemical screening tests. The research centre experts in the exploration group utilized 2 kinds of vacutainers to perform venepunctures; one contained EDTA for haematological tests, counting haemoglobin (Hgb), haematocrit, mean corpuscular haemoglobin, mean corpuscular volume, and mean corpuscular haemoglobin focus. These were performed utilizing a Beckman coulter LH750 machine which was consistently checked and aligned amid the

investigation as indicated by the standard quality confirmation conventions set toward the start of the exploration.

2.2.3 High Performance Liquid Chromatography (HPLC)

Superior fluid chromatography or high-weight fluid chromatography (HPLC) is a chromatographic strategy that is utilized to isolate a blend of mixes in investigative science to distinguish, evaluate or filter the individual segments of the blend. HPLC has been broadly utilized for a considerable length of time as a logical strategy for test division in pharmaceutical quality control and life science inquire about. Because of the power and adaptability of this system, various applications have been embraced for schedule use in the clinical lab.

HPLC is a detachment method in which the versatile stage is a watery arrangement or a natural dissolvable that is miscible with water, and the mixes of intrigue are disintegrated in water or a natural dissolvable, for example, methanol. The real preferred standpoint of the procedure is the adaptability it permits in picking the stationary furthermore, portable stages that best discrete the specific example parts. The greater part of the kinds of sections utilized in HPLC is invert stage; around a quarter are typical stage; about 14% are particle trade

sections. Getting high goals at high stream rates is the key necessity of any HPLC framework. Partition of the diverse segments of a blend is accomplished by enabling the example to relocate through the section at chosen trial conditions. Discrete groups are shaped since the diverse segments of the blend relocate through the section at various rates. The VARIANT a completely mechanized superior fluid chromatography (HPLC) framework utilizes twofold wave length identification (416 and 690 nm). This cationic-trade segment chromatography empowers subjective determinations of typical and strange haemoglobins in 6.5 minutes on a haemolyse arranged from 5 ml of venous blood.

2.2.4 Statistical analysis

Statistical analysis was carried out using the SPSS computer package version 21.0 (SPSS Inc., Chicago, IL, USA).

3. RESULTS

3.1 Subject information

Current study included one thousand eighty five (N = 1085) male and female participants. Among them 72 (6.64%) were anemic and 1013 (93.36%) were non-anemic (Table 1 and Figure 1).

3.2 Clinical parameters

3.2.1 Frequency distribution of anemic subjects as a function of range in clinical parameters namely HBA, HBA2, and HBF

Results of frequency distribution of anemic subjects (N = 72) revealed that 98.61%, subjects showed normal values, where as 1.3% and none showed below and above normal values for HBA (Table 2 and Figure 2). Among the all subjects 100%, subjects showed normal values, whereas none showed below and above normal values for HBA2 (Table 2 and Figure 2). Similarly, 20.83% subjects showed normal values, where as 76.38% and 2.77% showed below and above normal values for HBF (Table 2 and Figure 2).

3.2.2 Frequency distribution of non-anemic subjects as a function of range in clinical parameters namely HBA, HBA2, and HBF

Results of frequency distribution of non-anemic subjects (N = 1013) revealed that 97.03%, subjects showed normal values, where as 2.27% and 0.69% showed below and above normal values for HBA (Table 3 and Figure 3). Among the all subjects 98.12%, subjects showed normal values, whereas 0.09% and 1.77% showed below and above normal values for HBA2 (Table 3 and Figure 3). Similarly, 20.13% subjects showed normal values, where as 77.09% and 2.76% showed below and above normal values for HBF (Table 3 and Figure 3).

Table 1: Baseline characteristics	
Total subjects	N = 1085
Anemic/Non-anemic	
Anemic	N = 72 (6.64%)
Non-anemic	N = 1013 (93.36%)

Table 2: Frequency distribution of anemic subjects (N = 72) as a function of range in clinical parameters namely HBA, HBA2, and HBF	
HBA	
Normal range (96-98%)	N = 71 (98.61 %)
Below 96%	N = 01 (1.3 %)
Above 98%	Nil
HBA2	
Normal range (1-3.2%)	N = 72 (100 %)
Below 1%	Nil
Above 3.2%	Nil
HBf	
Normal range (0.5-1%)	N = 15 (20.83 %)
Below 0.5%	N = 55 (76.38 %)
Above 1%	N = 2 (2.77 %)

Table 3: Frequency distribution of non-anemic subjects (N = 1013) as a function of range in clinical parameters namely HBA, HBA2, and HBF	
HBA	
Normal range (96-98%)	N = 983 (97.03 %)
Below 96%	N = 23 (2.27 %)
Above 98%	N = 07 (0.69 %)
HBA2	
Normal range (1-3.2%)	N = 994 (98.12 %)
Below 1%	N = 01 (0.09 %)
Above 3.2%	N = 18 (1.77 %)
HBf	
Normal range (0.5-1%)	N = 204 (20.13 %)
Below 0.5%	N = 781 (77.09 %)
Above 1%	N = 28 (2.76 %)

4. DISCUSSION

In spite of the fact that sickliness in the ladies and youngsters has been considered broadly in many parts of the world, there is a deficiency of data on its commonness in the Saudi Arabia Population. Current examination included one thousand eighty five (N = 1085) male and female members. We have presumed that 72 (6.64%) were sickly and 1013 (93.36%) were non-pale. Among Saudi Arabian patients around 36% of hemoglobinopathies cases were β -thalassemia minor (Al-saeed, A.H., 2012). Further, in a southern city of Iran, Movahed et al., in a populace with hemoglobinopathies revealed the nearness of 70.2% cases with β -thalassemia minor that is higher than that in our examined populace (Movahed, An., et al., 2008). As indicated by Abolghasemi et al., report the thalassemia is increasingly common in the northern Caspian Sea drift, southern Persian Gulf and Oman Sea coasts regions of the nation (Abolghasemi, H., et al., 2007) that is in concurrence with the consequences of Movahed et al., 2008.

In our finding the recurrence appropriation of weak subjects (N = 72) uncovered that 98.61%, subjects demonstrated ordinary qualities, where as 1.3% and none appeared beneath or more typical qualities for HBA. Among the all subjects 100%, subjects

indicated typical qualities, though none appeared underneath or more ordinary qualities for HBA2. Additionally, 20.83% subjects indicated ordinary qualities, where as 76.38% and 2.77% appeared beneath or more typical qualities for HbF.

Among haemoglobin subtypes, HbF values did not demonstrate any critical change after iron treatment. This is like past investigations by El-Agouza, I., et al. (2002) and Wasi, P., et al. (1968). There have been clashing reports of the impact of iron lack on HbA2 levels. Barely any creators have detailed an altogether lower HbA2 levels in patients with attendant BTT and iron insufficiency contrasted with those with uncomplicated BTT (Alperin, J. B., et al., 1977; El-Agouza, I., et al. 2002 and Wasi, P., et al., 1968) while others have neglected to inspire such a distinction (Kurlekar, N. also, Mehta, B. C., 1981; Passarello, C., et al., 2012). One reason of these discrepant outcomes has all the earmarks of being the differences in cutoff dimension of serum ferritin into characterize press insufficiency. Passarello et al. (2008) thought about $30\mu\text{g/L}$ as cut off for iron insufficiency, which as we would like to think isn't appropriate since this dimension characterizes negative iron equalization and not press lack (Adamson, J. W., 2007). A few theories have been advanced for the reduced

HbA2 levels in iron insufficiency. Wasi et al. (1968) have proposed that beta chains might be more focused than delta chains in heme restricting prompting less HbA2 arrangement in heme insufficiency. Absence of iron may likewise meddle with delta chain union.

Based on these discovering we considered that recurrence appropriation of non-weak subjects out of 1013 uncovered that 97.03%, subjects demonstrated typical qualities, where as 2.27% and 0.69% appeared beneath or more ordinary qualities for HBA. Among the all subjects 98.12%, subjects demonstrated ordinary qualities, while 0.09% and 1.77% appeared beneath or more typical qualities for HBA2. Essentially, 20.13% subjects indicated typical qualities, where as 77.09% and 2.76% appeared underneath or more ordinary qualities for HBF.

5. CONCLUSION

From the results of current study, we conclude that majority of the subjects screened were non-anemic. Results of frequency distribution of anemic as well as non-anemic subjects revealed that majority of the subjects showed normal values for HBA and HBA2. However, in case of both the groups the values of HBF in most subjects fall below the minimum cut off. The present work has set a benchmark for conceivable future research as randomized preliminaries

which would manufacture more prominent comprehension of this medical problem what's more, would likewise bolster the advancement of a solid and appropriate general wellbeing arrangement which can effectively handle IDA. The discoveries recommend that engaged instruction and mindfulness systems are expected to enhance wholesome propensities by empowering the utilization of rich dietary iron sources and by bringing issues to light of the nourishment and beverages which encourage or prevent the bio accessibility of iron

6. REFERENCES

- [1] Peyrin-Biroulet, L., Williet, N., Cacoub, P. (2015). Guidelines on the diagnosis and treatment of iron deficiency across indications: a systematic review. *Am J Clin Nutr.* **102**:1585–1594.
- [2] Anand, T., Rahi, M., Sharma, P., Ingle, G.K. (2014). Issues in prevention of iron deficiency anemia in India. *Nutrition.* **30**:764–770.
- [3] Sirdah, M.M., Yaghi, A., Yaghi, A.R. (2014). Iron deficiency anemia among kindergarten children living in the marginalized areas of Gaza Strip, Palestine. *Rev Bras Hematol Hemoter.* **36**:132–138.

-
- [4] Chandyo, R.K., Henjum, S., Ulak, M., *et al.* (2016). The prevalence of anemia and iron deficiency is more common in breastfed infants than their mothers in Bhaktapur, Nepal. *Eur J Clin Nutr.* **70**:456–462.
- [5] Abu-Ouf, N.M., Jan, M.M. (2015). The impact of maternal iron deficiency and iron deficiency anemia on child's health. *Saudi Med J.***2**:146–149.
- [6] Abbaspour, N., Hurrell, R., Kelishadi, R. (2014). Review on iron and its importance for human health. *J Res Med Sci.***19**(2):164–174.
- [7] Hurrell, R., Egli, I. (2010). Iron bioavailability and dietary reference values. *Am J Clin Nutr.***91**:1461S–1467S.
- [8] Alquaiz, A.M., Gad, M.A., Khoja, T.A., *et al.* (2013). Prevalence of anemia and associated factors in child bearing age women in Riyadh, Saudi Arabia. *J Nutr Metab.* 636585.
- [9] Al-Quaiz, J.M. (2001). Iron deficiency anemia. A study of risk factors. *Saudi Med J.***22**:490–496.
- [10] Alquaiz, A.J., Khoja, T.A., Alsharif, A., *et al.* (2015). Prevalence and correlates of anaemia in adolescents in Riyadh city, Kingdom of Saudi Arabia. *Public Health Nutr.***18**:3192–3200.
- [11] World Health Organization. Worldwide Prevalence of Anemia 1993 to 2005. Geneva, Switzerland: WHO; 2008.
- [12] Shander, A., Goodnough, L.T., Javidroozi, M., *et al.* (2014). Iron deficiency anemia—bridging the knowledge and practice gap. *Transfus Med Rev.***28**:156–166.
- [13] Cao, C., and O'Brien, K.O. (2013). Pregnancy and iron homeostasis: an update. *Nutr Rev.* **71**:35–51.
- [14] Al-Hassan, N.N. (2015). The prevalence of iron deficiency anemia in Saudi University female students. *J Microsc Ultrastruct.***3**:25–28.
- [15] Mehdi, S.R., Pophali, A. and Al-Abdul Rahim, K.A. (2000). Prevalence of hepatitis B and C and blood donors. *Saudi medical journal*, **21**(10): 942-944.
- [16] El-Agouza, I., Abu Shahla, A. and Sirdah, M. (2002). The effect of iron deficiency anaemia on the levels of haemoglobin subtypes: possible on sequences for clinical diagnosis. *Clinical and Laboratory Haematology*, **24**(5): 285-289.
-

-
- [17] Wasi, P., Disthasongchan, P. and Nakorn, S. (1968). "The effect of iron deficiency on the levels of hemoglobins A2 and E. *The Journal of Laboratory and Clinical Medicine*, **71(1)**: 85-91.
- [18] Alperin, J. B., Dow, P. A. and Petteway, M. B. (1977). Hemoglobin A2 levels in health and various hematologic disorders, *American Journal of Clinical Pathology*, **67(3)**: 219-226.
- [19] Kurlekar, N. and Mehta, B. C., (1981). Haemoglobin A2 levels in iron deficiency anaemia, *Indian Journal of Medical Research*, **73(1)**: 77-81.
- [20] Passarello, C., Giambona, A., Cannata, M., Vinciguerra, M., Renda, D. and Maggio, A. (2012). Iron deficiency does not compromise the diagnosis of high HbA2 thalassemia trait, *Haematologica*, **97(3)**: 472-473.
- [21] Adamson, J. W. (2007). Iron deficiency and other hypo proliferative anemia's: introduction, in Harrison's Principles of Internal Medicine, A. S. Fauci, D. L. Kasper, E. Braunwald et al., Eds., McGraw Hill, NewYork, NY, USA, 17th edition.
- [22] Soleimani, N., and Abbaszadeh, N. (2011). Relationship between anaemia, caused from the iron deficiency, and academic achievement among third grade high school female students. *Procedia: Soc Behav Sci*. **29(0)**:1877-84.
- [23] Benoist, B., McLean, E., Egll, I., Cogswell, M. (2008). Worldwide prevalence of anaemia 1993-2005: WHO global database on anaemia. World Health Organization.
- [24] Soleimani, N. (2011). Relationship between anaemia, caused from the iron deficiency, and academic achievement among third grade high school female students. *Procedia: Soc Behav Sci*. **29**:1877-1884.
- [25] Verster, A., and van der Pols, J.C. (1995). Anaemia in the Eastern Mediterranean region. *East Mediterr Health J*. **1(1)**: 64-79.
- [26] Musaiger, A.O. (2002). Iron deficiency anaemia among children and pregnant women in the Arab Gulf countries: the need for action. *Nutr Health*. **16(3)**:161-171.
- [27] Al-saeed, A.H. (2012). Prevalence of Hemoglobinopathy Disorders in Adult Patients Sent for Diagnosis of
-

-
- Anemia in Saudi Arabia. *Genet Test and Mole Biomarkers*, **16**: 25-29.
- [28] Movahed, A., Obeidi, A., Khamisipour, G.R. (2008). Prevalence of hemoglobinopathies and their associations with different types of hemoglobin and mean cell volume in the pre university students of Bushehr, 2007. *South Med J* **12**: 54-59.
- [29] Abolghasemi, H., Amid, A., Zeinali, S., Radfar, M. H., Eshghi, P., Rahiminejad, M. S., Ehsani, M. A., Najmabadi, H., Akbari, M. T., Afrasiabi, A., Akhavan-Niaki, H., Hoorfar, H. (2007). Thalassemia in Iran: epidemiology, prevention, and management. *J Pediatr Hematol Oncol.*, **29**:233–238.
-