**International Journal of Biology, Pharmacy
and Allied Sciences (IJBPAS)**

'A Bridge Between Laboratory and Reader'

www.ijbpas.com

SIGNIFICANCE AND IMPORTANCE OF ACTINOMICETES AS AN ANTIBIOTIC: A REVIEW

**VALA P¹, UPADHAYAY D², MARCHAWALA F², BHATTACHARYA I² AND
ANDHARE P^{2*}**

1: Student, M. Sc. Microbiology, Parul Institute of Applied Sciences, Parul University, Post
Limda, Waghodia, Gujarat, 391760

2: Assistant Professor, Parul Institute of Applied Sciences, Parul University, Post Limda,
Waghodia, Gujarat, 391760

***Corresponding Author: E Mail: Dr. Prasad Andhare: prasad.andhare82145@paruluniversity.ac.in; Tel:
+918200614350**

Received 19th Jan. 2021; Revised 22nd Feb. 2021; Accepted 20th March 2021; Available online 1st April 2021

<https://doi.org/10.31032/IJBPAS/2021/10.4.1014>

ABSTRACT

Actinomycetes are generally gram-positive microorganisms and they are generally found in soil, water and marine water. Actinomycetes have different separated ecology and characteristics. Actinomycetes have many different classes. Recent studies of actinomycetes have widely separated into different ecology and soil is best source. This actinomycetes developed numerous bioactive compound which are used in prepare antibiotics this compound act as antibacterial therapeutic agent. The actinomycetes produces enzymes like pectinase, lipase, amylase, cellulase, agarases. The main aim of this review to identify new antibiotics against pathogens.

Keywords: Actinomycetes, Antibiotic, antimicrobial Activity

INTRODUCTION

Actinomycetes are gram-positive, free living, saprophytic bacteria, broadly dispensed in soil, water, and colonizing plants. From the 22,500 biologically lively compounds that have received shape

microbes, 45% are produced through actinomycetes, 38% through fungi, and 17% through unicellular bacteria [1]. The critical genera of actinomycetes are. Actinomycetes from the genera *Actinoplanes*,

Streptomyces, *Nocardia*, *Microonospora*, *Rhodococcus*, *Actinomadura*, *Streptoporangium* and *Gordona* were suggested to supply some of broad-spectrum antibiotics [2]. Actinomycetes are gram-positive microorganism with excessive guanine+cytosine content material of over 55% of their DNA. Actinomycetes have filamentous nature, branching pattern, and conidia formation, which can be just like the ones of fungi. Streptomyces are the dominant of all actinomycetes. A massive quantity of actinomycetes had been remoted and screened from soil withinside the beyond numerous decades, accounting for 70%-80% of applicable secondary metabolites to be had commercially [3]. Actinomycetes are the maximum extensively divided organization of microorganisms in nature which overall inhabit the soil. They have provided many vital bioactive compounds of high industrial cost and remain routinely screened for brand new bioactive compounds. These searches had been remarkably a hit and about thirds of evidently occurring antibiotics, together with a lot of medical importance, had been remoted from actinomycetes. According to the WHO, over-prescription and the incorrect use of antibiotics has brought about the era of antibiotic resistance in lots of bacterial pathogens. Nowadays, the drug against lines of pathogen emerges extra fast than the price of discovery of new tablets

and antibiotics. Because of this, many scientists and pharmaceutical enterprise have actively worried in isolation and screening of actinomycetes from one of a kind untouched habitat, for his or her manufacturing of antibiotics [4]. The large-scale screening of adverse actinomycetes beneath impartial situation has caused the invention of surely all present-day clinical antibiotics. During screening, the lines of actinomycetes that did now no longer display interest in impartial environments had been in all likelihood disregarded. Modern screening packages must don't forget the applicable traits of the actinomycetes and must additionally consist of opportunity situations for antibiotic isolation [5].

ACTINOMYCETES:

Actinomycetes were formerly thinking of as bacteria in the guise of fungi or vice versa. It was indeed an opposition to class them. several scientists considered them as Eubacteriales or excessive bacteria, while some thought them to belong to Hypomycetes or lower fungi. Selman Waksman suggested that these microorganisms could be grouped individually [6].

Actinomycetes, such as true bacteria (Eubacteria) are prokaryotes. Their growth (prothallus) is characterized by the development of usually branching threads and rods, frequently giving rise to a typical

mycelium, which is unicellular, especially during the early stages of growth [7]. Generally different the hyphae may turn septate under particular situation. The mycelium is prostrate, i.e., vegetative, and growing in the substrate, or aerial, when a special mycelium is produced above the vegetative growth [8]. Strains of streptomycetes are characterized by the existence of an aerial mycelium below laboratory conditions. On the other hand, the aerial mycelium of nocardia, often invisible to the naked eye, may be altogether absent. Sometimes it may consist of few short strands, which sometimes look like mere granules. Actinomycetes reproduce through special sporulating bodies or from parts of the vegetative mycelium [9].

IMPORTANCE OF ACTINOMYCETES:

The Actinomycetes, forming soil microflora have gained the considerable importance in lately years as producers of therapeutic substances. Many of the Actinomycetes could synthesize metabolites which hinder the growth of bacteria; these are called antibiotics, and, although harmful to bacteria are more or less harmless when presence into the human or animal body. Antibiotics have in modern times great therapeutical and industrial value [10].

The past decagon has seen significant interest in the Actinomycetes as producers

of antibiotic substances. The success uses in chemotherapy of streptomycin, chloramphenicol (Chloromycetin is the trade name of this substance), aureomycin and terramycin all metabolites of the Actinomycetes, has move the search for new Actinomycetes and new antibiotics between the Actinomycetes [11].

The significance of huge repertory of enzymes from actinomycetes and their ability in changing chemical catalysts is discussed. Successful commercialization of those enzymes is an essential step closer to revolutionizing “in experienced technology” Reduction with inside the price of enzyme manufacturing is verified via way of means of manufacturing of endoglucanases from *Streptomyces* sp. on low-price substrates. Such low-price manufacturing projects may be prolonged to different enzymes and metabolites [12]. Actinomycetes were constantly studied and hired for manufacturing of enzymes like that amylases, cellulases, proteases, chitinases, xylanases and pectinase. This evaluation summarized the manufacturing of industrially essential enzymes through Actinomycetes and consultant examples of those enzymes, their traits and commercial makes use of packages in biomedicine, food, detergent, pulp and paper, agriculture, fabric and waste management [13].

INDUSTRIAL SIGNIFICANT OF ACTINOMYCETES:

Actinomycetes are well accepted as they manufacture primary and secondary metabolites that are of economic significance. Actinomycetes produce enzymes such as lipase, cellulases and amylase which are used in industrial fermentation processes. They produce several valuable antibiotics including, amphotericin, neomycin, vancomycin, gentamicin, tetracycline, erythromycin, nystatin, novobiocin and chloramphenicol. Actinomycetes are also used as plant growth promoting agents. They produce biopesticide agents used to influence pests in farmlands. Actinomycetes also have application in bioremediation.

Actinomycetes produce protease enzyme which is used as anti-inflammation agents and in cancer treatment. Actinomycetes produce enzymes that have application in wine presentation [14].

ANTIBIOTICS PRODUCER ACTINOMICETES:

Interest in the direction of the sphere have been normally increasing, even though on occasion declining, hobby and the entire tale indicates a few cyclic functions with successes and screw ups and developed round converting medical wishes and new permitting technology. The current technology of antimicrobial chemotherapy started in 1929, with Fleming's discovery of the effective bactericidal substance, penicillin and Domagk's discovery in 1935

of artificial chemicals (sulfonamides) with broad antimicrobial activity [19]. Actinomycetes are nicely called an inexhaustible supply for antibiotics. Most of the regarded antimicrobials these days had been at first reported from actinomycetes, particularly from the genus *Streptomyces*. The produced materials encompass all essential drug instructions utilized in clinics these days, which includes β -lactams, tetracyclines, macrolides, aminoglycosides, or glycopeptides. In addition to that, the golden technology of antibiotic discovery has been over for pretty some time and the antibiotic pipeline became feared to ruddy. In 2002, the primary *Streptomyces* genome series became published. In addition to that, metagenomic statistics imply that there are a ways extra capacity antibiotic manufacturer in nature looking forward to isolation and investigation [15].

ANTIBIOTICS DISCOVERY:

Several scientific observations since late nineteenth century fueled speculation on the antagonistic nature of certain microbial metabolites on the growth of other bacteria. The first knowledge of some probable antibiotic product from the actinomycetes dates to 1875. Ferdinand Cohn named an organism he found in the tear duct of the human eye as *Streptothrix foersteri*. The French bacteriologist, Duchesne wrote on the antagonism between fungi and bacteria

in a thesis published in 1897. In 1928, Alexander Fleming discovered that the mould *Penicillium* inhibited bacterial growth. Although Fleming discovered Penicillin, the name was given to the antibiotic from the producer, *Penicillium*, it was Florey and Chain, who successfully managed to purify Penicillin in 1940, which paved the way for its commercial use. Yet, the decisive era when innumerable antibacterial would be unveiled was still waiting in the wings [16]. Professor Cook, the botanist, was indeed familiar with this group of microbes, but merely to the extent that they were

causative agents of potato scab. Moreover, he considered them as fungi, and not bacteria. Waksman, caught between and betwixt, fungi and bacteria, delved deeper in understanding these microbes. He felt that these organisms could be differentiated from both bacteria and fungi. After Waksman was awarded the Doctorate of Philosophy in 1918, he moved over to Rutgers University from University of California, Berkeley. It was at Rutgers that Waksman and his team discovered several antibiotics [17].

INDUSTRIAL IMPORTANT GENERA OF ACTINOMYCETES (Table 1)

Table 1: Representation of industrially important actinomycetes producing antibiotics

S. No	Industrially important genera of Actinomycetes	Representatives of antibacterial compounds	Reference
1	<i>Saccharopolyspora</i>	<i>Sporaricin, Saccharocin</i>	[18].
2	<i>Saccharothrix</i>	<i>Saccharocarcin A</i>	[19].
3	<i>Sebekia</i>	<i>Glycothiohexides</i>	[20].
4	<i>Streptomyces</i>	<i>Streptothricin B, Albomycetin, Isomaltose, Grisamine, Tertiomycin, Thiomycin, Tertiomycin B, Streptogramin, Rifamycin, Lincomycin, Clindamycin, Platenomycin, Glycinothricin, Thienamycin, Siomycin, Neomycin, Carbomycin, Streptomycin, Azinothricin, Thioxamycin, Streptothricin, Tetrodecamycin, Lactonomycin, Nocathiacin</i>	[21].
5	<i>Streptoverticillium</i>	<i>Erythronolide B</i>	[22].

ANTIMICROBIAL ACTIVITY OF ACTINOMYCETES:

A kind of strategies are located for this cause and due to the fact that now no longer they all primarily based totally on identical principles, consequences received will additionally be profoundly stimulated now no longer most effective with the aid of using the techniques selected however

additionally with the aid of using the microorganisms the diploma of solubility of every test compound o be had screening strategies for detection of antimicrobial activity. Since guy began out stricken by different sicknesses as a result of infectious microorganisms, the hunt for his or her remedies began out and brought about the invention of a huge wide variety of

antibiotics from microorganisms inclusive of actinomycetes. Microorganisms have made an outstanding contribution to the fitness and wellness of humans during the world. However, speedy emergence of antimicrobial resistance amongst pathogenic microorganisms has brought about a renewed look for new antimicrobial sellers from Streptomyces. The maximum resistant micro-organism inflicting vital network acquired infections consist of methicillin resistant *Staphylococcus aureus*, vancomycin - resistant *Staphylococcus aureus*, vancomycin-resistant Enterococcus, prolonged spectrum β lactamase producing micro-organism such as *E. coli* and *Klebsiella spp* and a couple of drugs resistant *Mycobacterium tuberculosis*. Therefore, it's far vital to look for new, efficacious and secure antibiotics from herbal reasserts to fight the threat of drug-resistant infections [23].

CONCLUSION

Actinomycetes separated soil environment and in recent findings actinomycetes developed various bioactive compounds those are particularly important for antibiotics developed. These studies concluded that actinomycetes are high source of producing antimicrobial agents against pathogens.

ACKNOWLEDGEMENT

It is our privilege and honour to express our sincerest gratitude to the Parul University,

Vadodara, Gujarat for providing me all the necessary support and facilities including state of the art infrastructural facilities with advanced technological scientific laboratories and everything else that was required to carry out this work.

REFERENCE

- [1] Rahman, Md. Ajijur, et al. "Antibacterial Activities of Actinomycete Isolates Collected from Soils of Rajshahi, Bangladesh." Biotechnology Research International, Hindawi Limited, Sept. 2011, pp. 1–6.
- [2] Devi Salam, Menaka. "Antimicrobial Potential of Actinomycetes Isolated from Soil Samples of Punjab, India." Journal of Microbiology & Experimentation, no. 2, MedCrave Group, LLC, May 2014
- [3] Chaudhary, HotamS, et al. "Antibacterial Activity of Actinomycetes Isolated from Different Soil Samples of Sheopur (A City of Central India)." Journal of Advanced Pharmaceutical Technology & Research, no. 2, Medknow, 2013, p. 118. Crossref, doi:10.4103/2231-4040.111528
- [4] Kumar, Narendra, et al. "Isolation and Screening of Soil Actinomycetes as Source of Anti-biotics Active against Bacteria." International Journal of Microbiology Research, no. 2, Bioinfo Publications, Dec. 2010, pp. 12–16. doi: 10.9735/0975-5276.2.2.12-16.

-
- [5] Trenozhnikova, Lyudmila, and Azliyati Azizan. "Discovery of Actinomycetes from Extreme Environments with Potential to Produce Novel Antibiotics." *Central Asian Journal of Global Health*, no. 1, University Library System, University of Pittsburgh, Dec. 2018. doi:10.5195/cajgh.2018.337.
- [6] Mahajan, G. B., & Balachandran, L. (2012). Antibacterial agents from actinomycetes-a review. *Frontiers in bioscience (Elite edition)*, 4, 240-253.
- [7] Bailey, J. (Ed.). (2014). *The facts on file dictionary of botany*. Infobase Publishing.
- [8] Kushwaha, D. A. (2020). *Fungi (Viruses, Bacteria and Mycoplasma) (Vol. 1)*. Dr. AK Kushwaha.
- [9] Bronner, M., & Bronner, M. (2016). *Actinomycosis*. Butterworth-Heinemann.
- [10] Gao, R., Liu, C., Zhao, J., Jia, F., Yu, C., Yang, L., & Xiang, W. (2014). *Micromonospora jinlongensis* sp. nov., isolated from muddy soil in China and emended description of the genus *Micromonospora*. *Antonie van Leeuwenhoek*, 105(2), 307-315.
- [11] Mohr, K. I. (2016). History of antibiotics research. In *How to Overcome the Antibiotic Crisis* (pp. 237-272). Springer, Cham.
- [12] Nawani, Neelu, et al. "Actinomycetes: Role in Biotechnology and Medicine." *Bio Med Research International*, Hindawi Limited, 2013, pp. 1–1. doi:10.1155/2013/687190.
- [13] Mukhtar, Salma, et al. "Actinomycetes: A Source of Industrially Important Enzymes." *Journal of Proteomics & Bioinformatics*, no. 12, OMICS Publishing Group, 2017. Crossref, doi:10.4172/jpb.1000456.
- [14] Mehnaz, D., Abdulla, K., Aiysha, D., Zaheer, A., & Mukhtar, S. (2017). *Actinomycetes: a source of industrially important enzymes*.
- [15] Mast, Yvonne, and Evi Stegmann. "Actinomycetes: The Antibiotics Producers." *Antibiotics*, no. 3, MDPI AG, July 2019, p. 105. Crossref, doi:10.3390/antibiotics8030105
- [16] Nai, C., & Meyer, V. (2018). From axenic to mixed cultures: technological advances accelerating a paradigm shift in microbiology. *Trends in microbiology*, 26(6), 538-554.
- [17] Moon, D., & Landa, E. R. (2017). The centenary of the journal soil science: reflections on the discipline in the United States and Russia around a hundred years ago. *Soil Science*, 182(6), 203-215.
- [18] Mertz, F. P., & Yao, R. C. (1990). *Saccharopolyspora spinosa* sp. nov. isolated from soil collected in a sugar mill rum still. *International Journal of Systematic and Evolutionary Microbiology*, 40(1), 34-39.
-

-
- [19] Labeda, D. P., Testa, R. T., Lechevalier, M. P., & Lechevalier, H. A. (1984). *Saccharothrix*: a new genus of the Actinomycetales related to *Nocardiosis*. *International Journal of Systematic and Evolutionary Microbiology*, 34(4), 426-431.
- [20] BESIII collaboration, Ablikim, M., Achasov, M. N., Ai, X. C., Albayrak, O., Albrecht, M., & Li, L. (2015). Measurement of the $e^+e^- \rightarrow \pi^+\pi^-$ cross section between 600 and 900 MeV using initial state radiation. In *30 Years of BES Physics: Proceedings of the Symposium* (pp. 250-259).
- [21] Ohnishi, Y., Ishikawa, J., Hara, H., Suzuki, H., Ikenoya, M., Ikeda, H., & Horinouchi, S. (2008). Genome sequence of the streptomycin-producing microorganism *Streptomyces griseus* IFO 13350. *Journal of Bacteriology*, 190(11), 4050-4060.
- [22] Puranik, P. R., & Paknikar, K. M. (1997). Biosorption of lead and zinc from solutions using *Streptoverticillium cinnamomeum* waste biomass. *Journal of biotechnology*, 55(2), 113-124.
- [23] Nedialkova, D., & Naidenova, M. (2005). Screening the antimicrobial activity of actinomycetes strains isolated from Antarctica.