Evaluation and Management of Accidental Intra-Arterial Injection in the Ante-Cubital Fossa

Jayesh Patel¹, Pratiksha Shah^{2*}, Fenil Gandhi², Ali Guy³, Fareeha Nasir⁴, Edlira Horjeti⁵, Erisa Kola⁶, Loran Rakovica⁷ and Juna Musa⁸

¹Head of Department, Department of Vascular Surgery, Pramukhswami Medical College, Shree Krishna Hospital, Karamsad, Gujarat, India

²Intern, Department of Vascular Surgery, Pramukhswami Medical College, Shree Krishna Hospital, Karamsad, Gujarat, India

³Department of Physical Medicine and Rehabilitation, New York University, School of Medicine - NYU Medical Center, New York, USA

⁴Intern, Department of General Surgery, Christian Hospital Quetta, Pakistan

⁵Family Doctor, Department of Family Medicine, Tirane, Albania

⁶Department of Pathology and Forensic Medicine, Albania

⁷Medical Doctor, Faculty of Medicine, University of Prishtina, Prishtina, Kosovo

⁸Research Fellow, Department of Surgery, Mayo Clinic, Rochester, Minnesota, USA

***Corresponding Author:** Pratiksha Shah, Intern, Department of Vascular Surgery, Pramukhswami Medical College, Shree Krishna Hospital, Karamsad, Gujarat, India.

Received: April 01, 2020

Published: April 21, 2020

© All rights are reserved by Pratiksha Shah, et al.

Abstract

Early evaluation and management of patients presenting with clinical features of accidental intra-arterial injection is of uttermost importance due to its numerous complications. Also, it is important to have a standard treatment plan, which can be employed for limb salvation. In accordance with other modalities of treatment, the use of sympathetic ganglion block of 0.1% Ropivacaine 50 ml for 72 hours was very specific to all the cases that were reported. Following Ropivacaine use, patients reported pain relief with some movement of the digits, and an increase in the local temperature of the affected digits.

Keywords: Intra-Arterial Injection (IA); Diclofenac Sodium; Sympathetic Ganglion Block; Fasciotomy; Ray's Amputation

Introduction

Intra-arterial drug injection is a rare, but a potentially serious medical emergency. It is most commonly seen as an iatrogenic complication when administering an intravenous medication. Accidental intra-arterial injection was defined as an intravenous injection administered in the upper limb for any illness, which was followed by sudden severe pain in the limb followed by bluish discoloration of any part of the affected limb.

The aim of the study is to identify the risk factors early on for limb amputation following intra-arterial injection. Also, to assess the efficacy of the various modalities of the treatments administered and establish a standardized treatment plan for intra-arterial injection to achieve limb salvaging.

Accidental intra-arterial injection, whether being self-administered or iatrogenic, is a medical emergency. The normal vascular anatomy, aberrant vasculature, difficult emergency situations and human error, all contribute to the iatrogenic intra-arterial injections in an attempt of getting an intravenous access. Van der Post first reported the condition in 1942 [1]. It is estimated that accidental intra-arterial injection is revealed in 1:3500 - 1:56000 patients visiting the emergency department [2].

Barbiturates, thiopental sodium, narcotics, and tranquilizers are common drugs that have caused medical catastrophes affecting the upper limb [3]. Diclofenac sodium is also a common over the counter drug being used for analgesia. It is a non-steroidal

anti-inflammatory drug (NSAIDs) that is commonly used to treat mild to moderate pain, hyperpyrexia, and inflammation. It is most commonly administered through oral, intramuscular, intravenous, transdermal and rectal routes. It can lead to acute ischemia of the affected limb if administered intra-arterially because of its vaso-spastic action on the arterial wall. Common sites for error are the ante-cubital fossa, groin, and forearm due to the proximity of arteries and the veins at these sites [4].

The immediate reporting of the symptoms and accurate diagnosis are of central importance, as therapeutic strategies range from conservative methods to amputation of the affected limb.

The medical sequel that has been most commonly seen is paraesthesia, severe pain, motor dysfunction, compartment syndrome, gangrene, and limb loss [5]. Acute and/or chronic manifestations can be seen due to accidental IA injection. Several patients complained of immediate discomfort within seconds to a few minutes. The pain may range from local irritation to intense pain, presenting distal to the site of injection. Soon thereafter, many patients also complained of tingling sensations, burning sensations and paraesthesia. Altered motor functions such as involuntary muscle contractions and muscle weaknesses, and cutaneous manifestations such as flushing and mottling have been also commonly reported.

In addition, it has been proposed that there are patients that are at a higher risk of iatrogenic IA injection. Some examples including morbidly obese, dark pigmented patients, thoracic outlet

syndrome, patients with indwelling arterial catheters for blood pressure measurement, and pre-existing vascular anomalies of the forearm [5].

A symptomatic classification was established by Rutherford in 1986, revised in 1987 [6]. Rutherford classified peripheral arterial

disease in two parts, acute and chronic limb ischemia. The classification associated patient clinical symptoms with objective findings, including Doppler studies, arterial brachial index, and pulse volume recordings. It also emphasized that each presentation requires different treatment algorithms (Table 1).

Category	Findings			Doppler signal	
	Description/Prognosis	Sensory loss	Muscle weakness	Arterial	Venous
Viable	Not immediately threatened	None	None	Audible	Audible
Threatened					
Marginally	Salvageable if promptly treated	Minimal (toes) or none	None	Inaudible	Audible
Immediately	Salvageable with immediate revascularization	More than toes, associated with rest pain	Mild, Moderate	Inaudible	Audible
Irreversible	Major tissue loss or permanent nerve damage inevitable	Profound, anesthetic	Profound, paralysis	Inaudible	Inaudible

Table 1: Rutherford Classification for Acute Limb Ischemia.

The above classification was used to diagnose and treat a patient presenting with acute limb ischemia. It has helped the medical professionals to identify high-risk patients and treat accordingly.

Objective of the Study

The objective of the study is to identify the risk factors early on for limb amputation following intra-arterial injection. Also, to assess the efficacy of the various modalities of the treatments administered and establish a standardized treatment plan for intra-arterial injection to achieve limb salvaging. Since, intra-arterial injections are not a common incidence, it is difficult to follow and establish a standard plan of treatment to prevent an amputation and/ or permanent disability.

Materials and Methods

Study design

A case control study was conducted in the Vascular Surgery department of Shree Krishna Hospital from January 17th, 2017 to December 30th, 2018. A total of twelve patients with iatrogenic intra-arterial injection were studied. Accidental intra- arterial injection was defined as an intravenous injection administered in the upper limb for any illness, which was followed by sudden severe pain in the limb followed by bluish discoloration of any part of the limb. A thorough clinical history of all the patients was taken, enquiring about the various clinical features they may have experienced, for example pain, weakness, numbness, swelling surrounding the area, blackening of the affected region. Furthermore, the patients were examined for paraesthesia, pallor, edema, local temperature of the affected region, muscle contractures, and signs of gangrene. Also, the previous records were checked to determine the drug that had been accidentally administered intra-arterially. Simultaneously, all patients had been classified according to the Rutherford Classification, in order to have a better understanding of the severity of the injury. The vitals were constantly monitored from the time of presentation. Following clinical examination, all patients underwent an arterial Doppler study to determine the extent of the injury. According to the presenting clinical features, results from the Doppler

study, Rutherford classification, the treatment for each patient was determined upon. Moreover, all the patients underwent a thorough clinical examination and arterial Doppler studies post-operatively.

Inclusion criteria

- Patients presenting with signs and symptoms falling under the Rutherford Classification of Acute Limb Ischemia.
- Patients having a recent history of intravenous drug administration, possibly indicating accidental intra-arterial injection.
- Patients having positive Doppler findings indicating acute limb ischemia.

Exclusion criteria

- Patients presenting with sensory and motor function loss but no recent history of intravenous drug administration.
- Patients showing no positive findings on Doppler examination.

Ethics

A verbal consent was taken from each patient to use his/ her details for the study. The personal information of all the patients was kept confidential and in no manner manipulated. No harm was done to the patients. The patients details were solely used for this study only and no other research studies.

Results

Among the twelve patients, eleven were males and one was a female. The age of the patients ranged from 19 - 55 years. Ten out of the twelve patients had presented with complains of pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, and swelling of the arm. In addition, upper limb ischemia with blackening of 2nd and 3rd digits and hypothermia of the arm was also present. The patients presented to Trauma and Emergency Care (TEC) within 1 hour of taking an intravenous line where Diclofenac sodium was administered. Upon general examination, the vitals of all the patients were within normal limits. The local examination of the arm showed that brachial,

radial and ulnar arteries were not palpable on the affected side; axillary artery was palpable. Palpation of all the arteries in the opposite upper limb was plausible. The patients were classified as Category IIb (Immediately, Threatened), according to the Rutherford Classification (Table 1 and 2).

The following treatment was employed immediately in the patients:

- 1. Injection Heparin 10,000-unit IV stat followed by 1000 unit/ hour infusion. The target APTT being 75 - 85 sec-

Patient	Age (Years)	Sex	H/O IV line taken	Drug Administered	Time from when the first symptoms appeared to when they presented to TEC
1)	19	Male	Yes	Diclofenac Sodium	30 minutes
2)	35	Male	Yes	Diclofenac Sodium	45 minutes
3)	35	Male	Yes	Diclofenac Sodium	30 minutes
4)	36	Female	Yes	Diclofenac Sodium	60 minutes
5)	38	Male	Yes	Diclofenac Sodium	1.5 hours
6)	41	Male	Yes	Diclofenac Sodium	60 minutes
7)	45	Male	Yes	Unknown	60 minutes
8)	50	Male	Yes	Diclofenac Sodium	30 minutes
9)	51	Male	Yes	Diclofenac Sodium	45 minutes
10)	55	Male	Yes	Diclofenac Sodium	30 minutes
11)	36	Male	Yes	Unknown	After 12 hours
12)	38	Male	Yes	Unknown	After 12 hours

Table 2: General details of the patients selected for this study.

Patient	Site of IV Line	Clinical Features	Rutherford Classification
1)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
2)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, swelling over the forearm 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
3)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
4)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, hypothermia present 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
5)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, swelling over the forearm 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
6)	Left Upper Limb in the Ante-cubital Fossa	1. Pain in the left forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint 2. Brachial, Radial and Ulnar arteries were not palpable on the left side; Axillary artery was palpable. Palpation of all arteries on the right side was plausible.	Category IIb = Immediately, Threatened
7)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, swelling over the forearm 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened

8)	Left Upper Limb in the Ante-cubital Fossa	1. Pain in the left forearm and hand, numbness and tingling sensations, swelling over the forearm 2. Brachial, Radial and Ulnar arteries were not palpable on the left side; Axillary artery was palpable. Palpation of all arteries on the right side was plausible.	Category IIb = Immediately, Threatened
9)	Right Upper Limb in the Ante-cubital Fossa	1. Pain in the right forearm and hand, numbness and tingling sensations, inability to move finger and wrist joint, swelling over the forearm, hypothermia present 2. Brachial, Radial and Ulnar arteries were not palpable on the right side; Axillary artery was palpable. Palpation of all arteries on the left side was plausible.	Category IIb = Immediately, Threatened
10)	Left Upper Limb in the Ante-cubital Fossa	1. Pain in the left forearm and hand, numbness and tingling sensations, swelling over the forearm, hypothermia present 2. Brachial, Radial and Ulnar arteries were not palpable on the left side; Axillary artery was palpable. Palpation of all arteries on the right side was plausible.	Category IIb = Immediately, Threatened
11)	Left Upper Limb in the Ante-cubital Fossa	Blackening of left 1 st , 2 nd and 3 rd fingers	Category III = Irreversible
12)	Right Upper Limb in the Ante-cubital Fossa	Blackening of right 1 st , 2 nd and 3 rd fingers	Category III = Irreversible

Table 3: Clinical features and Rutherford Classification.

- onds. Monitoring was done every 6 hours for 5 days. This was done to prevent thrombosis. A significant improvement was noticed, due to which the use of thrombolytic therapy was not required.
2. Injection Hydrocortisone 200 mg IV TDS for 3 days, followed by 100 mg TDS for 2 days and 100 mg OD for 2 days. It aided in enhancing tissue repair.
 3. Injection Sulfamethoxazole-Trimethoprim 40 ml/hour for 48 hours.
 4. Sympathetic ganglion block of 0.1% Ropivacaine 50ml for 72 hours. Patients reported pain relief with some movement of the digits. Also the local temperature of the digits began to increase.
 5. Injection lidocaine (lignocaine) hydrochloride 2% IV infusion for 5 days. It helped in vasodilation, hence decreasing vasospasm.
 6. Tablet Aspirin 75 mg once a day qPM.
 7. Intravenous Analgesics.
 8. Intravenous Antibiotics.
 9. Upper limb fasciotomy under general anesthesia.
 10. Limb Elevation to favor limb drainage and prevent edema from occurring.
 11. Physiotherapy.
 12. Doppler ultrasonography was done every 24 hours.

An upper limb Arterial Doppler was performed in all the patients. It showed echogenic areas in the distal most part of the affected brachial artery, signs of thrombosis within, and lack of flow in the radial and ulnar artery beyond. The patients were advised

to undergo upper limb fasciotomy under general anesthesia. Upon discharge all vitals of the patients were normal, and limb was successfully salvaged. The patients were advised to keep the limb elevated and physiotherapy was prescribed.

The following drugs were prescribed upon discharge:

1. Oral Antibiotics.
2. Oral Analgesics.
3. Tablet Aspirin 75 mg once a day qPM for 3 months.
4. Doppler USG was performed twice during the follow up to ensure proper functioning of brachial, radial and ulnar arteries.

The patients were asked to come for a follow up examination every week for two weeks and biweekly afterwards. Upon completion of three months, the wound was completely healed and healthy. In addition, the patients reported of a well-functioning limb.

The remaining two patients had presented with complain of blackening of 1st, 2nd and 3rd digits. It was gradual in onset and progressive in nature. It was also associated with pain. They also reported a history of a previous injury when an IV line was taken to administer medications. No significant past history was present. Upon general examination, all the vitals of the patients were normal. Local examination showed blackening of the 1st, 2nd and 3rd digits. The patients were classified as Category III (Irreversible), under the Rutherford Classification (Table 1).

The following medications were administered immediately as they presented to the Emergency Department:

1. Tablet Aspirin 75 mg once a day in qPM
2. Intravenous Antibiotics
3. Intravenous Analgesics

4. Intravenous Fluids

5. Ray’s amputation was performed under sedation and block. Bone nibbled and metacarpophalangeal joints and digits amputated. Sterile dressing was done with limb elevation.

6. Limb Elevation

7. Physiotherapy.

ensure that the wound was healthy and healing well. No other complaints were reported of. The following drugs were prescribed upon discharge:

1. Oral Antibiotics

2. Oral Analgesics

3. Physiotherapy

4. Limb Elevation
- Upon discharge the patients were advised a high protein and ketone diet. They were asked to come for a weekly follow up to

Patient	Doppler USG study	Treatment	Outcome
1)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
2)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
3)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
4)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
5)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
6)	Echogenic areas in the distal most part of left brachial artery, signs of thrombosis within, and lack of flow in the left radial and ulnar artery beyond.	Left Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
7)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
8)	Echogenic areas in the distal most part of left brachial artery, signs of thrombosis within, and lack of flow in the left radial and ulnar artery beyond.	Left Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
9)	Echogenic areas in the distal most part of right brachial artery, signs of thrombosis within, and lack of flow in the right radial and ulnar artery beyond.	Right Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
10)	Echogenic areas in the distal most part of left brachial artery, signs of thrombosis within, and lack of flow in the left radial and ulnar artery beyond.	Left Upper Limb Fasciotomy Limb Elevation Physiotherapy	Healthy wound upon follow up Functions of the limb were restored Limb was salvaged
11)	Amputation Done, Doppler Findings - Not applicable	Ray’s Amputation of left 1 st , 2 nd and 3 rd fingers Limb Elevation Physiotherapy	Amputation stump was healthy upon follow up 1 st , 2 nd , and 3 rd fingers not salvaged Left Upper Limb functioning well otherwise

12)	Amputation Done, Doppler Findings - Not applicable	Ray's amputation of right 1 st , 2 nd and 3 rd fingers Limb Elevation Physiotherapy	Amputation stump was healthy upon follow up 1 st , 2 nd , and 3 rd fingers not salvaged Right Upper Limb functioning well otherwise
-----	--	--	--

Table 4: Investigations and treatment.

Discussion

Intra-arterial injection is an infrequent event, leading to a serious medical emergency. It may be iatrogenic and/or in relation to drug abuse. The most common site for intra-arterial injection is the ante-cubital fossa where the branches of brachial and ulnar arteries are superficial and can be easily entered. Some of the proposed mechanisms that explain the emergency are vasospasm, intra-vascular thrombosis and chemical endarteritis [7]. The following events lead to acute thrombosis of the affected artery, followed by a decrease in blood supply to the digits. This will eventually lead to the development of gangrene of the affected limb. Also, it has been previously proposed that the benzyl alcohol preservative used in the non-aqueous preparation of diclofenac could be the cause of the emergency vasospasm due to endothelial edema and capillary endothelial dysfunction.

Conservative treatment comprised of vasodilators, steroids, nerve blocks, analgesics, and prophylactic antibiotics. Lidocaine hydrochloride is a local anesthetic agent, which is a sterile, non-pyrogenic solution. It was one of the drugs administered to the patients who presented early and with no signs of gangrene, where it leads to vasodilatation of the artery. Low molecular weight heparin was administered to prevent further thrombosis from occurring. Monitoring of coagulation parameters was also performed to anticipate and prevent serious bleeding complications. Hydrocortisone is a steroid, which prevents the release of substances that lead to inflammation. When administered in patients, it aided in enhancing tissue repair. The utmost important modality, which proved to be of significance, was the sympathetic ganglion block. Patients reported of a decrease in pain and an increase in local temperature. The sympathetic nerve block targets the sympathetic nervous system, which contains a series of nerves that spread out from the spine to the body, helping to control several involuntary bodily functions. Furthermore, rest, physiotherapy, and elevation to the affected limb are of importance as it leads to a decrease in edema and speeds the recovery process.

Treatment for IA injection also comprises of surgical modalities, which are employed in almost all the patients. Patients that presented early, with no clinical features of gangrene, can be easily operated upon to salvage the limb. Fasciotomy was the procedure performed, where the tissue surrounding the area was cut open to relieve pressure. It is a surgery to relieve swelling and pressure from any compartment of the body. This helps to increase blood flow, and to stop/ slow down the rate of damage to the local tissue. Some complications that may appear are excessive bleeding, scarring, infection, and chronic pain. Moreover, patients that presented late and with clinical features of gangrene were immediately treat-

ed with Ray's amputation of the affected digits. The gangrene was halted at the very minimum severity as possible, hence preserving some function of the affected digits. In a different article, Samantha and Samantha reported cases of accidental intra-arterial injection where the patient presented with necrosis of the finger for which amputation was performed [8]. Diclofenac sodium was injected in the radial artery, which was mistaken for a vein. The rate of amputation for accidental intra-arterial drug injection in such cases is 29% [9]. In addition, it is of utmost importance to follow up with the patients to ensure proper functioning of the limb and to discover any occurrence of further complications.

Complications occurring due to iatrogenic intra-arterial injection have an incidence between 1:3500 to 1:56000 as reported by several authors [10]. Some of the reported complications being acute limb ischemia, unbearable pain at the site of injection, numbness, tingling sensations, inability to move the limb, and lastly the onset of gangrene. All the patients that were studied reported of two or more of the above mentioned complications. This study has helped to identify a standardized treatment plan that has worked in salvaging the limb. Although, the treatment plan can always be modified according to the patients needs and their clinical features.

Conclusion

Clinically, it may not be easy to prevent cases of accidental intra-arterial injection from occurring, however incidence can be lowered with supervision and proper training of the medical personnel. It is important that a medically trained physician supervise this task, in order to prevent such catastrophes from happening. Furthermore, it is important to identify the signs early on, and administer appropriate treatment to reduce the possibility of amputation and prevent the patient from having a permanent disability. Lastly, a clinician must not forget to educate and inform the patient of the possible outcome of his/her condition. The patients must be taken care of with the utmost care.

Bibliography

1. Van der Post CWH. "A case of mistaken injection of pentothal sodium into an aberrant ulnar artery". *South African Medical Journal* 16 (1942): 182-184.
2. Malik A., et al. "Accidental Intra Arterial Injection And Limb Ischemia". *Journal of Ayub Medical College, Abbottabad* 29.2 (2017): 230-233.
3. Kumar M., et al. "Accidental Intra-Arterial Injection of Diclofenac-Case Report". *Journal of Clinical and Diagnostic Research* 9.1 (2015): PD16-PD17.

4. Caroline Lake and Christina L Beecroft. "Extravasation Injuries and Accidental Intra-Arterial Injection". *Continuing Education in Anaesthesia Critical Care and Pain* (2010).

5. Surjya Sen., *et al.* "Complications After Unintentional Intra-Arterial Injection of Drugs: Risks, Outcomes, and Management Strategies". *Mayo Clinic Proceedings* 80.6 (2005): 783-795.

6. Cohen SM. "Accidental Intra-Arterial Injection of Drugs". *Lancet* 2.6523 (1948): 361.

7. Ghouri AF, *et al.* "Accidental Intraarterial Drug Injections via Intravascular Catheters Placed on the Dorsum of the Hand". *Anesthesia and Analgesia* 95.2 (2002): 487-491.

8. Sukhen Samanta and Sujay Samanta. "Accidental intra-arterial injection of diclofenac sodium and their consequences: report of two cases". *Anaesthesia Pain and Intensive Care* 17.1 (2013): 101-102.

9. Devulapalli C., *et al.* "Inadvertent Intra-Arterial Drug Injections in the Upper Extremity: Systematic Review". *Journal of Hand Surgery* 40.11 (2015): 2262-2268.e5.

10. Stone HH and Donnelly CC. "The Accidental Intraarterial Injection of Thiopental". *Anesthesiology* 22 (1961): 995-1006.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: <https://www.actascientific.com/>

Submit Article: <https://www.actascientific.com/submission.php>

Email us: editor@actascientific.com

Contact us: +91 9182824667