Bioactive *Psiadia punctulata* Trachylobane Diterpenes Against Mastitis PathogensWanjau OM^{1*}, Mahungu SM² and Matasyoh JC¹¹Department of Chemistry, Faculty of Science, Egerton University, Kenya²Department of Dairy and Food Science, Faculty of Agriculture, Egerton University, Nakuru, Kenya***Corresponding Author:** Wanjau OM, Department of Chemistry, Faculty of Science, Egerton University, Kenya.

DOI: 10.31080/ASNH.2020.04.0653

Received: February 21, 2020**Published:** February 26, 2020© All rights are reserved by **Wanjau OM, et al.****Abstract**

Bacterial pathogens are a major challenge and a constant drawback in the management of both human and animal health. The dairy farmer may suffer huge losses due to infection of the udder of dairy cows with mastitis. Manifestation of mastitis in dairy cows occurs, resulting from poor personnel hygiene, unsuitable environment, unhygienic milking procedure or use of contaminated milking equipment and supplies. At times managerial measures have failed to give significant success, and use of post milking teat dipping solutions has been applied with success.

The presence of a leaf surface exudate on *Psiadia punctulata* is supposedly for a purpose, and this, coupled with the traditional uses for the plant warrant bioassay directed phytochemical studies on the plant.

Column chromatography of concentrated cold acetone leaf wash of *Psiadia punctulata* using silica gel; serially eluting the column with ethyl acetate-hexane mixtures, yielded two trachylobane diterpenoid compounds, PP5 and PP6 as clear crystalline compounds. *In vitro* bioassay tests applying the disc diffusion method and the agar diffusion methods using levofloxacin and chloramphenicol as positive controls found the trachylobane diterpenoid compounds to be active against *Staph aureus* and *Salmonella typhi*. Structural characteristics of the diterpenoid compounds were evaluated through phytochemical tests and correlation of proton and carbon-13 NMR spectra. The presence of antimicrobial agents in acetone leaf wash of *Psiadia punctulata*, active against *Staph aureus* and *Salmonella typhi* makes the plant invaluable as a source of a home remedy in eradication of mastitis manifestation in dairy cows.

Keywords: *Psiadia punctulata*; Antimicrobial; Trachylobane Diterpenes; Mastitis, *Staph aureus* and *Salmonella typhi*

Introduction

Mastitis, also known as udder inflammation, is a common problem in dairy herds and causes huge economic losses in the dairy industry [1]. *Staphylococcus aureus*, *Escherichia coli* and *Streptococcus uberis* are the main mastitis-causing pathogens [2]. More antimicrobial agents must be identified to replace some of the antimicrobial drugs that have developed drug resistance due to extensive use, misuse and overuse of antimicrobials in both human and animal health. *Staphylococcus aureus* is also one of the most frequent causal agents of lower respiratory tract infections. Methicillin-resistant *S. aureus* (MRSA) is of particular concern in the United States since approximately 60% of all staphylococcal infections in the ICU are caused by MRSA [3].

Plants are a reservoir of antimicrobial substances which are not only potent against target pathogens but also seem to stand a better chance to overcome the onslaught of microbial resistance mechanisms [4].

The medicinal value of plants lies in the ability of their constituent compounds to produce physiological action in the human body [5]. The most important of these bioactive compounds include alkaloids, terpenoids, steroids, flavonoids, tannins and phenolic compounds [6]. Leaf decoctions of *Psiadia punctulata* have

been used as a remedy for colds, headache and abdominal pains [7]. Maasai use the leaf decoction to get rid of fleas from calves [8] (Henke, 1994) while a report by Maurice, et al. (1999) [9] details on the use of the leaf decoction of *Psiadia punctulata* for the treatment of venereal diseases and as a tonic.

The genus *Psiadia* belongs to the family *Compositae* (*Asteraceae*). It is a shrub that is draught resistant and is widely spread in the arid regions of the Rift valley, Eastern region and the drier parts of central Kenya. Its leaves have a shiny look and develop a sticky feel soon after they have been plucked due to the presence of some leaf exudate which is found as 24% dry leaf weight [10]. The constituent compounds of leaf surface exudate [11] supposedly serve some protective role on the plant [12].

Meena Shamrao Deogade, et al. (2016) [13] reported antimicrobial activity of a methanol extract of fresh leaves of *Ocimum tenuiflorum* Linn. against *Escherichia coli*, *Staphylococcus aureus* and *Klebsiella pneumonia* at 20 mg/ml, corresponding to inhibition zones of 8 mm, 16 mm and 11 mm respectively. Shruthi, et al. (2012), [14] working on *Murraya Koenigii* observed antibacterial activity in ethanol extract at 12.43 mg/ml for *Klebsiella pneumonia* and 9.23 mg/ml for *Salmonella typhimurium*. Acetone leaf extracts of nine plant species (Elisha, et al. 2017) [15] were active against

Escherichia coli, *Salmonella typhimurium* and *Pseudomonas aeruginosa* at MIC levels ranging between 0.04 and 0.32 mg/ml. Methanol root extract *Clerodendrum myricoides* (Hochst.) R. Br. ex Vatke, locally known as “Munjuga-iria” in Mbeere community of Embu County in Kenya was active against methicillin resistant *S. aureus* (MRSA) at MIC level of 15.3 µg/ml [16].

Plant materials

The resinous leafy branches of *P. punctulata* were harvested in the neighborhood of lake Elementaita, about 35 kilometers east of Nakuru town, Kenya, at an altitude of 1,670 m. A voucher specimen is available at the National Museum of Kenya Herbarium.

Extraction

Leaves were detached from the branches of *P. punctulata* plant and were stuffed into a 5L erlmeyer flask while fresh, up to a weight of 1 Kg. for extraction. A 1.5 liter portion of cold acetone was introduced into the flask and shaken for two minutes and decanted as an orange solution [4,17]. Two more fresh portions of acetone were used to wash the leaves clear of the orange exudate. Extracted leaves were allowed to dry in open air to a constant weight of 438g. The acetone extract was decanted and filtered through some glass wool into a 2.5L Winchester bottle. The acetone was recovered using a rotary evaporator and the concentrate was preserved in an open brown glass bottle (100 ml) and stored in a vacuum desiccator until a constant weight of 47.5g of concentrate (10% wt. of dry extracted leaves) was attained.

Chromatographic methods

10 g of the extract was introduced into a glass column (4.5 cm diameter), packed with 230 - 400 mesh silica gel up-to a height of 27 cm. The column was sequentially eluted with 500 ml portions of 15%, 20%, 40%, 50% and 100% ethylacetate in hexane [18]. The eluant was collected as 40 ml fractions from which pure compounds were isolated through fractional crystallization from acetone-hexane mixtures. Anisaldehyde spray solution was used as the locating reagent on TLC plates developed in 8:2 ethylacetate-hexane solvent mixture. Melting points (uncorrected) were determined using a Gallen camp melting point apparatus. 1H and 13C NMR spectra were determined at 500 and 125 MHZ, respectively.

Identification of the compounds was achieved through correlation of spectral data, melting point values and reference to literature data [19].

Antimicrobial disc diffusion tests

Tests for antibacterial activity for pure compounds and the crude plant extract were carried out using the agar disk diffusion assay [20]. Agar medium was prepared from commercially available dehydrated base according to the manufacturer’s instruction. The media was reconstituted at 120°C for 15 - 20 minutes in an autoclave, after which it was immediately allowed to cool in a water bath maintained at 45 - 50°C. The freshly prepared and cooled medium was poured into disposable plastic flat-bottomed petri dishes to give a uniform depth of approximately 5 mm. The agar

medium was allowed to cool at room temperature and stored at 2 - 8°C. A representative sample of each batch was examined for sterility by incubating at 37°C for 24 hours.

Preparation of discs

Discs for the drug sensitivity test were prepared by punching discs of 6mm diameter using Whatman filter paper No.1. The discs were dispensed into a screw capped glass bottle and sterilized by dry heat in a hot air oven at 180°C for 2 hours.

Preparation of antimicrobial test solutions

Levofloxacin (500 mg) was used as positive control at concentration levels of 10 ppm, 5 ppm and. 2.5 ppm. One tablet (500 mg) was dissolved in cold DMSO: H₂O mixture in the ratio of 1:5 to make a 1000 ml stock solution of 500 ppm. 2 ml of the 500 ppm stock solution were diluted to 100 ml to give 10 ppm solution. 10 ml and 5 ml of the 10 ppm solution were consecutively diluted to 20 ml in separate volumetric flasks to give 5 ppm and 2.5 ppm solutions respectively.

Test solutions of PP5 and PP6 were prepared by dissolving 10 mg of each into 50 ml of 1:5 DMSO:H₂O mixture to make stock solutions of 200 ppm. Test solutions of 10 ppm, 5 ppm and 2.5 ppm were prepared through serial dilution of the 200 ppm stock solutions.

Discs treated with the test solutions were placed individually on the inoculated agar plates using the fine sterile forceps. The discs were distributed evenly so that they were no closer than 25 mm from center to center. Three discs were placed on a plate of 100mm in diameter. Within 15 minutes after the discs were applied, the plates were inverted and placed in an incubator set at 37°C. Results were noted and recorded after an incubation period of 24 hours.

Antimicrobial agar dilution tests

Varying concentrations of the antimicrobial agents were incorporated in series into agar plates by mixing 2 ml of each dilution with 18 ml of molten agar per plate, at 50°C. Four holes of 6 mm in diameter each were perpendicularly dug at evenly spaced positions into the agar mixture after setting. Each well was inoculated with the same volume (0.1 ml) of the standardized suspension of the bacteria under test - *Salmonella typhi* (1), *Staphylococcus aureus* (2), *Bacillus subtilis* (3) and *Proteus mirabilis* (4). The agar plates were incubated for 24 hours, after which MIC for each drug was determined as the lowest concentration that inhibited bacteria growth.

Results and Discussion

Column chromatography of the acetone leaf extract using hexane-ethylacetate solvent system mixtures yielded antibacterial trachylobane diterpenoid compounds PP5 and PP6 whose structural data correlate to compounds 1, 2 and 3 in (Figure 1) [21].

Doublets at δ 0.6 and δ 0.8 were characteristic of cyclopropane ring H-16 and H-12 with corresponding carbon resonances at δ 25.35 (C-16) and δ 23.53 (C-12).

Figure 5: Agar dilution test.

Antimicrobial activity has also been reported for both chloroform and methanol extracts of *Larrea tridentata* against clinical isolates of *Staphylococcus aureus* (n = 25) at MIC₈₀ = 125 µg/ml for both extracts [23].

Conclusion

The present study indicate PP5 and PP6 as effective antimicrobial agents against *Staph aureus* and *Salmonella typhi* at concentration levels below 2.5 ppm. Clinical trials are required for possible application of the trachyloban diterpenes in the formulation of hand washing soaps and milking salves in the management of mastitis manifestations in dairy cows.

Acknowledgements

The authors are grateful to the University of Kwa-Zulu Natal and Prof Dulcie Mullholand for support in laboratory space, chemicals and instruments.

Bibliography

1. Aleksandra Kalinska., et al. "Mastitis pathogens in dairy cattle – a review". *World Scientific News* 89 (2017): 22-31.

2. Misrak Birhanu., et al. "Prevalence of bovine subclinical mastitis and isolation of its major causes in Bishoftu Town, Ethiopia". *BMC Research Notes* 767 (2017).

3. Virgilio Bocanegra-García., et al. "The bioactivity of plant extracts against representative bacterial pathogens of the lower respiratory tract". *BMC Research Notes* 2 (2009): 95.

4. Kenneth G Ngwoke., et al. "Antimicrobial natural. Products". In book: *Science against microbial pathogens: communicating current research and technological Advances*. Published by Formatex Spain (2011).

5. Sebastien Block., et al. "Diterpenes from the leaves of *Croton zambesicus*". *Phytochemistry* 65 (2004): 1165-1171.

6. T Mythili and R Ravindhran. "Phytochemical screening and antimicrobial activity of *Sesbania sesban* (L.)". *Asian Journal of Pharmaceutical and Clinical Research* 5.4 (2012): 179-182.

7. J Ogwenso Midiwo., et al. "Bioactive compounds from some Kenyan ethnomedicinal plants: Myrsinaceae, Polygonaceae and *Psiadia punctulata*". *Phytochemistry Reviews* 1 (2002): 311-323.

8. Henke (1994).

9. Maurice., et al. (1999).

10. J Ogwenso Midiwo., et al. "Flavonoids of *Polygonum senegalense* part III: Isolation of dihydrochalcone glucoside and quercetin glucosides". *Bulletin of the Chemical Society of Ethiopia* 8.2 (1994): 79-84.

11. Midiwo JO., et al. "Diterpenes from the leaf exudates of *Psiadia punctulata*". *Phytochemistry* 45 (1997): 117-120.

12. Uma BF., et al. "Flavones and phenylpropanoids in the surface exudates of *Psiadia punctulata*". *Phytochemistry* 57 (2001): 571-574.

13. Meena Shamrao Deogade., et al. Antibacterial.

14. Shruthi., et al. (2012)

15. Ishaku Leo Elisha., et al. The antibacterial.

16. Sospeter Ngoci Njeru., et al. "Antimicrobial and cytotoxicity properties of the organic solvent fractions of *Clerodendrum myricoides* (Hochst.) R. Br. ex Vatke: Kenyan traditional medicinal plant". *Journal of Intercultural Ethnopharmacology* (2016).

17. Wanjau OM., et al. "Antioxidant Activity of Methoxylated Flavonoids in Oils in Deep Frying Processes". *Food and Nutrition Sciences* 9 (2018): 1273-1284.

18. Sergio R Ambrosio., et al. "Antibacterial Activity of Kaurane Diterpenes". *Zeitschrift für Naturforschung* 63 (2008): 326D330.

19. World Health Organization. Antimicrobial drug resistance report by Secretariat. Executive Board EB134/37, 134th Session 6 December (2013).

20. J Ochei and A Kolhatkar. Antimicrobial susceptibility tests in Medical Laboratory Science Theory and Practice (2007): 789-810.

21. JO Midiwo., *et al.* (1997b)

22. Ramzi AA., *et al.* "Assessment of selected Yemeni Medicinal plants for their in vitro antimicrobial, anticancer and antioxidant activities". *Pharmaceutical Biology* 49.2 (2011): 200-210.

23. Virgilio Bocanegra-García., *et al.* (2009b).

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: <https://www.actascientific.com/>

Submit Article: <https://www.actascientific.com/submission.php>

Email us: editor@actascientific.com

Contact us: +91 9182824667