A Unique Technique to Treat Comedones with the Combination of Platelet Rich Plasma with Microneedling on Face

Niyati Dhawan and Brajpal Singh Tyagi*

Head Department of ENT Columbia Asia Hospital Ghaziabad, CEO Harsh ENT Hospital, Centre for Deafness, ABR Health Care Solutions Limited, India

***Corresponding Author:** Brajpal Singh Tyagi, Professor, Head Department of ENT Columbia Asia Hospital Ghaziabad, CEO Harsh ENT Hospital, Centre for Deafness, ABR Health Care Solutions Limited, India.

Received: May 07, 2020

Published: June 01, 2020

© All rights are reserved by Niyati Dhawan and Brajpal Singh Tyagi.

Abstract

Comedo is Latin word for Glutton. It is collectively known as Acne, white heads, black heads, pustules, pimples and infected cysts. We are presenting the two cases of comedones successfully treated with PRP and microneedling at our center.

Keywords: Atrophic; Concentrate; Comedones Management; Platelet; Platelet-Rich Plasma (PRP); Scar

Objective of the Study

Aim is to study the effect of PRP and Micro Needling in treatment of comedones.

Mechanism of action

PRP: The PRP's mechanism of action involves the release of powerful antimicrobial peptides from the platelets' alpha granules.

PRP is a promising adjunct in scar management practice. Further research with long-term follow-up is warranted to delineate the value of this modality in different subtypes of scars.

Microneedling: Collagen induction therapy. It enhance the PRP delivery in transdermal region. These micro injuries lead to minimal superficial bleeding and set up a wound healing cascade with release of various growth factors such as platelet derived growth factor (PGF), transforming growth factor alpha and beta (TGF- α and TGF- β), connective tissue activating protein, connective tissue growth factor, and fibroblast growth factor (FGF) [1]. The needles also breakdown the old hardened scar strands and allow it to revascularize. Neovascularization and neocollagenesis is initiated by migration and proliferation of fibroblasts and laying down of intercellular matrix [2,3]. A fibronectin matrix forms after 5 days of injury that determines the deposition of collagen resulting in skin tightening persisting for 5 - 7 years in the form of collagen III. The depth of neocollagenesis has been found to be 5 - 600 μ m with a 1.5 mm length needle. Histological examination of the skin treated with 4 microneedling sessions 1 month apart shows upto 400% increase in collagen and elastin deposition at 6 months postoperatively, with a thickened stratum spinosum and normal rete ridges at 1 year postoperatively [4]. Collagen fibre bundles ap-

pear to have a normal lattice pattern rather than parallel bundles as in scar tissue [5].

Case-I

Figure 1

Case-II

Figure 2

Figure 2

Conclusion

Microneedling is an effective modality of treatment, especially in patients with Fitzpatrick’s IV and V skin types because it overcomes the side effects of scarring and hyperpigmentation resulting from other procedures in which the epidermis is compromised. It certainly promises to be a valuable technique with its numerous applications and its ever-expanding modifications as well as feasibility of home use.

Bibliography

1. Falabella AF and Falanga V. "Wound healing. The Biology of the Skin". Parethenon: New York (2001): 281-299.
2. Fabbrocini G., et al. "Acne scarring treatment using skin needling". *Clinical and Experimental Dermatology* 34.8 (2009): 874-879.
3. Majid I., et al. "Microneedling and its applications in dermatology". In Prime. 7. Volume 4. London: Informa Healthcare (2014): 44-49.
4. Aust MC., et al. "Percutaneous collagen induction therapy: An alternative treatment for scars, wrinkles, and skin laxity". *Plastic and Reconstructive Surgery* 121.4 (2008): 1421-1429.
5. Nair PA and Arora TH. "Microneedling using dermaroller: A means of collagen induction therapy". *Gujarat Medical Journal* 69.1 (2014): 24-27.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
 - Thorough Double blinded peer review
 - Rapid Publication
 - Issue of Publication Certificate
 - High visibility of your Published work
- Website: <https://www.actascientific.com/>
Submit Article: <https://www.actascientific.com/submission.php>
Email us: editor@actascientific.com
Contact us: +91 9182824667

1. Daniel Y. "Functional and aesthetic effects of spreader grafts technique in rhinoplasty". *Scripta Scientifica Medica* 46.1 (2014): 68-73.

2. Samaha M and RassouliA. "Spreader graft placement in endo-nasal rhinoplasty: Technique and a review of 100 cases". *Plastic Surgery* 23.4 (2015): 252-254.

3. Sheen JH. "Spreader graft: a method of reconstructing the roof of the middle nasal vault following rhinoplasty". *Plastic and Reconstructive Surgery* 73.2 (1984): 230-239.

4. Rohrich RJ and Ghavami A. "Rhinoplasty for middle eastern noses". *Plastic and Reconstructive Surgery* 123 (2009): 1343-1354.

5. Hassanpour SE., et al. "Comparison of Aesthetic and Functional Outcomes of Spreader Graft and Autospreader Flap in Rhinoplasty". *World Journal of Plastic Surgery* 5.2 (2016): 133-138.

6. Zojaji R., et al. "Quality of life in patients who underwent rhinoplasty". *Facial Plastic Surgery* 30 (2014): 593-596.

7. Bessler S., et al. "Anterior spreader flap technique: a new minimally invasive method for stabilizing and widening the nasal valve". *Journal of Plastic, Reconstructive and Aesthetic Surgery* 68.12 (2015): 1687-1693.

8. Gruber RP, et al. "The spreader flap in primary rhinoplasty". *Plastic and Reconstructive Surgery* 119.6 (2007): 1903-1910.

9. Byrd HS, et al. "Using the autospreader flap in primary rhinoplasty". *Plastic and Reconstructive Surgery* 119 (2007): 1897-1902.

10. Oneal RM and Berkowitz RL. "Upper lateral cartilage spreader flaps in rhinoplasty". *Aesthetic Surgery Journal* 18 (1998): 370-371.

11. FomonS and Caron A. "Collapsedala". *Acta Oto-Laryngologica*, 51 (1950): 465-484.

12. Yoo S and Most SP. "Nasal airway preservation using the autospreader technique. Analysis of outcomes using a disease-specific quality of life instruments". *Archives of Facial Plastic Surgery* 13.4 (2011): 231-233.

13. Lipan MJ and Most SP. "Development of a severity classification system for subjective nasal obstruction". *JAMA Facial Plastic Surgery* 15.5 (2013): 358-361.

14. Stewart MG., et al. "Development and validation of the Nasal Obstruction Symptom Evaluation (NOSE) scale". *Otolaryngology-Head and Neck Surgery* 130.2 (2004): 157-163.

15. Lam DJ., et al. "Comparison of anatomic, physiological, and subjective measures of the nasal airway". *The American Journal of Rhinology* 20.5 (2006): 463-470.

16. Stewart MG., et al. "Outcomes after nasal septoplasty: results from the Nasal Obstruction Septoplasty Effectiveness (NOSE) study". *Otolaryngology-Head and Neck Surgery* 130.3 (2004): 283-290.

17. Most SP. "Analysis of outcomes after functional rhinoplasty using a disease-specific quality-of-life instrument". *Archives of Facial Plastic Surgery* 8.5 (2006): 306-309.

18. Lerma J. "The "lapel" technique". *Plastic and Reconstructive Surgery* 102.6 (1998): 2274-2275.

19. Murrell GL. "Auricular cartilage grafts and nasal surgery". *Laryngoscope* 114.12 (2004): 2092-2102.

20. Jaykaran C and Tamoghna B. "How to Calculate Sample Size for Different Study Designs in Medical Research?". *Indian Journal of Psychological Medicine* 35.2 (2013): 121-126.

21. Standlee AG and Hohman MH. "Evaluating the Effect of Spreader Grafting on Nasal Obstruction Using the NOSE Scale". *Annals of Otolology, Rhinology, and Laryngology* 126.3 (2017): 219-223.

22. Hussein WK., et al. "Modified autospreader flap for nasal valve support: utilizing the spring effect of the upper lateral cartilage". *European Archives of Oto-Rhino-Laryngology* 272.2 (2015): 497-504.

23. Salyer KE. "Primary correction of the unilateral cleft lip nose: a 15-year experience". *Plastic and Reconstructive Surgery* 77 (1986): 558-566.

24. Varedi P, et al. "Spreader graft placement: A simplified technique for young surgeons". *International Journal of Oral and Maxillofacial Surgery* 43 (2014): 1216-1217.

25. Saedi B., et al. "Spreader flaps do not change early functional outcomes in reduction rhinoplasty: a randomized control trial". *American Journal of Rhinology and Allergy* 28 (2014): 70-74.

26. Manavbaşı YI and Başaran I. "The role of upper lateral cartilage in dorsal reconstruction after hump excision: section 1. Spreader flap modification with asymmetric mattress suture and extension of the spreading effect by cartilage graft". *Aesthetic Plastic Surgery* 35 (2011): 487-493.

27. Rhee JS., et al. "A systematic review of patient-reported nasal obstruction scores". *JAMA Facial Plastic Surgery* 16.3 (2014): 219-225.

28. Most SP. "Anterior septal reconstruction: outcomes after a modified extracorporeal septoplasty technique". *Archives of Facial Plastic Surgery* 8.3 (2006): 202-207.

29. Andre RF, et al. "Endonasal spreader graft placement as treat-

Citation: Eldeeb W., et al. "A Comparative Study of Open Septorhinoplasty with Spreader Graft and Auto-Spreader Flap in Patients with Nasal Obstruction". *Acta Scientific Otolaryngology* 2.6 (2020): 20-28.

ment for internal nasal valve insufficiency: no need to divide the upper lateral cartilages from the septum". *Archives of Facial Plastic Surgery* 6.1 (2004): 36-40.

30. Gruber RP, *et al.* "Dorsal reduction and spreader flaps". *Aesthetic Surgery Journal* 31.4 (2011): 456-464.

31. Helal MZ., *et al.* "Effects of rhinoplasty on the internal nasal