Letter in Response to Virtual Reality (VR) for Vestibular Rehabilitation: A Pilot Study Comparing VR with Conventional Rehabilitation with Conventional Rehabilitation Alone

Kusum Shankar Asnani* and Gurdas Viguruji Singh

GKT School of Medical Education, King's College London, UK

***Corresponding Author:** Kusum Shankar Asnani, GKT School of Medical Education, King's College London, UK.

Received: October 14, 2020

Published: October 30, 2020

© All rights are reserved by **Kusum Shankar Asnani and Gurdas Viguruji Singh**.

We read the innovative randomised control trial by Srinivas Doriasala, *et al.* on utilising Virtual Reality (VR) for Vestibular Rehabilitation with interest [1]. Whilst the study is important, there is scope for further research.

This study has not set out a clear enough exclusion criteria. Firstly, the paper states to investigate the outcome of VR on peripheral vestibular disorders. However, in groups A and B, patients 2, 3, 4, 7 and 8 suffer from a central vestibular loss such as migrainous vertigo and cerebellar degeneration. Additionally, motion sickness is on the list and that does not fall under peripheral nor central vestibular loss thus affecting the credibility of the results. The study could have cultivated more robust results had the patients been either suffering from solely peripheral or central vestibular problems. Earlier literature indicated that Meniere's Disease should not be included for VT due to its spontaneous presentation [2]. A clear cut description of the exclusion and inclusion criteria would be beneficial.

There is a limited explanation on how the VT would be implemented and the form of support and education given to patients prior and during the use. It would have been useful to know if the exercise replicated the traditional Cawthorne-Cooksey exercises [1]. This will highlight whether the results produced are a result of solely VT or the change in the actual content of vestibular rehabilitation.

The outcomes measured from this study were limited to only the Dizziness Handicap Inventory. Whilst it allows readers to notice the quantified reduction of dizziness in groups A and B, it would be helpful to measure other factors aiding or affecting the

improvement of symptoms. For example, in a patient with additional psychological features such as anxiety or depression, it could potentially hinder vestibular compression [2]. Therefore, to factor that in, a depression scale such as the Beck Depression Inventory, before, during and after treatment would highlight conflating factors which could confound any outcomes [3].

As the patients included in the study are of a wide age range, the study could have included feedback on user-friendliness and compliance of VT. Specifically, it would have been useful to analyse how the older population received VT and their feedback. This would allow further developments and progress on developing this technology for different groups.

For the benefit of future VR implementations, it is crucial to trial it out on a bigger sample size to note for statistically significant results and adverse effects or symptoms. Patients with vestibular impairment could experience nausea and vomiting. Using VT can cause cybersickness [4], however, the author's study did not include any cybersickness. Hence, a bigger sample size could capture these adverse effects.

Lastly, it would have been insightful if the authors had explored the costs of funding VR and its impacts on the health economy. Another point that could have been touched on was the cost-effectiveness and evidence to conduct VR on its own or alongside VRT [5].

Bibliography

1. Srinivas Doriasala, *et al.* "Virtual Reality (VR) for Vestibular Rehabilitation: A Pilot Study Comparing VR with Conventional Rehabilitation with Conventional Rehabilitation Alone". *Acta Scientific Otolaryngology* (2020): 18-22.

2. Byung In Han., *et al.* "Vestibular Rehabilitation Therapy: Review of Indications, Mechanisms, and Key Exercises". *Journal of Clinical Neurology* (2011): 184-196.
3. M Pavlou., *et al.* "The effect of virtual reality on visual vertigo symptoms in patients with peripheral vestibular dysfunction: A Pilot Study". *Journal of Vestibular Research* (2012): 273-281.
4. Tohru Kiryu and Richard HY So. "Sensation of presence and cybersickness in applications of virtual reality for advanced rehabilitation". *Journal of Neuroengineering and Rehabilitation* (2007).
5. Mathieu Bergeron., *et al.* "Use of Virtual Reality Tools for Vestibular Disorders Rehabilitation: A Comprehensive Analysis". *Advances in Medicine* (2015): 1-9.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667