Lens Dislocation in a 9-Year-Old Child with Marfan Syndrome

Japhet Pobanou Thera*, Pierre Dakouo, Jean Marc Lere Tiama and Ali Konipo

Institute of African Tropical Ophthalmology, Bamako, Mali

*Corresponding Author: Japhet Pobanou Thera, Institute of African Tropical Ophthalmology, Bamako, Mali.

Received: May 02, 2020

Published: May 22, 2020

© All rights are reserved by Japhet Pobanou Thera., et al.

Abstract

Introduction: Marfan syndrome is a rare genetic disorder linked to the mutation in the FBN1 gene, which codes for fibrillin. It is transmitted in the autosomal dominant mode. Its prevalence is estimated at 1/5000 individuals. Cardiovascular, skeletal and ophthalmic disorders are the most frequent

Clinical Observation: We report the case of a 9-year-old male child with uneventful family history brought by his parents for blurred vision. The visual acuity without correction was "count fingers at 2 meters". Slit lamp examination showed bilateral posterior dislocation of the lens in the vitreous, the Intra-ocular pressure (IOP) was 12 mmHg in both eyes, the fundus was normal. The general examination found a tall child measuring 1.48 m of height for 26 kg. He had slender limbs showing the sign of the thumb (Steinberg) and the sign of the wrist along with a moderate pectus excavatum. Cardiac ultrasound showed dilation of the ascending aorta. According to Gant's criteria (Ghent), we concluded to the diagnosis of Marfan syndrome.

Discussion: The diagnosis of Marfan syndrome is difficult to establish. At first, it is mainly clinical with the association of major and minor signs. Genetic research is possible as well as skin biopsy.

Lens ectopia is the most common ophthalmologic manifestation in Marfan syndrome. According to Gant's criteria, we concluded to the diagnosis of Marfan syndrome.

Conclusion: Marfan syndrome is oftentimes discovered during an ophthalmic consultation. The management of its various disorders and the follow-up imply a frank collaboration between many practitioners.

Keywords: Dislocation of the Lens; Marfan; Child

Introduction

Marfan syndrome is a genetic disorder linked to the mutation of the FBN1 gene, which codes for fibrillin on chromosome 15, sometimes to TGFBR1 or 2 [1-3]. It is transmitted in the autosomal dominant mode. Its prevalence is estimated at 1/5000 individuals [4]. It was first described by Antoine - Bernard Marfan in a young girl with unusual musculoskeletal characteristics in 1896, while Bürger first described ophthalmological characteristics in 1914 [5,6]. Multisystemic disorder, cardiovascular, skeletal and ophthalmological disorders are the most frequent [3,7]. The diagnosis of Marfan's syndrome is difficult to establish. At first, it is mainly clinical with the association of major and minor signs [8,9]. Genetic research is possible as well as a skin biopsy. Histological studies have shown an abnormal expression of fibrillin in the lens capsule and the ciliary epithelial surface at the attachment of the zonules [10]. Ectopia of the lens is the most common ocular manifestation

in Marfan syndrome. It is found in 50 to 80% of affected eyes and around 41% are initially seen for ocular pathology and diagnosed by an ophthalmologist [4,11,12]. Rare pathology, we report the case of a child with bilateral posterior dislocation of the lens.

Clinical Observation

He was a 9-year-old child with no known family history brought in our office by his parents for blurred vision. The visual acuity without correction was "count fingers at 2 meters". Slit lamp examination showed bilateral posterior dislocation of the lens in the vitreous (Figure 1), IOP was 12 mmHg in both eyes, the fundus was normal. The general examination shows a tall child measuring 1.48 m for 26 kg, slender limbs showing signs of the thumb (Steinberg's sign) and the wrist, a moderate pectus excavatum (Figure 2). Cardiac ultrasound showed dilation of the ascending aorta (Figure 2). According to Gant's criteria (Ghent), we made the diagnosis of Marfan syndrome.

Figure 1: Bilateral lens dislocation in the vitreous.

Figure 2: Skeletal features: slender limbs (a), sign of the thumb (b), sign of the wrist (c).

Discussion

Lens ectopia is the most common ophthalmologic manifestation in Marfan syndrome, 50 to 80% [4,11,12]. Although the disease is first described in a young girl [5], Charif-Chefchaoui, *et al.* reported a male predominance in a series of 20 cases with an average age of 14 years whose extremes range from 4 to 41 years [11]. A male predominance of 53% is reported by Mary J Roman, *et al* [13]. However, Kumar A., *et al.* reported a female predominance in a series of 4 cases including 3 women [14]. Laffargue F, *et al.* reported a series of 17 cases aged less than 10 years [15]. The ophthalmic complications of Marfan are among others, posterior dislocation of the clear or opacified lens, rhegmatogenic detachment of the retina, secondary glaucoma [11,14,16]. In our series the fundus was normal as in that of Omolase, *et al.* [4] in a 12-year-old girl. Cardiovascular and skeletal damage are also common. Dilation of the ascending aorta is a major sign of cardiac involvement [7] which can be complicated by dissection and aortic rupture, aortic or mitral insufficiency leading to cardiac insufficiency [2,15]. The skeletal manifestations reported by several authors are marked, among other things, by the large size of the subjects, ligament hyperlaxity, scoliosis, signs of the thumb and wrist, pectus carinatum and pectus excavatum, flat feet [2-8,10-15]. Scoliosis and deformation of the rib cage in the keel or funnel can lead to a restrictive disabling syndrome, thus the frequency of apical bubbles and spontaneous pneumothorax is increased [2]. Pneumothorax affects 5 to 11% of patients [6]. In our case, there was pectus excavatum without lung damage.

Conclusion

A multi-system disorder, Marfan syndrome is most often discovered during an ophthalmological consultation. The management of the various disorders and the follow-up imply a frank collaboration between ophthalmologists, cardiologists, orthopedists, pulmonologists and many others.

Conflict of Interest

No conflict of interest.

Bibliography

1. Dean JCS. "Marfan Syndrome : clinical, diagnosis and management". *European Journal of Human Genetics* 15 (2007): 724-333.
2. Le Parc J-M. "Le syndrome de Marfan". *Encyclopédie Orphanet* (2005): 1-74.
3. De Laage de Meux P, *et al.* "Déplacements congénitaux et acquis du cristallin". *EMC Ophtalmologie* (2005) : 21.
4. Omolase CO, *et al.* "Bilateral Ectopia Lentis in Marfan's Syndrome: A Case Report". *Clinics in Surgery* 3 (2018): 2244.
5. Gordana S, *et al.* "Ocular features of Marfan syndrome". *Medicine and Biology* 15.1 (2008): 37-40.
6. Neuville M, *et al.* "Manifestations respiratoires du syndrome de Marfan". *Revue des Maladies Respiratoires* 32.2 (2015): 173-181.
7. De Paepe A, *et al.* "Revise criteria for the Marfan syndrome". *American Journal of Medical Genetics* 62.4 (1996) :417-426.
8. Waduthantri S. "Ocular manifestations of Marfan's syndrome". *Medical Journal of Dr. D.Y. Patil Vidyapeeth* 10 (2017): 118-119.
9. Pyeritz RE, *et al.* "Immunohistochemical localization of fibrillin in human ocular tissues. Relevance to the Marfan Syndrome". *Archives of ophthalmology* 113 (1995): 103-109.
10. Linda M, *et al.* "Le syndrome de Marfan". *Cahiers de la Pédiatrie* 46.223 (2009): 37.
11. Charif-chefchaoui M, *et al.* "Les manifestations oculaires du syndrome de Marfan : à propos de 20 cas". *Médecine du Maghreb* (2001): 87.
12. Rubin SE, *et al.* "Ocular manifestations of autosomal dominant systemic conditions". *Duane's Clinical Ophthalmology* 3.58 (2006).

13. Mary J., *et al.* "Association of Age and Sex with Phenotype: The NHLBI GenTAC Registry". *Circulation: Cardiovascular Genetics* 10.13 (2007): 1647.
14. Kumar A., *et al.* "Bilateral posterior lens dislocation in Marfan's syndrome". *Indian Journal of Ophthalmology* 37 (1989): 202-204.
15. Laffargue F., *et al.* "Signes cliniques évocateurs d'un syndrome de Marfan chez l'enfant de moins de 10 ans". *Archives de Pédiatrie* 20.11 (2013): 1193-1200.
16. Mema V., *et al.* "Ocular complications of marfan syndrome. Report of two cases". *Hippokratia* 14.1 (2010): 45-47.

Assets from publication with us

- Prompt Acknowledgement after receiving the article
- Thorough Double blinded peer review
- Rapid Publication
- Issue of Publication Certificate
- High visibility of your Published work

Website: www.actascientific.com/

Submit Article: www.actascientific.com/submission.php

Email us: editor@actascientific.com

Contact us: +91 9182824667