---

*Pachydermoperiostosis:
Classic Presentation of a Rare Disease*

Rajat Kharbanda, TG Sundaram, Latika Gupta

Mediterr J Rheumatol 2020;31(2):214-5

Pachydermoperiostosis: Classic Presentation of a Rare Disease

Rajat Kharbanda, TG Sundaram, Latika Gupta

Department of Clinical Immunology & Rheumatology, Sanjay Gandhi Postgraduate Institute of Medical Sciences, Lucknow, India

Mediterr J Rheumatol 2020;31(2):214-5

<https://doi.org/10.31138/mjr.31.2.214>

Article Submitted: 16 Mar 2020; Revised Form: 10 Jun 2020; Article Accepted: 19 Jun 2020; Available Online: 30 Jun 2020

Keywords: Clubbing, pachydermoperiostosis, primary hypertrophic osteoarthropathy, hypertrophic osteoarthropathy

A thirty-four-year-old gentleman reported with non-inflammatory knee arthralgias of 30 years duration. He presented with limitation of movement in both knees for the past one year. He had experienced mechanical pain in the ankles and feet on occasion. He was a non-smoker with an unremarkable history.

On examination, the prominent folds of skin were noted on the patient's forehead (*Figure 1A*), apart from an oily facial skin and grade four clubbing of all fingers (*Figure 1B*) and toes. Terminal knee extension was restricted. The right ankle was swollen and tender with painful limitation of terminal movement, although movement was preserved on the left side, which was swollen but non-tender. Considering the onset of clubbing in childhood, with a family history of similar manifestations in the absence of lung or cardiac disease, a diagnosis of primary hypertrophic osteoarthropathy (pachydermoperiostosis) was deemed likely.

On further review, he reported noting disproportionately larger hands and feet as compared with peers, a perception shared by his sister as well. Besides, his sister has

a similar affliction of the fingers. Laboratory investigations and chest radiograph were unremarkable. The radiograph showed irregular cortical thickening of the bilateral femur, the subperiosteal bone formation of proximal tibia and fibula, with irregular-

subperiosteal bone formation and cortical thickening of the distal tibia, fibula (*Figures 2B,C*). Radiographs of bilateral hands showed enlargement of distal ulna and radius, metacarpals, phalanges (*Figure 2A*). He was prescribed Nonsteroidal anti-inflammatory drugs (NSAIDs) with a plan to consider bisphosphonate in the event of inadequate response. On follow up visits, the patient had relief in his articular symptoms.

Pachydermoperiostosis, also known as Touraine-Solente-Golé syndrome, is an autosomal-dominant disorder with variable penetrance. In its complete form, it is characterized by pachyderma (thickening of the facial skin), skeletal changes (periostosis), excessive sweating (hyperhidrosis), and acropachy (digital clubbing).^{1,2} Pachydermia is the most frequent skin symptom. Digital clubbing occurs in 89% of cases, and the classic radiologic findings in 80-97% of patients.³ Borochowitz and Rimoïn¹ proposed the presence of at least two of the four criteria - a history of familial transmission; pachydermia; digital clubbing, and skeletal manifestations such as pain or signs of radiographic periostitis for diagnosis. This patient fulfilled all four criteria.

The differential diagnoses include lung cancer, acromegaly, psoriatic arthritis, carcinomatous polyarthritis, thyroid acropachy, fluorosis and hypervitaminosis A.⁴ With classic findings, and no history or clinical findings explaining any of the above diagnoses, the diagnosis was never in doubt. Since prostaglandin E2 plays a central role in the pathogenesis of this disease, NSAIDs are the first line of treatment.⁵ Pamidronate is reportedly effective in NSAID-resistant cases.⁶

Corresponding Author:

Latika Gupta
Assistant Professor
Department of Clinical Immunology &
Rheumatology
Sanjay Gandhi Postgraduate Institute of
Medical Sciences
Lucknow, India, 226014
E-mail: dratikagupta@gmail.com
Tel.: +91 522 2495182

Figure 1(A). Image depicting clubbing of all digits.

Figure 1(B). Image of prominent folds on the patient's forehead.

Figure 2(A). Radiograph of hand demonstrating widening of distal end of radius and ulna (white arrow).

Figure 2(B). Radiograph of knee depicting irregular cortical thickening of bilateral femur (white arrow).

Figure 2(C). Radiograph of legs showing sub-periosteal bone formation (white arrow).

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We would like to thank the department of radiology at SGPGIMS, Lucknow for the radiographs.

REFERENCES

1. Borochowitz Z, Limoind DL. Pachydermoperiostosis. Birth Defect Encyclopaedia 1990;1349-50.
2. Supradeeptha C, Shandilya SM, Vikram Reddy K, Satyaprasad J. Pachydermoperiostosis - a case report of complete form and literature review. J Clin Orthop Trauma 2014;5:27-32.
3. Schumacher HR, Jr. Hypertrophic osteoarthropathy: rheumatologic manifestations. Clin Exp Rheumatol 1992;10:35-40.

4. Pathak H, Lonsdale R, Dhatariya K, Mukhtyar C. Carcinomatous polyarthritides as a presenting manifestation of papillary carcinoma of thyroid gland. Indian J Rheumatol 2016;11:164-6.
5. Zhang H, Yang B. Successful treatment of pachydermoperiostosis patients with etoricoxib, aescin, and arthroscopic synovectomy: Two case reports. Medicine 2017;96(47):e8865.
6. Guyot-Drouot MH, Solau-Gervais E, Cortet B. Rheumatologic manifestations of pachydermoperiostosis and preliminary experience with bisphosphonates. J Rheumatol 2000;27:2418-2423.