Biological, morphological and phytobiogeographic diversity of *Malva subovata* (DC.) Molero & J.M. Monts. (Malvales Malvaceae) in the Tlemcen region, Algeria

Sarra Ghalem*, Faïçal Hassani, Sid Ahmed Aouadj & Ibtissam Sarah Bouayed

Laboratory of Ecology and Management of Natural Ecosystems, Department of Ecology and Environment, Faculty of Nature and Life Sciences and Earth and Universe Sciences, Abou Bakr Belkaid University, Tlemcen, Algeria

*Corresponding author, email: sarah.ghalem@outlook.com

ABSTRACT

The vegetation of the region of western Algeria presents a good example of the study of phyto-diversity. The method of floristic analysis is a main element in the knowledge of natural environments and its rich flora. In my two stations, we carried out floristic surveys, these are a set of observations on the environment and on the vegetation. The study presented is only a part of the objectives of the Laboratory of Plant Ecology focused on the floristic diversity of *Malva subovata* (DC.) Molero & J.M. Monts. (Malvales Malvaceae) on the mastery of the biological, morphological and phytobiogeographic capital of the two stations “Boussdra and Hammam Boughrara” in the Tlemcen region. From the treatment of biodiversity indices and the floristic surveys, we were able to conclude that the study area is undergoing a regressive dynamic of vegetation.

KEY WORDS

Biology; *Malva subovata*; morphology; phytobiogeographic studies; Tlemcen.

Received 31.08.2020; accepted 30.01.2021; published online 15.02.2021

INTRODUCTION

Phytogeographic and chorological studies are a very important basis for any conservation and restoration attempt (Quezel, 1999; Aouadj et al., 2020a). They allow us to know the history of a given region and is a true model for interpreting the phenomena of regression (Aouadj et al., 2020b, c).

Aouadj et al. (2020d, e) report that: the floristic sectors and subsectors subdivision is a reflection of the narrow climate-vegetation dependence, based on the phytogeographic division of Barry et al. (1974). Vegetation plays a fundamental role in the structure and functioning of ecosystems, of which

it constitutes an expression of the biological potential (Bouayed et al., 2019).

From an ecological point of view, the Mediterranean basin is considered a hotspot because of its very important biological diversity, including endemic plant species. The estimated plant richness is of 25000 species. Mediterranean-type forests account for about twice as many woody species as European-type forests (247 versus 135) (Barbero et al., 1990; Quézel & Medail, 1995; Medail & Quézel, 1999). This plant diversity varies from country to country depending on geographical location. The plant richness of the North Africa is estimated between 5000 to 5300 species, represents 15% of that of the Mediterranean basin (Quézel, 1999). This

wealth is represented by 3800 species, 900 of which are endemic to Morocco. According to Quézel & Santa (1962–1963 in Vela & Benhouhou, 2007), the Algerian flora comprises 3139 species (3744 taxa) including 464 endemic species.

In the assessment established by Medail & Quézel (1999) and Aouadj et al. (2020f), the Mediterranean forest is made up of approximately 247 woody species compared to European forests (13 species).

Mediterranean plant biodiversity is produced, for many, from a traditional and harmonious use of the environment by humans (Medail & Quézel, 1999). Despite the incessant attacks they have suffered for a millennium, the Mediterranean forests still offer, in places, appreciable development. There are many methods of studying vegetation. The choice of a technique generally depends on the nature of the vegetation and the object of the study (Kerzabi, 2017). Numerous research programs, through international publications, have underlined the major role of various regions of Tlemcen as an essential reservoir of plant biodiversity (Bouazza et al., 1996).

Mankind, in its nature, has always used its entourage, in particular its natural environment, to meet its needs. It is characterized by its size, density, growth, cultivation, construction, feeding, gathering and predation. All these functions have a negative effect on natural resources. Thus, it is the vital activities of populations that directly influence the components of the environment and generally modify it (Bensenane, 2015; Aouadj et al., 2020g).

In order to better understand the characteristics of plant formations, the study carried out is essentially based on the analysis of exhaustive inventories of the species encountered in the *Malva subovata* (DC.) Molero & J.M. Monts (= *Lavatera maritima* Gouan) (Malvales Malvaceae) (Fig. 1) formations in the 2 study stations, which are characterized by a count of the species, with an identification of their morphological and biological types, and their floristic characters. We will also get to know the study region, which has a very diverse flora and is closely linked to various disturbance factors through biodiversity indices (Ghalem et al., 2020).

MATERIAL AND METHODS

The analysis of the plant structure takes into ac-

Figure 1. *Malva subovata* (from Ghalem et al., 2020).

Figures 2, 3. Study areas.

count the method of floristic surveys, which requires us to list all the plant species present. This floristic list changes from one station to another, from one year to the next in the same station (Raunkiaer, 1934; Bouayad, 2017; Kerzabi, 2017).

Our surveys were carried out during the optimal vegetation period from March to May during the year 2019.

The coordinates of the stations were obtained using a GPS (Bousdra: 001°37 W - 34°42 N; Hammam Boughrara: 001°40 W - 34°52 N; Tlemcen, Algeria). Each of these readings includes the general characteristics linked to the station itself (Figs. 2, 3). These are: place and date; the altitude; slope; exposure t; plant cover.

Currently, the survey method is based on the Braun-Blanquet method (1952) known as Zuricho-Montpellier which consists in determining the smallest surface called "Minimum area" (Braun-Blanquet, 1952; Gounot, 1969).

RESULTS

The floristic composition (Bousdra station)

The floristic inventory of the study area allowed us to invent 36 taxa divided into 16 families and 20 genera (Table 1).

The floristic composition (Hammam Boughrara station)

The floristic inventory of the study area allowed us to invent 68 taxa divided into 26 families and 36 genera (Table 2).

DISCUSSION

Bousdra station

Our study area has about 36 species, they belong

Taxa	Family	Biological type	Morphol. type
<i>Asteriscus maritimus</i>	Asteraceae	Ch	L.v
<i>Asperula hirsuta</i>	Rubiaceae	Th	H.a
<i>Avena sterilis</i>	Poaceae	Th	H.a
<i>Anarrhinum fruticosum</i> subsp. <i>fruticosum</i>	Plantaginaceae	Ch	L.v
<i>Ballota hirsuta</i>	Lamiaceae	He	H.v
<i>Campanula dichotoma</i>	Campanulaceae	Th	H.a
<i>Calycotome intermedia</i>	Fabaceae	Ch	L.v
<i>Centaurea involucreta</i>	Asteraceae	He	H.v
<i>Centaurea pullata</i>	Asteraceae	He	H.v
<i>Ceratonia siliqua</i>	Fabaceae	Ph	L.v
<i>Convolvulus althaeoides</i>	Convolvulaceae	Th	H.a
<i>Convolvulus cantabrica</i>	Convolvulaceae	He	H.v
<i>Cistus villosus</i>	Cistaceae	Ch	L.v
<i>Echium vulgare</i>	Boraginaceae	He	H.v
<i>Eruca vesicaria</i>	Brassicaceae	Th	H.a
<i>Fagonia cretica</i>	Zygophyllaceae	Th	H.a
<i>Fumanathymifolia</i>	Cistaceae	Ch	L.v

<i>Galactites duriae</i>	Asteraceae	Th	H.a
<i>Genista tricuspidata</i>	Fabaceae	Ch	L.v
<i>Inula crithmoides</i>	Asteraceae	Ch	L.v
<i>Malva subovata</i>	Malvaceae	Ch	L.v
<i>Lavandula dentata</i>	Lamiaceae	Ch	L.v
<i>Lavandula stoechas</i>	Lamiaceae	Ch	L.v
<i>Lobularia maritima</i>	Brassicaceae	Th	H.a
<i>Nepeta multibracteata</i>	Lamiaceae	He	H.v
<i>Olea europea</i>	Oleaceae	Ph	L.v
<i>Quercus coccifera</i>	Fagaceae	Ph	L.v
<i>Sedum sediforme</i>	Crassulaceae	Ch	H.v
<i>Sedum acre</i>	Crassulaceae	Ch	H.v
<i>Stipa parviflora</i>	Poaceae	Ch	H.v
<i>Stachys ocymastrum</i>	Lamiaceae	Th	H.a
<i>Tetraclinis articulata</i>	Cupressaceae	Ph	L.v
<i>Pallenis spinosa</i>	Asteraceae	He	H.v
<i>Phylleria angustifolia</i>	Oleaceae	Ph	L.v
<i>Plantago albicans</i>	Plantaginaceae	He	H.v
<i>Pistacia lentiscus</i>	Anacardiaceae	Ph	L.v

Table 1. Floristic composition of Bousdra station.

Figure 4. Percentage of morphological types of Bousdra station.

Figure 5. Biological types of the floral procession of Bousdra station.

to the subphylum Gymnosperms and Angiosperms with 18 families. The families most represented on a specific level are the Asteraceae with a percentage of 16.66%, the Lamiaceae with the percentage of 13.88%, the Fabaceae remain very little dominant with a percentage of 8.33% then the Cistaceae, Crassulaceae, Oleaceae, Brassicaceae, Poaceae and Plantaginaceae with the same percentage of 5.55, the other families have the same number with a very low rate (Figs. 4, 5).

On the morphological level, we notice that the perennial woody plants present the highest number in the station Bousdra station with a percentage of 44%, then the perennial grasses with 31% and finally the annual grasses with the percentage of 25%.

In this station we notice that the Chamaephytes are the best represented with a percentage of 36%, the Therophytes with 25%, the Hemicryptophytes 22%, the Phanerophytes with a low percentage of 17%. We also notice that there is a absence of Geophytes.

- The specific richness of the Bousdra resort: 18
- Total number of species: 36

$$D = S - 1 / \ln N$$

$$D = 18 - 1 / \ln 36 \quad D = 4.74$$

Simpson indices (Is):

$$Is = 1 / \sum Pi^2$$

$$Pi = ni/N = 0.982 \quad \rightarrow \quad Pi^2 = 0.0782$$

$$Is = 12.78$$

Equitability of the simpson index: $Es = Is - 1 / S - 1 = 0.693$

$$\text{Shannon-Weaver indicator: } H' = - \sum pi \log_2 pi$$

$$H' = -0.982 \log 0.982 \quad H' = 0.0257$$

The equity index measures the distribution of individuals within species, regardless of species richness.

$$E = H' / \log S = 0.020 \quad \rightarrow \quad \text{Abundance of individuals of each species are equal}$$

Hamam Boughrara station

Our study area has about 68 species, they belong to the sub-phylum Gymnosperms and Angiosperms with 26 families. The families most represented on a specific level are the Asteraceae with a percentage of 20.58%, and the Poaceae with the percentage of 14.7%, then the Apiaceae 8.82%, the Fabaceae, Brassicaceae, Liliaceae, Cistaceae and Plantaginaceae with the same percentage of 4.47%, the other families have the same number with a very low rate (Figs. 6, 7).

Morphologically, we notice that annual herbs have the highest number in Hamam Boughrara station with a percentage of 47%, then perennial herbs with 44% and finally perennial woody with a percentage of 9%. In this station we notice that the Therophytes are the best represented with the same percentage of 49%, the Hemicryptophytes

Taxa	Family	Biological type	Morphol. type
<i>Aegilops triuncialis</i>	Poaceae	Th	H.a
<i>Ajuga iva</i>	Lamiaceae	Th	H.a
<i>Allium porrum</i> subsp. <i>polyanthum</i>	Alliaceae	He	H.v
<i>Allium roseum</i>	Alliaceae	He	H.v
<i>Ammoides verticillata</i>	Apiaceae	Th	H.a
<i>Anacyclus vallentinus</i>	Asteraceae	Th	H.a
<i>Anagalis arvensis</i> subsp. <i>latifolia</i>	Primulaceae	TH	H.a
<i>Anagalis arvensis</i> subsp. <i>phonicea</i>	Primulaceae	Th	H.a
<i>Anthyllis tetraphylla</i>	Fabaceae	Th	H.a
<i>Arisarum vulgare</i>	Araceae	Ge	H.v
<i>Asphodelus microcarpus</i>	Liliaceae	Ge	H.v
<i>Asteriscus maritimus</i>	Asteraceae	Ch	L.v
<i>Asparagus acutifolius</i>	Liliaceae	Ge	H.v
<i>Atractylis carduus</i>	Asteraceae	He	H.v
<i>Atractylis concellata</i>	Asteraceae	He	H.v
<i>Avena sterilis</i>	Poaceae	Th	H.a
<i>Asparagus stipularis</i>	Liliaceae	Ge	H.v
<i>Ballota hirsuta</i>	Lamiaceae	He	H.v
<i>Bellis annua</i>	Asteraceae	Th	H.a
<i>Brachypodium distachyon</i>	Poaceae	Th	H.a
<i>Briza minor</i>	Poaceae	Th	H.a
<i>Bromus madritensis</i>	Poaceae	Th	H.a
<i>Bromus rubens</i>	Poaceae	Th	H.a
<i>Cakile maritima</i>	Brassicaceae	Th	H.a
<i>Calendula arvensis</i>	Asteraceae	Th	H.a
<i>Calendula suffruticosa</i>	Asteraceae	Th	H.a
<i>Carduus pycnocephalus</i>	Asteraceae	Th	H.a
<i>Carthamus coerulus</i>	Asteraceae	Th	H.a
<i>Convolvulus althaeoides</i>	Convolvulaceae	Th	H.v
<i>Convolvulus tricolor</i>	Convolvulaceae	Th	H.a
<i>Cynodon dactylon</i>	Poaceae	Ge	H.v
<i>Cytisus villosus</i>	Fabaceae	Ch	L.v
<i>Dactylis glomerata</i>	Poaceae	Ge	H.v
<i>Daucus carota</i> subsp. <i>maximus</i>	Apiaceae	He	H.v
<i>Echinophora spinosa</i>	Apiaceae	Ch	H.v
<i>Echinops spinosus</i>	Asteraceae	He	H.v
<i>Echium vulgare</i>	Boraginaceae	He	H.v
<i>Fagonia cretica</i>	Zygophyllaceae	He	H.v
<i>Ferula communis</i>	Apiaceae	He	H.v
<i>Ferula lutea</i>	Apiaceae	He	H.v
<i>Fumana thymifolia</i>	Cistaceae	Ch	L.v
<i>Galactites duriae</i>	Asteraceae	Th	H.A
<i>Helianthemum apertum</i>	Cistaceae	Th	H.a
<i>Helianthemum hirtum</i>	Cistaceae	Th	H.a
<i>Hordeum murinum</i>	Poaceae	Th	H.a
<i>Lagurus ovatus</i>	Poaceae	Th	H.a
<i>Lavandula multifida</i>	Lamiaceae	Ch	H.v
<i>Malva subovata</i>	Malvaceae	Ch	L.v
<i>Limonium sinuatum</i> subsp. <i>eusinuatum</i>	Plumbaginaceae	He	H.v
<i>Lobularia maritima</i>	Brassicaceae	Th	H.a
<i>Olea europaea</i>	Oleaceae	Ph	L.v
<i>Phagnalon saxatile</i>	Asteraceae	Ch	H.v
<i>Pallenis spinosa</i>	Asteraceae	He	H.v
<i>Papaver rhoeas</i>	Paveraceae	Th	H.a
<i>Paronychia argentea</i>	Caryophyllaceae	He	H.v
<i>Plantago albicans</i>	Plantaginaceae	He	H.v
<i>Plantago lagopus</i>	Plantaginaceae	He	H.v
<i>Plantago psyllium</i>	Plantaginaceae	Th	H.a
<i>Pistacia lentiscus</i>	Anacardiaceae	Ph	L.v
<i>Thapsia garganica</i>	Apiaceae	He	H.v
<i>Reseda alba</i>	Resedaceae	Th	H.a
<i>Rumex bucephalophorus</i>	Polygonaceae	Th	H.a
<i>Ruta chalepensis</i>	Rutaceae	Ch	H.v
<i>Sedum sediforme</i>	Crassulaceae	He	H.v
<i>Sonchus asper</i>	Asteraceae	Th	H.a
<i>Sinapis alba</i>	Brassicaceae	Th	H.a
<i>Ononis pubescens</i>	Fabaceae	Th	H.a
<i>Opuntia maxima</i> = <i>O. ficus-indica</i>	Cactaceae	Ch	H.v

Table 2. Floristic composition of Hammam Boughrara station.

Figure 6. Percentage of morphological types of the Hammam Boughrara station.

Figure 7. Biological types of the floral procession of the Hammam Boughrara station.

with 26%, the Chamaephytes with a percentage of 13%, the Geophytes 9% and finally we notice that there is a very low percentage of Phanerophytes of 3%.

Total number of species: 68

$$D = S - 1 / \ln N \rightarrow D = 26 - 1 / \ln 68 \rightarrow D = 5.92$$

Simpson indices (Is):

$$Is = 1 / \sum Pi^2 \rightarrow Pi = ni/N = 0.987 \rightarrow Pi^2 = 0.0872 \rightarrow Is = 11.46$$

Equitability of the Simpson index:

$$Es = Is - 1 / S - 1 = 0.418$$

Shannon-Weaver indicator: $H' = - \sum pi \log_2 pi$

$$H' = -0.987 \log 0.987 \rightarrow H' = 0.0186$$

So only one species is present.

The equity index measures the distribution of individuals within species, regardless of species richness.

$E = H' / \log S = 0.013 \rightarrow$ Abundance of individuals of each species are equal

CONCLUSIONS

The exhaustive inventory carried out at the two study stations (Bousdra, Hammam Boughrara) enabled us to study the following characterizations:

biological and morphological, followed by the distribution of families.

This study allowed us to bring out the following results:

dominance of perennial woody plants in the Bousdra resort and the dominance of annual herbaceous plants in the Hammam Boughrara resort from a morphological point of view;

importance of therophytes which confirms the phenomenon of therophytization in the Hammam boughrara station;

type vegetation: TH > He > Ch > Ge > PH for the Hama boughrara station;

special attention is generally paid to the distribution of Therophytes, the proportion of which in the Mediterranean region is around 50%;

the rate of theophytes is linked, whatever the scale of the analysis and the level of perception adopted, to the openness of the vegetation and the overall humidity of the environment;

type vegetation: Ch > Th > He > Ph for the Bousdra station;

grazing favors the overall establishment of the Chamaephytes often refused by the herd;

we consider the Chamaephytes to be better adapted to low temperatures and aridity;

the most dominant families in the field for the two stations are: Asteraceae, Poaceae and Lamiaceae;

the biological types constitute indices of the life strategy of species.

REFERENCES

- Aouadj S.A., Nasrallah Y. & Hasnoui O., 2020a. Regional phytogeographic analysis of the flora of the Mounts of Saida (western Algeria): evaluation-restoration report. *Biodiversity Journal*, 11: 25–34. <https://doi.org/10.31396/Biodiv.Jour.2020.11.1.25>.
- Aouadj S.A., Hasnoui O. & Nasrallah Y., 2020b. Approche ethnobotanique et inventaire floristique des plantes médicinales dans la région de Doui Thabet (Saida-Ouest Algérien). *Phyto Chem & Bio Sub Journal*, 14: 92–104. <https://doi.org/10.163.pcbsj/2020.14.-1-92>
- Aouadj S.A., Nasrallah Y. & Hasnoui O., 2020c. Ecological characterization and evaluation of the floristic potential of the forest of Doui Thabet (Saida Western Algeria) in the context of the restoration. *Ecology, Environment and Conservation*, 26: 266–278.
- Aouadj S.A., Nasrallah Y. & Hasnoui O., 2020d. Note on the orchids of mounts of Saida (Saida Western Algeria) in the context of the restoration. *Ecology, Environment and Conservation*, 26: 37–45.
- Aouadj S.A., Nasrallah Y., Hasnoui O. & Khatir H., 2020e. Impacts of anthropogenic pressure on the degradation of the forest of Doui Thabet (Saida, Western Algeria) in the context of the restoration. *Acta scientifica naturalis*, 7: 88–97. <https://doi.org/10.2478/asn-2020-0022>
- Aouadj S.A., Nasrallah Y., Hasnoui O. & Khatir H., 2020f. Impact of ecological restoration techniques on the dynamics of degraded ecosystems of the mounts of Saida: Case of the forests of Doui Thabet (West Algeria). *Acta scientifica naturalis*, 7: 98–121. <https://doi.org/10.2478/asn-2020-0023>
- Aouadj S.A., Nasrallah Y., Hasnoui O. & Khatir H., 2020g. Rare, endemic and threatened flora of the mounts of Saida (western Algeria). *Agrobiologia*, 11: 45–57.
- Barbero M., Quézel P. & Loisel R., 1990. Les apports de la phytoécologie dans l'interprétation des changements et perturbations induits par l'homme sur les écosystèmes forestiers méditerranéens. *Forêt Méditerranéenne*, 12: 194–215.
- Barry J.P., Celles J.C. & Faurel L., 1974. Carte internationale du tapis végétal. Ech : 1/1000000. Alger. 1 carte.
- Bensenane I., 2015. Assessment of anthropogenic actions of the last three decades in the steppe region of El-Gor and Sidi-Djilali (West Algeria); eco-floristic aspects. Thèse Doct. Université de Tlemcen, 120 pp.
- Bouazza M., Benabadji N., Metge G. & Loisel R., 1996. Description et aspect des sols en région semi-aride et aride au Sud de Sebdou (Oranie-Algérie). *Bulletin Institut Sciences de Rabat*, 20: 77–86.
- Braun-Blanquet J., 1952. Les groupements végétaux de la France méditerranéenne. Editions C.N.R.S., Montpellier, 297 pp.
- Bouayed I.S., Hassani F. & Ghalem S., 2019. Etude morphométrique d'une espèce menacée (fortement anthropisée): *Withania frutescens* "solanacées" dans l'Algérie Occidentale. *Algerian Journal of Environmental Science and Technology*, 5: 1121–1126.
- Ghalem S., Hassani F., Bensouna A., Khatir H. & Aouadj S.A., 2020. Floristic cortège of the genre *Lavatera* a Malvaceae for the two species: *Lavatera maritima* and *L. flava* in the region of Sabra (Tlemcen, Western of Algeria). *Ecology, environment and Conservation*, 26: S392–S397.
- Gounot M., 1969. Methods of quantitative study of vegetation. 1 Vol. Ed. March. Paris, pp. 314–341.
- Médail P. & Quézel P., 1999. Biodiversity hotspots in the Mediterranean basin: setting global conservation priorities. *Conservation Biology*, 13: 1510–1513.
- Quézel P., 1999. Large vegetation structures in the Mediterranean region: determining factors in their post-glacial development. *Geobios*, 32: 19–32.
- Quézel P. & Médail, 1995. La région circum-méditerranéenne: centre mondial majeur de biodiversité végétale. In: Actes des 6e Rencontres de l'ARPE Provence-Alpes-Côte d'Azur, Colloque scientifique international "Bio'Mes", Gap, pp. 152–160.
- Kerzabi R., 2017. The anthropogenic effect on halophyte vegetation in the semi-arid and arid environment of Orania; Vegetation dynamics in relation to edaphic factors, 123 pp.
- Raunkiaer C., 1934. The life forms of plants and statistical plant geography. Ed. Clarendon Press, Oxford, 632 pp.
- Véla E. & Benhouhou S., 2007. Evaluation d'un nouveau point chaud de biodiversité végétale dans le Bassin méditerranéen (Afrique du Nord). *C.R. Biologies*, 330: 589–605. <https://doi.org/10.1016/j.crv.2007.04.006>