The Fischer-Clifford Matrices and Character Table of the Split Extension Group $2^5:SL(5, 2)$

Rauhi I. Elkhatib

Dept. of Mathematics, Faculty of Applied Science, Thamar University, Yemen
Rauhie@yahoo.com

Abstract The group $\bar{G} = 2^5:SL(5,2)$ is a maximal subgroup of the special linear group $SL(6,2)$ of index 2016. This Group has two inertia factor groups namely, $SL(5,2)$ and $2^4:SL(4,2)$ of indices 1 and 31 respectively in $SL(5, 2)$. The aim of this paper is to construct the Fischer-Clifford matrices of $\bar{G}$, which together with the associated partial character tables of the inertia groups, are used to compute the full character table of $\bar{G}$. There are 27 Fischer-Clifford matrices with sizes between 1×1 and 3×3 .

Keywords: linear groups, group extensions, character table, Clifford theory, inertia groups, Fischer-Clifford matrix.

Mathematics Subject Classification: 20G15, 20C40.

1 Introduction

The theory of Clifford-Fischer matrices, which is based on Clifford Theory (see [1]) Was developed by B. Fischer (see [2]). Let $\bar{G} = 2^5:SL(5,2)$ be the split extension of $N = 2^5$ by $G=SL(5,2)$ where N is the vector space of dimension 5 over $GF(2)$ on which G acts naturally. The aim of this paper is to construct the character table of $\bar{G}$ by using the technique of Fischer-Clifford matrix $M(g)$ for each class representative g of G and the character tables of the inertia factor groups H_i of the inertia groups $\bar{H}_i = 2^5:H_i$. we use the properties of the Fischer-Clifford matrices discussed in ([3], [4], [5], [9], [12]) to compute entries of these matrices.

The Fischer-Clifford matrix $M(g)$ will be portioned row-wise into blocks, where each block corresponds to an inertia group $\bar{H}_i$. Now using the columns of character table of the inertia factor H_i of $\bar{H}_i$ which correspond to classes of H_i which fuse to the class $[g]$ in G and multiply these columns by the rows of the Fischer-Clifford matrix $M(g)$ that correspond to $\bar{H}_i$. Thus, we construct the portion of the character table of $\bar{G}$ which is in the block corresponding to $\bar{H}_i$ for the classes of $\bar{G}$ that come from the coset Ng . For more information about this technique see ([3], [4], [5], [9], [12]).

The character table of $\bar{G}$ will be divided row-wise into blocks where each block corresponding to an inertia group $\bar{H}_i = N:H_i$. The computations have been carried out with the aid of computer algebra system MAGMA [11] and GAP [10] and We will follow the notion of Atlas [8].

2 Theory of Fischer-Clifford Matrices

Let $\bar{G} = N : G$ be a split extension of N by G . Then for $\theta \in \text{Irr}(N)$, we define $\bar{H} = \{x \in \bar{G} : \theta^x = \theta\} = I_{\bar{G}}(\theta)$ and $H = \{x \in G : \theta^g = \theta\} = I_G(\theta)$ where $I_{\bar{G}}(\theta)$ is the stabilizer of θ in the action of $\bar{G}$ on $\text{Irr}(N)$, we have that $I_{\bar{G}}(\theta)$ is a subgroup of $\bar{G}$ and N is normal in $I_{\bar{G}}(\theta)$. Also $[\bar{G} : I_{\bar{G}}(\theta)]$ is the size of the orbit containing θ . Then it can be shown that $\bar{H} = N : H$, where $\bar{H}$ is the inertia group of θ in $\bar{G}$. The inertia factor $\bar{H}/N \cong H$ can be regarded as the inertia group of θ in the factor group $\bar{G}/N \cong G$. Define θ^g by $\theta^g(n) = \theta(gng^{-1})$ for $g \in \bar{G}, n \in N, \theta^g \in \text{Irr}(N)$. We say that θ is extendible to $\bar{H}$ if there exists $\varphi \in \text{Irr}(\bar{H})$ such that $\varphi \downarrow_N = \theta$. If θ is extendible to $\bar{H}$, then by Gallagher [7], we have $\{\alpha : \alpha \in \text{Irr}(\bar{H}), \langle \alpha \downarrow_N, \theta \rangle \neq 0\} = \{\beta\varphi : \beta \in \text{Irr}(\bar{H}/N)\}$. Let $\bar{G}$ has the property that every irreducible character of N can be extended to its inertia group. Now let $\theta_1 = 1_N, \theta_2, \dots, \theta_t$ be representatives of the orbits of $\bar{G}$ on $\text{Irr}(N)$, $\bar{H}_i = I_{\bar{G}}(\varphi_i), 1 \leq i \leq t, \varphi_i \in \text{Irr}(\bar{H}_i)$ be an extension of θ_i to $\bar{H}_i$ and $\beta \in \text{Irr}(\bar{H}_i)$ such that $N \subseteq \text{Ker}(\beta)$. Then it can be shown that

$$\text{Irr}(\bar{G}) = \bigcup_{i=1}^t \{(\beta\varphi_i)^{\bar{G}} : \beta \in \text{Irr}(\bar{H}_i), N \subseteq \text{Ker}(\beta)\} = \bigcup_{i=1}^t \{(\beta\varphi_i)^{\bar{G}} : \beta \in (\bar{H}_i/N)\}$$

Hence the irreducible characters of $\bar{G}$ will be divided into blocks, where each block corresponds to an inertia group $\bar{H}_i$. Let $\bar{H}_i$ be the inertia factor group and φ_i be an extension of θ_i to $\bar{H}_i$. Take $\theta_1 = 1_N$ as the identity character of N , then $\bar{H}_1 = \bar{G}$ and $H_1 \cong G$. Let $X(g) = \{x_1, x_2, \dots, x_{c(g)}\}$ be a set of representatives of the conjugacy classes of $\bar{G}$ from the coset $N\bar{g}$ whose images under the natural homomorphism $\bar{G} \rightarrow G$ are in $[g]$ and we take $x_1 = \bar{g}$. We define

$$R(g) = \{(i, y_k) : 1 \leq i \leq t, H_i \cap [g] \neq \emptyset, 1 \leq k \leq r\}$$

and we note that y_k runs over representatives of the conjugacy classes of elements of H_i which fuse into $[g]$ in G . Then we define the Fischer-Clifford matrix $M(g)$ by $M(g) = (a_{(i,y_k)}^j)$, where $a_{(i,y_k)}^j = \sum_l \frac{|C_{\bar{G}}(x_j)|}{|C(y_{lk})|} \varphi_i(y_{lk})$ with columns indexed by $X(g)$ and rows indexed by $R(g)$ and where $\sum_l^t$ is the summation over all l for which $y_{lk} \sim x_j$ in $\bar{G}$. Then the partial character table of $\bar{G}$ on the classes

$$\{x_1, x_2, \dots, x_{c(g)}\} \text{ is given by } \begin{bmatrix} C_1(g)M_1(g) \\ C_2(g)M_2(g) \\ \vdots \\ C_t(g)M_t(g) \end{bmatrix} \text{ where the Fischer-Clifford matrix } M(g) = \begin{bmatrix} M_1(g) \\ M_2(g) \\ \vdots \\ M_t(g) \end{bmatrix} \text{ is}$$

divided into blocks $M(g)$ with each block corresponding to an inertia group $\bar{H}_i$ and $C_i(g)$ is the partial character table of H_i consisting of the columns corresponding to the classes that fuse into $[g]$ in G . We can also observe that the number of irreducible characters of $\bar{G}$ is the sum of the number of irreducible characters of the inertia factors H_i 's. The group $\bar{G} = 2^5 : SL(5, 2)$ is a split extension with 2^5 abelian and therefore by Mackey's theorem (see [4] - Theorem 4.1.12), we have each irreducible character of 2^5 can be extended to its inertia group in $\bar{G}$. Hence by the above theoretical outline we can fully determine the character table of $\bar{G} = 2^5 : SL(5, 2)$.

3 The conjugacy classes of $2^5 : SL(5, 2)$

In this section, we will use the method of coset analysis to determine the conjugacy classes of $\bar{G} = 2^5 : SL(5, 2)$. We refer the reader to ([3], [4], [5], [9], [12]) for full details and background material regarding the method of coset analysis. Most of the information, which involved the conjugacy classes and permutation characters, were obtained by using direct computations in GAP [10] and MAGMA [11].

The special linear group $G := SL(5, 2)$ of Order = 9999360 = $2^{10} \cdot 3^2 \cdot 5 \cdot 7 \cdot 31$. We generate the group $SL(5, 2)$ by the two 5×5 matrices:

$$a = \begin{bmatrix} 0 & 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \end{bmatrix}, \quad b = \begin{bmatrix} 1 & 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Thus, by using MAGMA [11], We can find the conjugacy classes representative of $SL(5,2)$ in terms of 5×5 matrices over $GF(2)$. $SL(5,2)$ has 27 conjugacy classes and under the action of $SL(5,2)$ on 2^5 , we obtain two orbits of lengths 1 and 31. The point stabilizers for orbits of lengths 1 and 31 are the subgroups $SL(5,2)$ and $2^4:SL(4,2)$ of indices 1 and 31, respectively in $SL(5,2)$. Let $\chi(SL(5,2)|2^5)$ be the permutation character of $SL(5,2)$ on 2^5 . The values of $\chi(SL(5,2)|2^5)$ on different classes of $SL(5,2)$ determine the number of k of fixed points of each conjugacy class of $SL(5,2)$ in 2^5 . These values of the k 's will help us to calculate the conjugacy classes of $2^5:SL(5,2)$ and are listed in Table 1. Consequentially, having obtained the values of the k 's for the various classes of G , we then need to calculate the values of f_i 's corresponding to the various k 's, where f_i 's are the number of orbits Q_i 's for $1 \leq i \leq k$, that fuse together under the action of $C_G(g)$ to form one orbit Δ_j . For this purpose we used Programme A [4]. For a class representative $dg \in \bar{G}$, where $d \in 2^5$, $g \in SL(5,2)$ and $o(g) = m$, by Theorem 3.3.10 in [12] we have

$$o(dg) = \begin{cases} m & \text{if } w = 1_N \\ 2m & \text{otherwise} \end{cases}$$

To calculate the orders of the class representative $dg \in \bar{G}$, we used Programme B [4]. If $o(g) = m$ and $w = 1_N$ then $o(dg) = 2m$ otherwise if $w \neq 1_N$, then $o(dg) = 2m$. To each class of $\bar{G}$, we have attached some weight m_{ij} which will be used later in computing the Fischer matrices of the extension. These weights are computed by the formula

$$m_{ij} = [N_{\bar{G}}(N_{\bar{G}_i}):C_{\bar{G}}(g_{ij})] = |N| \frac{|C_G(g_i)|}{|C_{\bar{G}}(g_{ij})|}$$

Thus, we obtained 52 conjugacy classes for the group $2^5:SL(5,2)$ and we list these results about the conjugacy classes of $\bar{G}$ in Table 1.

Table1: Conjugacy Classes of $2^5:SL(5,2)$

$g_i \in SL(5,2)$	k_i	f_i	$m_{i,j}$	$[x]_{2^5:SL(5,2)}$	$ [x]_{2^5:SL(5,2)} $	$ C_{2^5:SL(5,2)}(x) $
$g_1 = 1a$	32	1 31	$m_{1,1} = 1$ $m_{1,2} = 31$	$g_{1,1} = 1a$ $g_{1,2} = 2a$	1 31	319979520 10321920
$g_2 = 2a$	16	1 7 8	$m_{2,1} = 2$ $m_{2,2} = 14$ $m_{2,3} = 16$	$g_{2,1} = 2b$ $g_{2,2} = 2c$ $g_{2,3} = 4a$	930 6510 7440	344064 49152 43008
$g_3 = 2b$	8	1 1 6	$m_{3,1} = 4$ $m_{3,2} = 4$ $m_{3,3} = 24$	$g_{3,1} = 2d$ $g_{3,2} = 2e$ $g_{3,3} = 4b$	26040 26040 156240	12288 12288 2048
$g_4 = 3a$	8	1 7	$m_{4,1} = 4$ $m_{4,2} = 28$	$g_{4,1} = 3a$ $g_{4,2} = 6a$	79360 555520	4032 576
$g_5 = 3b$	2	1 1	$m_{5,1} = 16$ $m_{5,2} = 16$	$g_{5,1} = 3b$ $g_{5,2} = 6b$	888832 888832	360 360
$g_6 = 4a$	8	1 3 4	$m_{6,1} = 4$ $m_{6,2} = 12$ $m_{6,3} = 16$	$g_{6,1} = 4c$ $g_{6,2} = 4d$ $g_{6,3} = 4e$	104160 312480 416640	3072 1024 768
$g_7 = 4b$	4	1 1 2	$m_{7,1} = 8$ $m_{7,2} = 8$ $m_{7,3} = 16$	$g_{7,1} = 4f$ $g_{7,2} = 4g$ $g_{7,3} = 4h$	624960 624960 1249920	512 512 256

$g_8 = 4c$	4	1 1 2	$m_{8,1} = 8$ $m_{8,2} = 8$ $m_{8,3} = 16$	$g_{8,1} = 4l$ $g_{8,2} = 4m$ $g_{8,3} = 8a$	2499840 2499840 4999680	128 128 64
$g_9 = 5a$	2	1 1	$m_{9,1} = 16$ $m_{9,2} = 16$	$g_{9,1} = 5a$ $g_{9,2} = 10a$	10665984 10665984	30 30
$g_{10} = 6a$	4	1 1 2	$m_{10,1} = 8$ $m_{10,2} = 8$ $m_{10,3} = 16$	$g_{10,1} = 6c$ $g_{10,2} = 6d$ $g_{10,3} = 12a$	3333120 3333120 6666240	96 96 48
$g_{11} = 6b$	2	1 1	$m_{11,1} = 16$ $m_{11,2} = 16$	$g_{11,1} = 6e$ $g_{11,2} = 6f$	13332480 13332480	24 24
$g_{12} = 7a$	4	1 3	$m_{12,1} = 8$ $m_{12,2} = 24$	$g_{12,1} = 7a$ $g_{12,2} = 14a$	1904640 5713920	168 56
$g_{13} = 7b$	4	1 3	$m_{13,1} = 8$ $m_{13,2} = 24$	$g_{13,1} = 7b$ $g_{13,2} = 14b$	1904640 5713920	168 56
$g_{14} = 8a$	2	1 1	$m_{14,1} = 16$ $m_{14,2} = 16$	$g_{14,1} = 8b$ $g_{14,2} = 8c$	9999360 9999360	32 32
$g_{15} = 12a$	2	1 1	$m_{15,1} = 16$ $m_{15,2} = 16$	$g_{15,1} = 12b$ $g_{15,2} = 12c$	13332480 13332480	24 24
$g_{16} = 14a$	2	1 1	$m_{16,1} = 16$ $m_{16,2} = 16$	$g_{16,1} = 14c$ $g_{16,2} = 28a$	11427840 11427840	28 28
$g_{17} = 14b$	2	1 1	$m_{17,1} = 16$ $m_{17,2} = 16$	$g_{17,1} = 14d$ $g_{17,2} = 28b$	11427840 11427840	28 28
$g_{18} = 15a$	2	1 1	$m_{18,1} = 16$ $m_{18,2} = 16$	$g_{18,1} = 15a$ $g_{18,2} = 30a$	10665984 10665984	30 30
$g_{19} = 15b$	2	1 1	$m_{19,1} = 16$ $m_{19,2} = 16$	$g_{19,1} = 15b$ $g_{19,2} = 30b$	10665984 10665984	30 30
$g_{20} = 21a$	1	1	$m_{20,1} = 32$	$g_{20,1} = 21a$	15237120	21
$g_{21} = 21b$	1	1	$m_{21,1} = 32$	$g_{21,1} = 21b$	15237120	21
$g_{22} = 31a$	1	1	$m_{22,1} = 32$	$g_{22,1} = 31a$	10321920	31
$g_{23} = 31b$	1	1	$m_{23,1} = 32$	$g_{23,1} = 31b$	10321920	31
$g_{24} = 31c$	1	1	$m_{24,1} = 32$	$g_{24,1} = 31c$	10321920	31
$g_{25} = 31d$	1	1	$m_{25,1} = 32$	$g_{25,1} = 31d$	10321920	31
$g_{26} = 31e$	1	1	$m_{26,1} = 32$	$g_{26,1} = 31e$	10321920	31
$g_{27} = 31f$	1	1	$m_{27,1} = 32$	$g_{27,1} = 31f$	10321920	31

4 The Inertia Factor Group of $\bar{G} = 2^5:SL(5, 2)$

The action of G on N produce two orbits of lengths 1 and 31. Hence by Brauer's theorem (see Theorem 5.1.4 in [12]) G acts on $\text{Irr}(N)$ with the same number of orbits. The lengths of the two orbits will be $1, r$ where $1 + r = 32$, with corresponding point stabilizers H_1 and H_2 as subgroups of G such that $[G:H_1] = 1$ and $[G:H_2] = 31$. Considering the indices of these subgroups in G and investigating the maximal and submaximal subgroups of G , we have the Group $SL(5, 2)$ acting on $\text{Irr}(2^5)$ produce two inertia factor groups:

- (1) $H_1 = SL(5, 2)$ of index equal 1,
- (2) $H_2 = 2^4:SL(4, 2)$ of index equal 31.

Using GAP [10], we can generate the group $2^4:SL(4, 2)$ in terms of 5×5 matrices over $GF(2)$ as follows:

$$a = \begin{bmatrix} 1 & 0 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, b = \begin{bmatrix} 1 & 0 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}, c = \begin{bmatrix} 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \\ 1 & 1 & 1 & 0 & 0 \\ 0 & 1 & 1 & 0 & 0 \end{bmatrix}$$

Using these generators in MAGMA [11], we obtained the conjugacy classes and the character tables for these groups.

5 The Fusion of Inertia Factor Group $H = 2^4:SL(4, 2)$ into the group $G = SL(5, 2)$

We can complete the fusion maps by using matrix conjugation in $SL(5, 2)$, This fusion map is listed in table2:

Table 2: The Fusion $H = 2^4:SL(4, 2)$ into $G = SL(5, 2)$

$[g]_H \Rightarrow$	$[g]_G$	$[g]_H \Rightarrow$	$[g]_G$	$[g]_H \Rightarrow$	$[g]_G$
1a	1a	2a	2a	2b	2a
2c	2b	2d	2b	3a	3b
3b	3a	4a	4a	4b	4b
4c	4a	4d	4b	4e	4c
4f	4c	5a	5a	6a	6a
6b	6b	6c	6a	7a	7a
7b	7b	8a	8a	12a	12a
14a	14b	14b	14a	15a	15a
15b	15b				

6 The Fischer-Clifford Matrices of $\bar{G} = 2^5:SL(5, 2)$

For each conjugacy class $[g]$ of G with representative $g \in G$, we construct the corresponding Fischer-Clifford matrix $M(g)$ of $\bar{G} = 2^5:SL(5, 2)$. We use the properties of the Fischer-Clifford matrices (see ([3], [4], [5], [9], [12])) together with fusions of H_2 into H_1 (Table 2) to compute the entries of these matrices and to construct an algebraic system of linear and non-linear equations with the help of the symbolic mathematical package Maxima [6], we can solve these system of equations and compute all the Fischer matrices of $\bar{G}$. The Fischer-Clifford matrix will be partitioned row-wise into blocks, where each block corresponding to an inertia group $\bar{H}_i$. We list the Fischer-Clifford matrices of $\bar{G}$ in Table 3.

Table 3: The Fischer-Clifford Matrices of $2^5:SL(5, 2)$

$M(g)$	$M(g)$
$M(1a) = \begin{bmatrix} 1 & 1 \\ 31 & -1 \end{bmatrix}$	$M(2a) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 14 & -2 & 0 \end{bmatrix}$
$M(2b) = \begin{bmatrix} 1 & 1 & 1 \\ 3 & 3 & -1 \\ 4 & -4 & 0 \end{bmatrix}$	$M(3a) = \begin{bmatrix} 1 & 1 \\ 7 & -1 \end{bmatrix}$
$M(3b) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(4a) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 6 & -2 & 0 \end{bmatrix}$
$M(4b) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 2 & -2 & 0 \end{bmatrix}$	$M(4c) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 2 & -2 & 0 \end{bmatrix}$

$M(5a) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(6a) = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 1 & -1 \\ 2 & -2 & 0 \end{bmatrix}$
$M(6b) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(7a) = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$
$M(7b) = \begin{bmatrix} 1 & 1 \\ 3 & -1 \end{bmatrix}$	$M(8a) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
$M(12a) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(14a) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
$M(14b) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(15a) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
$M(15b) = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$M(21a) = [1]$
$M(21b) = [1]$	$M(31a) = [1]$
$M(31b) = [1]$	$M(31c) = [1]$
$M(31d) = [1]$	$M(31e) = [1]$
$M(31f) = [1]$	

7 The Character Table of $\bar{G} = 2^5:SL(5, 2)$

Now, we have:

- (1) The conjugacy classes of $\bar{G} = 2^5:SL(5,2)$ (Table 1);
- (2) The character tables of all the inertia factors (by using MAGMA [11]);
- (3) The fusions of conjugacy classes of the inertia factors into classes of $SL(5,2)$ (Table 2) ;
- (4) The Fischer matrices of classes of $\bar{G} = 2^5:SL(5,2)$ (Table 3);

Thus, we can construct the character table of the $2^5:SL(5,2)$

Table 4. The Character Table of $\bar{G} = 2^5:SL(5, 2)$

Class	1a	2a	2b	2c	2d	2e	3a	3b	4a	4b
Order	1	2	2	2	2	2	3	3	4	4
Size	1	31	930	6510	26040	26040	79360	888832	7440	156240
X1	1	1	1	1	1	1	1	1	1	1
X2	30	30	14	14	6	6	6	0	14	6
X3	124	124	28,	28	12	12	1	4	28	12
X4	155	155	27	27	-5	-5	8	5	27	-5
X5	217	217	-7	-7	9	9	7	4	-7	9
X6	280	280	56	56	8	8	7	-5	56	8
X7	315	315	-21	-21	3	3	0	0	-21	3
X8	315	315	-21	-21	3	3	0	0	-21	3
X9	315	315	-21	-21	3	3	0	0	21	3
X10	315	315	-21	-21	3	3	0	0	-21	3
X11	315	315	-21	-21	3	3	0	0	-21	3
X12	315	315	-21	-21	3	3	0	0	-21	3
X13	465	465	17	17	-15	-15	3	0	17	-15
X14	465	465	-31	-31	9	9	3	0	-31	9
X15	465	465	17	17	-15	-15	3	0	17	-15
X16	465	465	-31	-31	9	9	3	0	-31	9
X17	496	496	48	48	16	16	-8	1	48	16
X18	651	651	-21	-21	-5	-5	0	6	-21	-5
X19	651	651	-21	-21	-5	-5	0	-3	-21	-5
X20	651	651	-21	-21	-5	-5	0	-3	-21	-5
X21	868	868	-28	-28	4	4	7	1	-28	4
X22	930	930	50	50	-6	-6	6	0	50	-6
X23	930	930	-14	-14	-6	-6	-3	0	-14	-6
X24	930	930	-14	-14	-6	-6	-3	0	-14	-6
X25	960	960	64	64	0	0	-6	0	64	0

X26	1024	1024	0,	0	0	0	-8	4	0	0
X27	1240	1240	-8	-8	8	8	1	-5	-8	8
X28	31	-1	15	-1	7	-1	7,	1	-1	-1
X29	217	-7	-7	9	9	-15	7	4	-7	1
X30	434	-14	98	2	26	10	14	-1	-14	-6
X31	465	-15	97	-15	9	-15	21	0	1	1
X32	620	-20	76	12	28	-4,	-7	5	-20	-4
X33	651	-21	-21	27	-5	-13	0	6	-21	3
X34	651	-21	-21	27	-5	-13	0	-3	-21	3
X35	651	-21	-21	27	-5	-13	0	-3	-21	3
X36	868	-28	-28	36	4	-28	7	1	-28	4
X37	1085	-35	77	29	-11	29	14	5	-35	-3
X38	1395	-45	-45	3	3	27	0	0	3	-5
X39	1395	-45	3	51	-21	3	0	0	-45	3
X40	1395	-45	-45	3	3	27	0	0	3	-5
X41	1395	-45	3	51	-21	3	0	0	-45	3
X42	1736	-56	168	40	24	24	-7	-4	-56	-8
X43	1984	-64	64	64	0	0	-14	4	-64	0
X44	2170	-70	42	74	2	-14	7	-5	-70	2
X45	2790	-90	246	-42	30	-18	0	0	6	-2
X46	3255	-105	231	-41	-33	-9	21	0	7	7
X47	3255	-105	7	-9	15,	39	21	0	7	-9
X48	3255	-105	-105	7	39	-33	21	0	7	-1
X49	3720	-120	104	-24	24	24	-21	0	8	-8
X50	6510	-210	126	-34	6	-42	-21	0	14	6
X51	9765	-315	-315	21	21	-3	0	0	21	-3
X52	9765	-315	21	-27	-51	21	0	0	21	5

Class	4c	4d	4e	4f	4g	4h	4l	4m	5a
Order	4	4	4	4	4	4	4	4	5
Size	104160	312480	416640	624960	624960	1249920	2499840	2499840	10665984
X1	1	1	1	1	1	1	1	1	1
X2	6	6	6	2	2	2	2	2	0
X3	4	4	4	4	4	4	0	0	-1
X4	3	3	3	-5	-5	-5	-1	-1	0
X5	-7	-7	-7	1	1	1	1	1	2
X6	8	8	8	0	0	0	0	0	0
X7	3	3	3	-1	-1	-1	-1	-1	0
X8	3	3	3	-1	-1	-1	-1	-1	0
X9	3	3	3	-1	-1	-1	-1	-1	0
X10	3	3	3	-1	-1	-1	-1	-1	0
X11	3	3	3	-1	-1	-1	-1	-1	0
X12	3	3	3	-1	-1	-1	-1	-1	0
X13	1	1	1	1	1	1	1	1	0
X14	1	1	1	-3	-3	-3	1	1	0
X15	1	1	1	1	1	1	1	1	0
X16	1	1	1	-3	-3	-3	1	1	0
X17	0	0	0	0	0	0	0	0	1
X18	3	3	3	3	3	3	-1	-1	1
X19	3	3	3	3	3	3	-1	-1	1
X20	3	3	3	3	3	3	-1	-1	1
X21	-4	-4	-4	4	4	4	0	0	-2
X22	-6	-6	-6	-2	-2	-2	-2	-2	0
X23	2	2	2	-2	-2	-2	2	2	0
X24	2	2	2	-2	-2	-2	2	2	0
X25	0	0	0	0	0	0	0	0	0
X26	0	0	0	0	0	0	0	0	-1
X27	-8	-8	-8	0	0	0	0	0	0
X28	7	-1	-1	3	-1	-1	3	-1	1
X29	-7	1	1	1	5	-3	1	-3	2
X30	18	2	-6	6	-2	-2	2	2	-1

X31	17	-7	1	-3	1	1	1	-3	0
X32	4	4	-4	4	4	-4	0	0	0
X33	3	-5	3	3	-1	-1	-1	3	1
X34	3	-5	3	3	-1	-1	-1	3	1
X35	3	-5	3	3	-1	-1	-1	3	1
X36	-4	-4	4	4	4	-4	0	0	-2
X37	-3	5	-3	-7	-3	5	-3	1	0
X38	3	-5	3	-5	7	-1	3	-1	0
X39	3	-5	3	-1	-5	3	3	-1	0
X40	3	-5	3	-5	7	-1	3	-1	0
X41	3	-5	3	-1	-5	3	3	1	0
X42	8	8	-8	0	0	0	0	0	1
X43	0	0	0	0	0	0	0	0	-1
X44	-14	2	2	-2	6	-2	-2	-2	0
X45	6	-10	6	2	-6	2	2	2	0
X46	7	-1	-1	-5	7	-1	-1	3	0
X47	-25	-1	7	3	-1	-1	-1	3	0
X48	-17	7	-1	-1	-5	3	3	-1	0
X49	-8	-8	8	0	0	0	0	0	0
X50	-10	6	-2	-6	2	2	2	2	0
X51	21	-3	-3	-3	1	1	-3	1	0
X52	-3	5	-3	9	-3	-3	1	-3	0

Class	6a	6b	6c	6d	6e	6f	8a
Order	6	6	6	6	6	6	8
Size	555520	888832	3333120	3333120	13332480	13332480	4999680
X1	1	1	1	1	1	1	1
X2	6	0	2	2	0	0	2
X3	1	4	1	1	0	0	0
X4	8	5	0	0	1	1	-1
X5	7	4	-1	-1	0	0	1
X6	7	-5	-1	-1	-1	-1	0
X7	0	0	0	0	0	0	-1
X8	0	0	0	0	0	0	-1
X9	0	0	0	0	0	0	-1
X10	0	0	0	0	0	0	-1
X11	0	0	0	0	0	0	-1
X12	0	0	0	0	0	0	-1
X13	3	0	-1	-1	0	0	1
X14	3	0	-1	-1	0	0	1
X15	3	0	-1	-1	0	0	1
X16	3	0	-1	-1	0	0	1
X17	-8	1	0	0	1	1	0
X18	0	6	0	0	-2	-2	-1
X19	0	-3	0	0	1	1	-1
X20	0	-3	0	0	1	1	-1
X21	7	1	-1	-1	1	1	0
X22	6	0	2	2	0	0	-2
X23	-3	0	1,	1,	0	0	2
X24	-3	0	1	1	0	0	2
X25	-6	0	-2	-2	0	0	0
X26	-8	4	0	0	0	0	0
X27	1	-5	1	1	-1	-1	0
X28	-1	-1	3	-1	1	-1	-1
X29	-1	-4	1	1	-1	1	1
X30	-2	1	0	4	0	0	-2
X31	-3	0	-1	-1	1	-1	1
X32	1	-5	1	-3	1	-1	0
X33	0	-6	-4	4	0	0	-1
X34	0	3	2	-2	0	0	-1

X35	0	3	2	-2	0	0	-1
X36	-1	-1	3	-1	-1	1	0
X37	-2	-5	4	0	0	0	1
X38	0	0	0	0	0	0	-1
X39	0	0	0	0	0	0	-1
X40	0	0	0	0	0	0	-1
X41	0	0	0	0	0	0	-1
X42	1	4	-1	-1	-1	1	0
X43	2	-4	-2	-2	0	0	0
X44	-1	5	-1	3	1	-1	2
X45	0	0	0	0	0	0	-2
X46	-3	0	-1	-1	-1	1	-1
X47	-3	0	-1	-1	1	-1	-1
X48	-3	0	-1	-1	-1	1	-1
X49	3	0	1	1	-1	1	0
X50	3	0	1	1	1	-1	-2
X51	0	0	0	0	0	0	1,
X52	0	0	0	0	0	0	1

Class	7a	7b	8b	8c	10a	12a	12b
Order	7	7	8	8	10	12	12
Size	1904640	1904640	9999360	9999360	10665984	6666240	13332480
X1	1	1	1	1	1	1	1
X2	2	2	0	0	0	2	0
X3	-2	-2	0	0	-1	1	1
X4	1	1	-1	-1	0	0	0
X5	0	0,	1	1	2	-1	-1
X6	0	0	0	0	0	-1	-1
X7	0	0	1	1	0	0	0
X8	0	0	1	1	0	0	0
X9	0	0	1	1	0	0	0
X10	0	0	1	1	0	0	0
X11	0	0	1	1	0	0	0
X12	0	0	1	1	0	0	0
X13	z1	z1#3	1	1	0	-1	1
X14	z1#3	z1	-1	-1	0	-1	1
X15	z1#3	z1	1	1	0	-1	1
X16	z1	z1#3	-1	-1	0	-1	1
X17	-1	-1	0	0	1	0	0
X18	0	0	-1	-1	1	0	0
X19	0	0	-1	-1	1	0	0
X20	0	0	-1	-1	1	0	0
X21	0	0	0	0	-2	-1	-1
X22	-1	-1	0	0	0	2	0
X23	z2	z2#3	0	0	0	1,	-1
X24	z2#3	z2	0	0	0	1	-1
X25	1	1	0	0	0	-2	0
X26	2	2	0	0	-1	0	0
X27	1	1	0	0	0	1	1
X28	3	3	1	-1	-1	-1	1
X29	0	0	1	-1	-2	-1	-1
X30	0	0	0	0	1	-2	0
X31	3	3	-1	1	0	1	-1
X32	-3	-3	0	0	0	1	1
X33	0	0	-1	1	-1	0	0
X34	0	0	-1	1	-1	0	0
X35	0	0	-1	1	-1	0	0
X36	0	0	0	0	2	-1	-1
X37	0	0	-1	1	0	-2	0
X38	3*z1	3*(z1#3)	1	-1	0	0	0
X39	3*(z1#3)	3*z1	-1	1	0	0	0

X40	$3^*(z1\#3)$	3^*z1	1	-1	0	0	0
X41	3^*z1	$3^*(z1\#3)$	-1	1	0	0	0
X42	0	0	0	0	-1	1	-1
X43	3	3	0	0	1	2	0
X44	0	0	0	0	0	-1	1
X45	-3	-3	0	0	0	0	0
X46	0	0	1	-1	0	1	1
X47	0	0	1	-1	0	1	-1
X48	0	0	-1	1	0	1	1
X49	3	3	0	0	0	-1	1
X50	0	0	0	0	0	-1	-1
X51	0	0	1	-1	0	0	0
X52	0	0	-1	1	0	0	0

Class	12c	14a	14b	14c	14d	15a	15b
Order	12	14	14	14	14	15	15
Size	13332480	5713920	5713920	11427840	11427840	10665984	10665984
X1	1	1	1	1	1	1	1
X2	0	2	2	0	0	0	0
X3	1	-2	-2	0	0	-1	-1
X4	0	1	1	-1	-1	0	0
X5	-1	0	0,	0	0	-1	-1
X6	-1	0	0	0	0	0	0
X7	0	0	0	0	0	0	0
X8	0	0	0	0	0	0	0
X9	0	0	0	0	0	0	0
X10	0	0	0	0	0	0	0
X11	0	0	0	0	0	0	0
X12	0	0	0	0	0	0	0
X13	1	$z1$	$z1\#3$	$z1$	$z1\#3$	0	0
X14	1	$z1\#3$	$z1$	$-z1\#3$	$-z1$	0	0
X15	1	$z1\#3$	$z1$	$z1\#3$	$z1$	0	0
X16	1	$z1$	$z1\#3$	$-z1$	$-z1\#3$	0	0
X17	0	-1	-1	-1	-1	1	1
X18	0	0	0	0	0	1	1
X19	0	0	0	0	0	$z3$	$z3\#7$
X20	0	0	0	0	0	$z3\#7$	$z3$
X21	-1	0	0	0	0	1	1
X22	0	-1	-1	1	1	0	0
X23	-1	$z2$	$z2\#3$	0	0	0	0
X24	-1	$z2\#3$	$z2$	0	0	0	0
X25	0	1	1	1	1	0	0
X26	0	2	2	0	0	-1	-1
X27	1	1	1	-1	-1	0	0
X28	-1	-1	-1	1	1	1	1
X29	1	0	0	0	0	-1	-1
X30	0	0	0	0	0	-1	-1
X31	1	-1	-1	-1	-1	0	0
X32	-1	1	1	-1	-1	0	0
X33	0	0	0	0	0	1	1
X34	0	0	0	0	0	$z2$	$z2\#7$
X35	0	0	0	0	0	$z2\#7$	$z2$
X36	1	0	0	0	0	1	1
X37	0	0	0	0	0	0	0
X38	0	$-z1$	$-(z1\#3)$	$z1$	$z1\#3$	0	0
X39	0	$-(z1\#3)$	$-z1$	$-z1\#3$	$-z1$	0	0
X40	0	$-(z1\#3)$	$-z1$	$z1\#3$	$z1$	0	0
X41	0	$-z1$	$-(z1\#3)$	$-z1$	$-z1\#3$	0	0
X42	1	0	0	0	0	1	1
X43	0	-1	-1	1	1	-1	-1
X44	-1	0	0	0	0	0	0

X45	0	1	1	1	1	0	0
X46	-1	0	0	0	0	0	0
X47	1	0	0	0	0	0	0
X48	-1	0	0	0	0	0	0
X49	-1	-1	-1	-1	-1	0	0
X50	1	0	0	0	0	0	0
X51	0	0	0	0	0	0	0
X52	0	0	0	0	0	0	0

Class	21a	21b	28a	28b	30a	30b
Order	21	21	28	28	30	30
Size	15237120	15237120	11427840	11427840	10665984	10665984
X1	1	1	1	1	1	1
X2	-1	-1	0	0	0	0
X3	1	1	0	0	-1	-1
X4	1	1	-1	-1	0	0
X5	0	0	0	0	-1	-1
X6	0	0	0	0	0	0
X7	0	0	0	0	0	0
X8	0	0	0	0	0	0
X9	0	0	0	0	0	0
X10	0	0	0	0	0	0
X11	0	0	0	0	0	0
X12	0	0	0	0	0	0
X13	Z1	Z1#3	z1	z1#3	0	0
X14	Z1#3	Z1	-z1#3	-z1	0	0
X15	Z1#3	Z1	z1#3	z1	0	0
X16	Z1	Z1#3	-z1	-z1#3	0	0
X17	-1	-1	-1	-1	1	1
X18	0	0	0	0	1	1
X19	0	0	0	0	z3	z3#7
X20	0	0	0	0	z3#7	z3
X21	0	0	0	0	1	1
X22	-1	-1	1	1	0	0
X23	-z1	-z1#3	0	0	0	0
X24	-z1#3	-z1	0	0	0	0
X25	1	1	1	1	0	0
X26	-1	-1	0	0	-1	-1
X27	1	1	-1	-1	0	0
X28	0	0	-1	-1	-1	-1
X29	0	0	0	0	1	1
X30	0	0	0	0	1	1
X31	0	0	1	1	0	0
X32	0	0	1	1	0	0
X33	0	0	0	0	-1	-1
X34	0	0	0	0	-z2	-(z2#7)
X35	0	0	0	0	-(z2#7)	-z2
X36	0	0	0	0	-1	-1
X37	0	0	0	0	0	0
X38	0	0	-z1	-(z1#3)	0	0
X39	0	0	-(-z1#3)	z1	0	0
X40	0	0	-(z1#3)	-z1	0	0
X41	0	0	z1	-(-z1#3)	0	0
X42	0	0	0	0	-1	-1
X43	0	0	-1	-1	1	1
X44	0	0	0	0	0	0
X45	0	0	-1	-1	0	0
X46	0	0	0	0	0	0
X47	0	0	0	0	0	0
X48	0	0	0	0	0	0
X49	0	0	1	1	0	0
X50	0	0	0	0	0	0

X51	0	0	0	0	0	0
X52	0	0	0	0	0	0
Class	31a	31b	31c	31d	31e	31f
Order	31	31	31	31	31	31
Size	10321920	10321920	10321920	10321920	10321920	10321920
X1	1	1	1	1	1	1
X2	-1	-1	-1	-1	-1	-1
X3	0	0	0	0	0	0
X4	0	0	0	0	0	0
X5	0	0	0	0	0	0
X6	1	1	1	1	1	1
X7	Z4	Z4#15	Z4#3	Z4#7	Z4#5	Z4#11
X8	Z4#3	Z4#7	Z4#5	Z4#11	Z4#15	Z4
X9	Z4#5	Z4#11	Z4#15	Z4	Z4#7	Z4#3
X10	Z4#11	Z4#5	Z4	Z4#15	Z4#3	Z4#7
X11	Z4#15	Z4	Z4#7	Z4#3	Z4#11	Z4#5
X12	Z4#7	Z4#3	Z4#11	Z4#5	Z4	Z4#15
X13	0	0	0	0	0	0
X14	0	0	0	0	0	0
X15	0	0	0	0	0	0
X16	0	0	0	0	0	0
X17	0	0	0	0	0	0
X18	0	0	0	0	0	0
X19	0	0	0	0	0	0
X20	0	0	0	0	0	0
X21	0	0	0	0	0	0
X22	0	0	0	0	0	0
X23	0	0	0	0	0	0
X24	0	0	0	0	0	0
X25	-1	-1	-1	-1	-1	-1
X26	1	1	1	1	1	1
X27	0	0	0	0	0	0
X28	0	0	0	0	0	0
X29	0	0	0	0	0	0
X30	0	0	0	0	0	0
X31	0	0	0	0	0	0
X32	0	0	0	0	0	0
X33	0	0	0	0	0	0
X34	0	0	0	0	0	0
X35	0	0	0	0	0	0
X36	0	0	0	0	0	0
X37	0	0	0	0	0	0
X38	0	0	0	0	0	0
X39	0	0	0	0	0	0
X40	0	0	0	0	0	0
X41	0	0	0	0	0	0
X42	0	0	0	0	0	0
X43	0	0	0	0	0	0
X44	0	0	0	0	0	0
X45	0	0	0	0	0	0
X46	0	0	0	0	0	0
X47	0	0	0	0	0	0
X48	0	0	0	0	0	0
X49	0	0	0	0	0	0
X50	0	0	0	0	0	0
X51	0	0	0	0	0	0
X52	0	0	0	0	0	0

Explanation of Character Value Symbols:

denotes algebraic conjugation, #k indicates replacing the root of unity w by w^k .

Z1 = (CyclotomicField (7: Sparse := true)) ! [RationalField() | 0, 1, 1, 0,1, 0];

Z2 = (CyclotomicField (7: Sparse := true)) ! [RationalField() | 0, 2, 2, 0,2, 0];

Z3 = (CyclotomicField (15: Sparse := true)) ! [RationalField() | -1, -1, 0,0, -1, -2, -1, -2];

Z4= (CyclotomicField (31: Sparse := true)) ! [RationalField() | -1, -1, -1,-1, -1, -1, -1, -1, -1, -1, -1, -1, -1, -1, 0, -1, -1, -1, -1, -1, -1, -1,0, -1, -1, -1, 0, -1, 0].

References:

- [1] A. H. Clifford, Representations induced in an invariant subgroup, Ann. of Math. 38 (1937), 533 - 550.
- [2] B. Fischer, Clifford matrices, Representation theory of finite groups and finite-dimensional Lie algebras (eds G. O. Michler and C. M. Ringel; Birkhäuser, Basel, 1991), 1 - 16.
- [3] B. M. Basheer, Clifford-Fischer Theory Applied to Certain Groups Associated with Symplectic, Unitary and Thompson Groups (PhD Thesis, University of KwaZulu-Natal, Pietermaritzburg 2012.
- [4] F. Ali, Fischer-Clifford Theory for Split and Non-Split Group Extensions, PhD Thesis, University of Natal, Pietermaritzburg, 2001.
- [5] J. Moori, On the Groups G^+ and G of the form 210:M22 and 210:M22, PhD Thesis, University of Birmingham, 1975.
- [6] Maxima, A Computer Algebra System. Version 5.18.1; 2009, (<http://maxima.sourceforge.net>).
- [7] P. X. Gallagher, Group characters and normal Hall subgroups, Nagoya Math. J. **21** (1962), 223 - 230.
- [8] R. A. Wilson et al., Atlas of finite group representations, (<http://brauer.maths.qmul.ac.uk/Atlas/v3/>)
- [9] T. T. Seretlo, Fischer-Clifford Matrices and Character Tables of Certain Groups Associated with Simple Groups $O^+_8(2)$, HS and Ly , PhD Thesis, University of KwaZulu-Natal, Pietermaritzburg, 2012.
- [10] The GAP Group, GAP – Groups, Algorithms, and Programming, Version 4.8.6; 2016. (<http://www.gap-system.org>).
- [11] W. Bosma and J. J. Cannon, Handbook of Magma Functions, Department of Mathematics, University of Sydney, November 1994.
- [12] Z. E. Mpono, Fischer Clifford Theory and Character Tables of Group Extensions, PhD Thesis, University of Natal, Pietermaritzburg, 1998.