Explicit Expression for a First Integral for a Class of Two-dimensional Differential System

Rachid Boukoucha ^{*1}, Mouna Yahiaoui ²

^{1,2} Lab. de Mathématiques Appliquées, Université de Bejaia, 06000 Bejaia, Algérie.

¹rachid.boukecha@yahoo.fr, ²mouna.yahiaoui89@gmail.com

Abstract. In this paper we are interested in studying the existence of a first integral and to the curves which are formed by the trajectories of the two-dimensional differential systems of the form

$$\begin{cases} x' = P(x, y) + x \left(\lambda x \exp \left(\frac{M(x, y)}{N(x, y)} \right) + \beta y \exp \left(\frac{R(x, y)}{S(x, y)} \right) \right), \\ y' = Q(x, y) + y \left(\lambda x \exp \left(\frac{M(x, y)}{N(x, y)} \right) + \beta y \exp \left(\frac{R(x, y)}{S(x, y)} \right) \right), \end{cases}$$

where $P(x, y)$, $Q(x, y)$, $M(x, y)$, $N(x, y)$, $R(x, y)$, $S(x, y)$ are homogeneous polynomials of degree a , a , b , b , c , c respectively and $\lambda, \beta \in \mathbb{R}$. Concrete examples exhibiting the applicability of our result are introduced.

Keywords: *Differential system, First integral, Curves.*

2010 MSC No: 34C05, 34C07, 37C27, 37K10.

1 Introduction

We consider two-dimensional autonomous systems of differential equations of the form

$$\begin{cases} x' = \frac{dx}{dt} = F(x(t), y(t)), \\ y' = \frac{dy}{dt} = G(x(t), y(t)), \end{cases} \quad (1)$$

where $F(x, y)$ and $G(x, y)$ are real functions. In the qualitative theory of planar dynamical systems [1, 9, 10, 16], one of the most important topics is related to the second part of the unsolved Hilbert 16th problem [15]. There is a huge literature about limit cycles, most of them deal essentially with their detection, their number and their stability and rare are papers concerned by giving them explicitly [2, 4, 7, 13]. There exist three main open problems in the qualitative theory of real planar differential systems, the distinction between a centre and a focus, the determination of the number of limit cycles and their distribution, and the determination of its integrability. The importance for searching first integrals of a given system was already noted by Poincaré in his discussion on a method to obtain polynomial or rational first integrals. One of the classical tools in the classification of all trajectories of a dynamical system is to find first integrals. For more details about first integral see for instance [3, 5, 8, 11, 12, 14], see the

^{*}Corresponding author. Rachid Boukoucha ¹ rachid.boukecha@yahoo.fr

references quoted in those articles. We recall that in the phase plane, a limit cycle of system (1) is an isolated periodic orbit in the set of all periodic orbits of system (1)

System (1) is integrable on an open set Ω of $\mathbb{R}^2$ if there exists a non constant C^1 function $H : \Omega \rightarrow \mathbb{R}$, called a first integral of the system on Ω , which is constant on the trajectories of the system (1) contained in Ω , i.e. if

$$\frac{dH(x, y)}{dt} = \frac{\partial H(x, y)}{\partial x} F(x, y) + \frac{\partial H(x, y)}{\partial y} G(x, y) \equiv 0 \quad \text{in the points of } \Omega.$$

Moreover, $H = h$ is the general solution of this equation, where h is an arbitrary constant. It is well known that for differential systems defined on the plane $\mathbb{R}^2$ the existence of a first integral determines their phase portrait [6].

In this paper we are interested in studying the existence of a first integral and to the curves which are formed by the trajectories of the two-dimensional differential systems of the form

$$\begin{cases} x' = P(x, y) + x \left(\lambda x \exp \left(\frac{M(x, y)}{N(x, y)} \right) + \beta y \exp \left(\frac{R(x, y)}{S(x, y)} \right) \right), \\ y' = Q(x, y) + y \left(\lambda x \exp \left(\frac{M(x, y)}{N(x, y)} \right) + \beta y \exp \left(\frac{R(x, y)}{S(x, y)} \right) \right), \end{cases} \quad (2)$$

where $P(x, y)$, $Q(x, y)$, $M(x, y)$, $N(x, y)$, $R(x, y)$, $S(x, y)$ are homogeneous polynomials of degree a , a , b , b , c , c respectively and $\lambda, \beta \in \mathbb{R}$.

We define the trigonometric functions

$$\begin{aligned} f_1(\theta) &= \lambda (\cos \theta) \exp \left(\frac{M(\cos \theta, \sin \theta)}{N(\cos \theta, \sin \theta)} \right) + \beta (\sin \theta) \exp \left(\frac{R(\cos \theta, \sin \theta)}{S(\cos \theta, \sin \theta)} \right), \\ f_2(\theta) &= P(\cos \theta, \sin \theta) \cos \theta + Q(\cos \theta, \sin \theta) \sin \theta, \\ f_3(\theta) &= (\cos \theta) Q(\cos \theta, \sin \theta) - (\sin \theta) P(\cos \theta, \sin \theta). \end{aligned}$$

2 Main result

Our main result on the integrability of differential system (2) is the following :

Theorem 1. *Consider a differential system (2), then the following statements hold.*

(1) *If $f_3(\theta) N(\cos \theta, \sin \theta) S(\cos \theta, \sin \theta) \neq 0$ and $a \neq 2$, then system (2) has the first integral*

$$\begin{aligned} H(x, y) &= (x^2 + y^2)^{\frac{a-2}{2}} \exp \left((2-a) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega \right) - \\ &\quad (a-2) \int_0^{\arctan \frac{y}{x}} \exp \left((2-a) \int_0^w A(\omega) d\omega \right) B(w) dw, \end{aligned}$$

where $A(\theta) = \frac{f_2(\theta)}{f_3(\theta)}$, $B(\theta) = \frac{f_1(\theta)}{f_3(\theta)}$, and the curves which are formed by the trajectories of the differential system (2), in cartesian coordinates are written as

$$x^2 + y^2 = \left(\begin{array}{c} h \exp \left((a-2) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega \right) + \\ (a-2) \exp \left((a-2) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega \right) \\ \int_0^{\arctan \frac{y}{x}} \exp \left((2-a) \int_0^w A(\omega) d\omega \right) B(w) dw \end{array} \right)^{\frac{2}{a-2}},$$

where $h \in \mathbb{R}$.

(2) *If $f_3(\theta) N(\cos \theta, \sin \theta) S(\cos \theta, \sin \theta) \neq 0$ and $a = 2$, then system (2) has the first integral*

$$H(x, y) = (x^2 + y^2)^{\frac{1}{2}} \exp \left(- \int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right),$$

and the curves which are formed by the trajectories of the differential system (2), in cartesian coordinates are written as

$$(x^2 + y^2)^{\frac{1}{2}} - h \exp \left(\int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right) = 0,$$

where $h \in \mathbb{R}$.

(3) *If $f_3(\theta) = 0$ for all $\theta \in \mathbb{R}$, then system (2) has the first integral $H = \frac{y}{x}$, and the curves which are formed by the trajectories of the differential system (2), in cartesian coordinates are written as $y - hx = 0$, where $h \in \mathbb{R}$.*

Proof. In order to prove our results we write the polynomial differential system (2) in polar coordinates (r, θ) , defined by $x = r \cos \theta$ and $y = r \sin \theta$, then system (2) becomes

$$\begin{cases} r' = f_1(\theta) r^2 + f_2(\theta) r^a, \\ \theta' = f_3(\theta) r^{a-1}, \end{cases} \quad (3)$$

where the trigonometric functions $f_1(\theta)$, $f_2(\theta)$, $f_3(\theta)$ are given in introduction, $r' = \frac{dr}{dt}$ and $\theta' = \frac{d\theta}{dt}$.

If $f_3(\theta) N(\cos \theta, \sin \theta) S(\cos \theta, \sin \theta) \neq 0$ and $a \neq 2$.

Taking as independent variable the coordinate θ , this differential system (3) writes

$$\frac{dr}{d\theta} = A(\theta) r + B(\theta) r^{3-a}, \quad (4)$$

where $A(\theta) = \frac{f_2(\theta)}{f_3(\theta)}$ and $B(\theta) = \frac{f_1(\theta)}{f_3(\theta)}$, which is a Bernoulli equation.

By introducing the standard change of variables $\rho = r^{a-2}$ we obtain the linear equation

$$\frac{d\rho}{d\theta} = (a-2)(A(\theta)\rho + B(\theta)). \quad (5)$$

The general solution of linear equation (5) is

$$\begin{aligned} \rho(\theta) &= \exp\left((a-2) \int_0^\theta A(\omega) d\omega\right) \\ &\quad \left(\mu + (a-2) \int_0^\theta \exp\left((2-a) \int_0^w A(\omega) d\omega\right) B(w) dw\right), \end{aligned}$$

where $\mu \in \mathbb{R}$, which has the first integral

$$\begin{aligned} H(x, y) &= (x^2 + y^2)^{\frac{a-2}{2}} \exp\left((2-a) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega\right) - \\ &\quad (a-2) \int_0^{\arctan \frac{y}{x}} \exp\left((2-a) \int_0^w A(\omega) d\omega\right) B(w) dw. \end{aligned}$$

Let Γ be a periodic orbit surrounding an equilibrium located in one of the open quadrants, and let $h_\Gamma = H(\Gamma)$.

The curves $H = h$ with $h \in \mathbb{R}$, which are formed by trajectories of the differential system (2), in cartesian coordinates are written as

$$x^2 + y^2 = \left(\begin{array}{c} h \exp\left((a-2) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega\right) + \\ (a-2) \exp\left((a-2) \int_0^{\arctan \frac{y}{x}} A(\omega) d\omega\right) \\ \int_0^{\arctan \frac{y}{x}} \exp\left((2-a) \int_0^w A(\omega) d\omega\right) B(w) dw \end{array} \right)^{\frac{2}{a-2}},$$

where $h \in \mathbb{R}$.

Hence statement (1) of Theorem 1 is proved.

Suppose now that $f_3(\theta) N(\cos \theta, \sin \theta) S(\cos \theta, \sin \theta) \neq 0$ and $a = 2$.

Taking as independent variable the coordinate θ , this differential system (3) writes

$$\frac{dr}{d\theta} = (A(\theta) + B(\theta)) r. \quad (6)$$

The general solution of equation (6) is

$$r(\theta) = \mu \exp\left(\int_0^\theta (A(\omega) + B(\omega)) d\omega\right),$$

where $\mu \in \mathbb{R}$, which has the first integral

$$H(x, y) = (x^2 + y^2)^{\frac{1}{2}} \exp \left(- \int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right).$$

The curves $H = h$ with $h \in \mathbb{R}$, which are formed by trajectories of the differential system (2), in cartesian coordinates are written as

$$(x^2 + y^2)^{\frac{1}{2}} - h \exp \left(\int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right) = 0,$$

where $h \in \mathbb{R}$.

Hence statement (2) of Theorem 1 is proved.

Assume now that $f_3(\theta) = 0$ for all $\theta \in \mathbb{R}$, then from system (3) it follows that $\theta' = 0$. So the straight lines through the origin of coordinates of the differential system (2) are invariant by the flow of this system. Hence, $\frac{y}{x}$ is a first integral of the system, then curves which are formed by the trajectories of the differential system (2), in cartesian coordinates are written as $y - hx = 0$, where $h \in \mathbb{R}$, since all straight lines through the origin are formed by trajectories.

This completes the proof of statement (3) of Theorem 1. $\square$

3 Examples

The following examples are given to illustrate our result

Example 1 If we take $\lambda = 1$, $\beta = -2$, $P(x, y) = 2x - 3y$, $Q(x, y) = 3x + 2y$, $M(x, y) = x^2 + 2y^2$, $N(x, y) = x^2 + y^2$, $R(x, y) = x^4 + 3x^2y^2 + y^4$ and $S(x, y) = x^4 + 2x^2y^2 + y^4$, then system (2) reads

$$\begin{cases} x' = 2x - 3y + x \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{x^4 + 3x^2y^2 + y^4}{x^4 + 2x^2y^2 + y^4} \right) \right), \\ y' = 3x + 2y + y \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{x^4 + 3x^2y^2 + y^4}{x^4 + 2x^2y^2 + y^4} \right) \right), \end{cases} \quad (7)$$

the differential system (7) in polar coordinates (r, θ) becomes

$$\begin{cases} r' = ((\cos \theta) \exp(1 + \sin^2 \theta) - 2(\sin \theta) \exp(\frac{9}{8} - \frac{1}{8} \cos 4\theta)) r^2 + 2r, \\ \theta' = 3, \end{cases}$$

here $f_1(\theta) = (\cos \theta) \exp(1 + \sin^2 \theta) - 2(\sin \theta) \exp(\frac{9}{8} - \frac{1}{8} \cos 4\theta)$, $f_2(\theta) = 2$ and $f_3(\theta) = 3$, it is the case (1) of the Theorem 1.

The differential system (7) has the first integral

$$H(x, y) = (x^2 + y^2)^{\frac{-1}{2}} \exp \left(\frac{2}{3} \arctan \frac{y}{x} \right) - \int_0^{\arctan \frac{y}{x}} \exp \left(\frac{2}{3} w \right) B(w) dw,$$

where $B(w) = \frac{(\cos w) \exp(1 + \sin^2 w) - 2(\sin w) \exp(\frac{9}{8} - \frac{1}{8} \cos 4w)}{3}$.

The curves $H = h$ with $h \in \mathbb{R}$, which are formed by trajectories of the differential system (7), in cartesian coordinates are written as

$$x^2 + y^2 = \left[\left(h + \int_0^{\arctan \frac{y}{x}} \exp \left(\frac{2}{3} w \right) B(w) dw \right) \exp \left(\frac{-2}{3} \arctan \frac{y}{x} \right) \right]^{-2},$$

where $h \in \mathbb{R}$.

Example 2 If we take $\lambda = 1$, $\beta = -2$, $P(x, y) = 5x^2 + 2xy$, $Q(x, y) = -2xy + 5y^2$, $M(x, y) = x^2 + 2y^2$, $N(x, y) = x^2 + y^2$, $R(x, y) = y$ and $S(x, y) = x$, then system (2) reads

$$\begin{cases} x' = 5x^2 + 2xy + x \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{y}{x} \right) \right), \\ y' = -2xy + 5y^2 + y \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{y}{x} \right) \right), \end{cases} \quad (8)$$

the differential system (8) in polar coordinates (r, θ) becomes

$$\begin{cases} r' = (7 \cos^3 \theta + 3 \sin^3 \theta - 2 (\sin \theta) (-1 + \exp \tan \theta) + (\cos \theta) (-2 + \exp (1 + \sin^2 \theta))) r^2, \\ \theta' = (3 \cos \theta \sin^2 \theta - 7 \cos^2 \theta \sin \theta) r, \end{cases}$$

then the differential system (8) has the first integral

$$H(x, y) = (x^2 + y^2)^{\frac{1}{2}} \exp \left(- \int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right)$$

where $A(\omega) + B(\omega) = \frac{7 \cos^3 \omega + 3 \sin^3 \omega - 2 (\sin \omega) (-1 + \exp \tan \omega) + (\cos \omega) (-2 + \exp (1 + \sin^2 \omega))}{3 \cos \omega \sin^2 \omega - 7 \cos^2 \omega \sin \omega}$, it is the case (2) of the Theorem 1,

The curves $H = h$ with $h \in \mathbb{R}$, which are formed by trajectories of the differential system (8), in cartesian coordinates are written as

$$(x^2 + y^2)^{\frac{1}{2}} - h \exp \left(\int_0^{\arctan \frac{y}{x}} (A(\omega) + B(\omega)) d\omega \right) = 0,$$

where $h \in \mathbb{R}$.

Example 3 If we take $\lambda = 1$, $\beta = -2$, $P(x, y) = x^3 + xy^2$, $Q(x, y) = y^3 + yx^2$, $M(x, y) = x^2 + 2y^2$, $N(x, y) = x^2 + y^2$, $R(x, y) = x^4 + 3x^2y^2 + y^4$ and $S(x, y) = x^4 + 2x^2y^2 + y^4$, then system (2) reads

$$\begin{cases} x' = x^3 + xy^2 + x \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{x^4 + 3x^2y^2 + y^4}{x^4 + 2x^2y^2 + y^4} \right) \right), \\ y' = y^3 + yx^2 + y \left(x \exp \left(\frac{x^2 + 2y^2}{x^2 + y^2} \right) - 2y \exp \left(\frac{x^4 + 3x^2y^2 + y^4}{x^4 + 2x^2y^2 + y^4} \right) \right), \end{cases} \quad (9)$$

the differential system (9) in polar coordinates (r, θ) becomes

$$\begin{cases} r' = r' = r^3 + ((\cos \theta) \exp (1 + \sin^2 \theta) - 2 (\sin \theta) \exp (\frac{9}{8} - \frac{1}{8} \cos 4\theta)) r^2, \\ \theta' = 0, \end{cases} \quad (10)$$

it is the case (3) of the Theorem 1, then from system (10) it follows that $\theta' = 0$. Hence, $\frac{y}{x}$ is a first integral of the system, then curves which are formed by the trajectories of the differential system (9), in Cartesian coordinates are written as $y - hx = 0$, where $h \in \mathbb{R}$, since all straight lines through the origin are formed by trajectories.

4 Conclusion

The elementary method used in this paper seems to be fruitful to investigate planar differential systems of ODEs in order to obtain explicit expression for a first integral and characterizes its trajectories, this is a one of the classical tools in the classification of all trajectories of dynamical systems.

References

- [1] A. A. Andronov, E. A. Leontovich, I. I. Gordon, and A. L. Maier, *Qualitative Theory of Second-Order Dynamical Systems*, Wiley, New York, 1973. <https://doi.org/10.31559/glm2016.3.2.3>
- [2] A. Bendjeddou, R. Boukoucha, *Explicit limit cycles of a cubic polynomial differential systems*, Stud. Univ. Babes-Bolyai Math, 2016, vol. 61, no. 1, pp. 77-85.
- [3] R. Boukoucha, *On the Dynamics of a Class of Kolmogorov Systems*, Journal of Siberian Federal University. Mathematics & Physics 2016, vol. 9, no. 1, pp. 11-16. <https://doi.org/10.4236/ajcm.2013.33026>
- [4] R. Boukoucha, *Explicit limit cycles of a family of polynomial differential systems*, Electronic Journal of Differential Equations, 2017, vol. 2017, no. 217, pp. 1-7. <https://doi.org/10.1016/j.amc.2005.07.072>
- [5] R. Boukoucha, A. Bendjeddou, *On the dynamics of a class of rational Kolmogorov systems*, Journal of Nonlinear Mathematical Physics, 2016, vol. 23, no. 1, pp. 21-27.

- [6] L. Cairó, J. Llibre, *Phase portraits of cubic polynomial vector fields of Lotka–Volterra type having a rational first integral of degree 2*, J. Phys, 2007, A 40, pp. 6329–6348.
- [7] M. Carbonell, B. Coll and J. Llibre, *Limit cycles of polynomial systems with homogeneous nonlinearities*, J. of Math. Anal. and Appl, 1989, vol. 142, pp. 573–590.
- [8] J. Chavarriga, H. Giacomini, J. Giné, J. Llibre, *Darboux integrability and the inverse integrating factor*, J. Differential Equations, 2003, vol. 194, pp. 116–139.
- [9] C. Christopher, C. Li, S. Yakovenko, *Advanced Course On Limit Cycles of Differential Equations*, Centre de Recerca Matemàtica Bellaterra (Spain), (2006).
- [10] F. Dumortier, J. Llibre and J. Artés, *Qualitative Theory of Planar Differential Systems*, (Universitext) Berlin, Springer, (2006).
- [11] P. Gao, *Hamiltonian structure and first integrals for the Lotka–Volterra systems*, Phys. Lett, 2000, A 273, pp. 85–96.
- [12] J. Giné. *Polynomial first integrals via the Poincaré series*, J. Comput. Appl. Math, 2005, vol. 184 , pp. 428–441.
- [13] J. Giné, M. Grau, *Coexistence of algebraic and non-algebraic limit cycles, explicitly given, using Riccati equations*. Nonlinearity, 2006, vol. 19, pp. 1939–1950.
- [14] N. A. Korol, *The integral curves of a certain differential equation*, (in Russian), Minsk. Gos. Ped. Inst. Minsk, 1973, pp. 47–51.
- [15] D. Hilbert, *Mathematische Probleme*, Lecture, Second Internat, English transl, Bull. Amer. Math. Soc, 1902, vol. 8, pp. 437–479.
- [16] L. Perko, *Differential equations and dynamical systems*, Third edition. Texts in Applied Mathematics, 7. Springer-Verlag, New York, 2001.