General Letters in Mathematics (GLM)

Journal Homepage: <http://www.refaad.com/views/GLM/home.aspx>

ISSN: 2519-9277 (Online) 2519-9269 (Print)

Micro β -open sets in Micro topology

Hariwan Z. Ibrahim¹

Department of Mathematics, Faculty of Education, University of Zakho, Zakho, Kurdistan Region-Iraq.

Abstract

In this paper, Micro β -open in micro topological space is introduced and basic properties of Micro β -open sets are analyzed and also this set used to introduce and study the new types of functions called Micro β -irresolute, Micro β -continuous and Micro β -continuous functions..©2020 All rights reserved.

Keywords: Nano topology, Micro topology, Micro open set, Micro β -open set .

2010 MSC: Primary: 54A05, 54A10; Secondary: 54C05

1. Introduction

The concept of nano topology was introduced by Thivagar et al [12] which is defined in terms of the lower and upper approximations and the boundary region of a subset of an universe. The notion of approximations and boundary region of a set was originally proposed by Pawlak [17] in order to introduce the concept of rough set theory. In 2019, Chandrasekar [1] introduced the concept of micro topology which is a simple extension of nano topology and he also studied the concepts of Micro pre-open and Micro semi-open sets. Chandrasekar and Swathi [2] introduced Micro α -open in micro topological space. Thivagar and Richard [14] established the weak forms of nano open set and also they introduced nano-regular open set in a nano topological space. Parimala et al [16] introduced the notion nano β -open set. Ibrahim [11] introduced the notion Micro β -open set. In the past few years, different forms of β -open sets have been studied Ibrahim [3, 4, 5, 6, 7, 8, 9, 10].

2. Preliminaries

Definition 2.1. [15] Let U be a nonempty finite set of objects called the universe and R be an equivalence relation on U named as the indiscernibility relation. Elements belonging to the same equivalence class are said to be indiscernible with one another. The pair (U, R) is said to be the approximation space. Let $X \subseteq U$.

1. The lower approximation of U with respect to R is the set of all objects, which can be for certain classified as X with respect to R and its is denoted by $L_R(X)$. That is, $L_R(X) = \bigcup_{x \in U} \{R(x) : R(x) \subseteq X\}$, where $R(x)$ denotes the equivalence class determined by x .

*Corresponding author

Email address: hariwan.ibrahim@uoz.edu.krd (Hariwan Z. Ibrahim)

doi:[10.31559/glm2020.8.1.2](https://doi.org/10.31559/glm2020.8.1.2)

Received 2 Dec 2019 : Accepted 24 Mar 2020

2. The upper approximation of U with respect to R is the set of all objects, which can be possibly classified as X with respect to R and it is denoted by $U_R(X)$. That is, $U_R(X) = \bigcup_{x \in U} \{R(x) : R(x) \cap X \neq \emptyset\}$.
3. The boundary region of U with respect to R is the set of all objects, which can be classified neither as X nor as not- X with respect to R and it is denoted by $B_R(X)$. That is, $B_R(X) = U_R(X) - L_R(X)$.

Definition 2.2. [12, 13] Let U be the universe, R be an equivalence relation on U and $\tau_R(X) = \{U, \emptyset, L_R(X), U_R(X), B_R(X)\}$, where $X \subseteq U$. Then, $\tau_R(X)$ satisfies the following axioms:

1. U and $\emptyset \in \tau_R(X)$.
2. The union of the elements of any subcollection of $\tau_R(X)$ is in $\tau_R(X)$.
3. The intersection of the elements of any finite subcollection of $\tau_R(X)$ is in $\tau_R(X)$.

That is, $\tau_R(X)$ is a topology on U called the nano topology on U with respect to X . We call $(U, \tau_R(X))$ as the nano topological space. The elements of $\tau_R(X)$ are called as nano open sets. A subset F of U is nano closed if its complement is nano open.

Definition 2.3. [12] Let $(U, \tau_R(X))$ be a nano topological space with respect to X where $X \subseteq U$ and $K \subseteq U$, then the nano interior of K is defined as the union of all nano open subsets of K and it is denoted by $NInt(K)$. The nano closure of K is defined as the intersection of all nano closed sets containing K and it is denoted by $NCl(K)$.

Definition 2.4. A subset K of a nano topological space $(U, \tau_R(X))$ is called:

1. nano-regular open if $NInt(NCl(K)) = K$ [14].
2. nano β -open if $K \subseteq NCl(NInt(K)) \cup NInt(NCl(K))$ [16].

Definition 2.5. [1] Let $(U, \tau_R(X))$ be a nano topological space. Then, $\mu_R(X) = \{N \cup (N' \cap \mu) : N, N' \in \tau_R(X) \text{ and } \mu \notin \tau_R(X)\}$ is called the micro topology on U with respect to X . The triplet $(U, \tau_R(X), \mu_R(X))$ is called micro topological space and the elements of $\mu_R(X)$ are called Micro open sets and the complement of a Micro open set is called a Micro closed set. Every nano topology is micro topology.

Definition 2.6. [1] The Micro closure of a set A is denoted by $Mic-cl(A)$ and is defined as $Mic-cl(A) = \bigcap \{B : B \text{ is Micro closed and } A \subseteq B\}$. The Micro interior of a set A is denoted by $Mic-int(A)$ and is defined as $Mic-int(A) = \bigcup \{B : B \text{ is Micro open and } A \supseteq B\}$.

Definition 2.7. Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space and $A \subseteq U$. Then,

1. A is called Micro α -open if $A \subseteq Mic-int(Mic-cl(Mic-int(A)))$ [2].
2. A is called Micro pre-open if $A \subseteq Mic-int(Mic-cl(A))$ [1].
3. A is called Micro semi-open if $A \subseteq Mic-cl(Mic-int(A))$ [1].
4. A is called Micro b -open if $A \subseteq Mic-int(Mic-cl(A)) \cup Mic-cl(Mic-int(A))$ [11].

Theorem 2.8. [1] Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space. Let A and B be any two subsets of U . Then:

1. A is a Micro open set if and only if $Mic-int(A) = A$.
2. A is a Micro closed set if and only if $Mic-cl(A) = A$.
3. $Mic-int(U \setminus A) = U \setminus Mic-cl(A)$.
4. $Mic-cl(U \setminus A) = U \setminus Mic-int(A)$.

Remark 2.9. [11] Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space. Let A and B be any two subsets of U . Then:

1. $Mic-cl(A)$ is Micro closed.

2. $A \subseteq \text{Mic-cl}(A)$.
3. $U \setminus \text{Mic-cl}(U \setminus A) = \text{Mic-int}(A)$.
4. $U \setminus \text{Mic-int}(U \setminus A) = \text{Mic-cl}(A)$.
5. $x \in \text{Mic-cl}(A)$ if and only if for every Micro open subset L of U containing x , $A \cap L \neq \emptyset$.

Remark 2.10. In [11], Ibrahim obtain the following diagram of implications:

3. Micro β -open Sets

Definition 3.1. Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space and $A \subseteq U$. Then, A is called a Micro β -open set if $A \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A)))$. $\text{Mic-}\beta O(U, X)$ denotes the collection of all Micro β -open sets. A subset B of U is called Micro β -closed if and only if its complement is Micro β -open. Moreover, $\text{Mic-}\beta C(U, X)$ denotes the collection of all Micro β -closed sets.

Remark 3.2. It can be shown that a subset B of U is Micro β -closed if and only if $\text{Mic-int}(\text{Mic-cl}(\text{Mic-int}(B))) \subseteq B$.

Theorem 3.3. Every Micro b -open is Micro β -open.

Proof. Let A be Micro b -open, then $A \subseteq \text{Mic-int}(\text{Mic-cl}(A)) \cup \text{Mic-cl}(\text{Mic-int}(A)) \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A))) \cup \text{Mic-cl}(\text{Mic-int}(A)) \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A)) \cup \text{Mic-int}(A)) \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A)) \cup \text{Mic-int}(\text{Mic-cl}(A))) \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A)))$. Thus, A is Micro β -open.

The converse of the above theorem need not be true in general as it is shown below.

Example 3.4. Consider $U = \{a, b, c, d, e\}$ with $U/R = \{\{a\}, \{b\}, \{c\}, \{d\}, \{e\}\}$ and $X = \{a, c\}$. Then, $\tau_R(X) = \{U, \emptyset, \{a, c\}\}$. If $\mu = \{e\}$, then $\mu_R(X) = \{U, \emptyset, \{a, c\}, \{a, c, e\}, \{e\}\}$ and $\{b, c, d\}$ is Micro β -open but not Micro b -open.

Remark 3.5. The concepts of Micro β -open and nano β -open sets are independent.

Example 3.6. Consider $U = \{a, b, c\}$ with $U/R = \{\{a\}, \{b, c\}\}$ and $X = \{a\}$. Then, $\tau_R(X) = \{U, \emptyset, \{a\}\}$. If $\mu = \{c\}$, then $\mu_R(X) = \{U, \emptyset, \{a\}, \{c\}, \{a, c\}\}$ and $\{c\}$ is Micro β -open but $\{c\}$ is not nano β -open.

Example 3.7. Consider $U = \{a, b, c\}$ with $U/R = \{\{a, b, c\}\}$ and $X = \{b, c\}$. Then, $\tau_R(X) = \{U, \emptyset\}$. If $\mu = \{a\}$, then $\mu_R(X) = \{U, \emptyset, \{a\}\}$ and $\{b\}$ is nano β -open but $\{b\}$ is not Micro β -open.

Remark 3.8. Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space. Then,

1. Every Micro open is Micro β -open.
2. Every nano open is Micro β -open.
3. Every nano-regular open is Micro β -open.
4. Every Micro α -open is Micro β -open.
5. Every Micro pre-open is Micro β -open.
6. Every Micro semi-open is Micro β -open.

Remark 3.9. From Remark 2.10 and Theorem 3.3, we obtain the following diagram of implications:

Theorem 3.10. An arbitrary union of Micro β -open sets is Micro β -open.

Proof. Let $\{A_i : i \in I\}$ be a family of Micro β -open sets. Then for each i , $A_i \subseteq \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A_i)))$ and so

$$\begin{aligned}
 \cup_{i \in I} A_i &\subseteq \cup_{i \in I} [\text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A_i)))] \\
 &\subseteq [\cup_{i \in I} \text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(A_i)))] \\
 &\subseteq [\text{Mic-cl}(\cup_{i \in I} \text{Mic-int}(\text{Mic-cl}(A_i)))] \\
 &\subseteq [\text{Mic-cl}(\text{Mic-int}(\cup_{i \in I} \text{Mic-cl}(A_i)))] \\
 &\subseteq [\text{Mic-cl}(\text{Mic-int}(\text{Mic-cl}(\cup_{i \in I} A_i)))] .
 \end{aligned}$$

Thus, $\cup_{i \in I} A_i$ is a Micro β -open set.

Remark 3.11. 1. An arbitrary intersection of Micro β -closed sets is Micro β -closed.

2. The intersection of two Micro β -open sets may not be Micro β -open.

Example 3.12. From Example 3.6, $A = \{a, b\}$ and $B = \{b, c\}$ are Micro- β -open, but $A \cap B = \{b\}$ which is not Micro- β -open.

Definition 3.13. Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space. Then:

1. The union of all Micro β -open sets contained in A is called the Micro β -interior of A and denoted by $\text{Mic-}\beta\text{int}(A)$.
2. The intersection of all Micro β -closed sets containing A is called the Micro β -closure of A and denoted by $\text{Mic-}\beta\text{cl}(A)$.

Now, we state the following theorem without proof.

Theorem 3.14. Let $(U, \tau_R(X), \mu_R(X))$ be a micro topological space. For any subsets A, B of U , we have the following:

1. A is Micro β -open if and only if $A = \text{Mic-}\beta\text{int}(A)$.
2. A is Micro β -closed if and only if $A = \text{Mic-}\beta\text{cl}(A)$.
3. If $A \subseteq B$, then $\text{Mic-}\beta\text{int}(A) \subseteq \text{Mic-}\beta\text{int}(B)$ and $\text{Mic-}\beta\text{cl}(A) \subseteq \text{Mic-}\beta\text{cl}(B)$.
4. $\text{Mic-}\beta\text{int}(A) \cup \text{Mic-}\beta\text{int}(B) \subseteq \text{Mic-}\beta\text{int}(A \cup B)$.
5. $\text{Mic-}\beta\text{int}(A \cap B) \subseteq \text{Mic-}\beta\text{int}(A) \cap \text{Mic-}\beta\text{int}(B)$.
6. $\text{Mic-}\beta\text{cl}(A) \cup \text{Mic-}\beta\text{cl}(B) \subseteq \text{Mic-}\beta\text{cl}(A \cup B)$.
7. $\text{Mic-}\beta\text{cl}(A \cap B) \subseteq \text{Mic-}\beta\text{cl}(A) \cap \text{Mic-}\beta\text{cl}(B)$.
8. $\text{Mic-}\beta\text{int}(U \setminus A) = U \setminus \text{Mic-}\beta\text{cl}(A)$.

9. $\text{Mic-}\beta\text{cl}(\mathcal{U} \setminus A) = \mathcal{U} \setminus \text{Mic-}\beta\text{int}(A)$.
10. $\mathcal{U} \setminus \text{Mic-}\beta\text{cl}(\mathcal{U} \setminus A) = \text{Mic-}\beta\text{int}(A)$.
11. $\mathcal{U} \setminus \text{Mic-}\beta\text{int}(\mathcal{U} \setminus A) = \text{Mic-}\beta\text{cl}(A)$.
12. $x \in \text{Mic-}\beta\text{int}(A)$ if and only if there exists a Micro β -open set L such that $x \in L \subseteq A$.

Theorem 3.15. Let A be a subset of a micro topological space $(\mathcal{U}, \tau_R(X), \mu_R(X))$. Then, $x \in \text{Mic-}\beta\text{cl}(A)$ if and only if for every Micro β -open subset L of $\mathcal{U}$ containing x , $A \cap L \neq \emptyset$.

Proof. Let $x \in \text{Mic-}\beta\text{cl}(A)$ and suppose that $L \cap A = \emptyset$ for some Micro β -open set L which contains x . Then, $(\mathcal{U} \setminus L)$ is Micro β -closed and $A \subset (\mathcal{U} \setminus L)$, thus $\text{Mic-}\beta\text{cl}(A) \subset (\mathcal{U} \setminus L)$. But this implies that $x \in (\mathcal{U} \setminus L)$, a contradiction. Thus, $L \cap A \neq \emptyset$.

Conversely, let $A \subseteq \mathcal{U}$ and $x \in \mathcal{U}$ such that for each Micro β -open set L_1 which contains x , $L_1 \cap A \neq \emptyset$. If $x \notin \text{Mic-}\beta\text{cl}(A)$, there is a Micro β -closed set F such that $A \subseteq F$ and $x \notin F$. Then, $(\mathcal{U} \setminus F)$ is a Micro β -open set with $x \in (\mathcal{U} \setminus F)$, and thus $(\mathcal{U} \setminus F) \cap A \neq \emptyset$, which is a contradiction.

Theorem 3.16. Let $(\mathcal{U}, \tau_R(X), \mu_R(X))$ be a micro topological space and $L \subseteq \mathcal{U}$. Then, L is Micro β -open if and only if for each $s \in L$, there exists a Micro β -open set D such that $s \in D \subseteq L$.

Proof. It is obvious.

Definition 3.17. A function $f : \mathcal{U} \rightarrow V$ is said to be Micro β -irresolute if $f^{-1}(K)$ is a Micro β -open set in $\mathcal{U}$, for every Micro β -open set K in V .

Theorem 3.18. Let $f : \mathcal{U} \rightarrow V$ be a function, then the following statements are equivalent:

1. f is Micro β -irresolute.
2. $f(\text{Mic-}\beta\text{cl}(A)) \subseteq \text{Mic-}\beta\text{cl}(f(A))$ holds for every subset A of $\mathcal{U}$.
3. $f^{-1}(B)$ is Micro β -closed in $\mathcal{U}$, for every Micro β -closed subset B of V .

Proof. (2) $\Rightarrow$ (3): Let B be a Micro β -closed set in V , then $\text{Mic-}\beta\text{cl}(B) = B$. By using (2), we have $f(\text{Mic-}\beta\text{cl}(f^{-1}(B))) \subseteq \text{Mic-}\beta\text{cl}(B) = B$. Thus, $\text{Mic-}\beta\text{cl}(f^{-1}(B)) \subseteq f^{-1}(B)$ and hence $f^{-1}(B)$ is Micro β -closed in $\mathcal{U}$.

(3) $\Rightarrow$ (2): If $A \subseteq \mathcal{U}$, then $\text{Mic-}\beta\text{cl}(f(A))$ is Micro β -closed in V and by (3) $f^{-1}(\text{Mic-}\beta\text{cl}(f(A)))$ is Micro β -closed in $\mathcal{U}$. Furthermore, $A \subseteq f^{-1}(f(A)) \subseteq f^{-1}(\text{Mic-}\beta\text{cl}(f(A)))$. Thus, $\text{Mic-}\beta\text{cl}(A) \subseteq f^{-1}(\text{Mic-}\beta\text{cl}(f(A)))$, consequently, $f(\text{Mic-}\beta\text{cl}(A)) \subseteq f(f^{-1}(\text{Mic-}\beta\text{cl}(f(A)))) \subseteq \text{Mic-}\beta\text{cl}(f(A))$.

(3) $\Leftrightarrow$ (1): Obvious.

Definition 3.19. A function $f : \mathcal{U} \rightarrow V$ is said to be Micro β -continuous at a point $x \in \mathcal{U}$ if for each Micro open subset K of V containing $f(x)$, there exists a Micro β -open subset L of $\mathcal{U}$ containing x such that $f(L) \subseteq K$. The function f is said to be Micro β -continuous if it has this property at each $x \in \mathcal{U}$.

Definition 3.20. A function $f : \mathcal{U} \rightarrow V$ is said to be Micro Wb-continuous at a point $x \in \mathcal{U}$ if for each Micro open subset K of V containing $f(x)$, there exists a Micro β -open subset L of $\mathcal{U}$ containing x such that $f(L) \subseteq \text{Mic-cl}(K)$. The function f is said to be Micro Wb-continuous if it has this property at each $x \in \mathcal{U}$.

Remark 3.21. It is obvious from the above definition that Micro β -continuous implies Micro Wb-continuous. However, the converse is not true in general as it is shown in the following example.

Example 3.22. Consider $\mathcal{U} = \{a, b, c\}$ with $\mathcal{U}/R = \{\{a\}, \{b, c\}\}$ and $X = \{a\}$. Then, $\tau_R(X) = \{\mathcal{U}, \emptyset, \{a\}\}$. If $\mu = \{b\}$, then $\mu_R(X) = \{\mathcal{U}, \emptyset, \{a\}, \{b\}, \{a, b\}\}$ and $\text{Mic-}\beta\text{O}(\mathcal{U}, X) = \{\mathcal{U}, \emptyset, \{a\}, \{b\}, \{a, b\}, \{a, c\}, \{b, c\}\}$. Consider $V = \{a, b, c, d\}$ with $V/R = \{\{a\}, \{c\}, \{b, d\}\}$ and $Y = \{b, d\}$. Then, $\tau_R(Y) = \{V, \emptyset, \{b, d\}\}$. If $\mu = \{a\}$, then $\mu_R(Y) = \{V, \emptyset, \{a\}, \{b, d\}, \{a, b, d\}\}$. Define a function $f : \mathcal{U} \rightarrow V$ as follows:

$$f(x) = \begin{cases} c & \text{if } x = a \\ a & \text{if } x = b \\ d & \text{if } x = c \end{cases}$$

Then, f is Micro Wb -continuous, but not Micro β -continuous, because $\{b, d\}$ is a Micro open set in V containing $f(c) = d$, but there exist no Micro β -open set L in U containing c such that $f(L) \subseteq \{b, d\}$.

Theorem 3.23. A function $f : U \rightarrow V$ is Micro β -continuous if and only if the inverse image of every Micro open set in V is Micro β -open in U .

Proof. Let f be Micro β -continuous and K be any Micro open set in V . If $f^{-1}(K) = \emptyset$, then $f^{-1}(K)$ is a Micro β -open set in U but if $f^{-1}(K) \neq \emptyset$, then there exists $x \in f^{-1}(K)$ which implies $f(x) \in K$. Since f is Micro β -continuous, then there exists a Micro β -open set L in U containing x such that $f(L) \subseteq K$. This implies that $x \in L \subseteq f^{-1}(K)$ and hence $f^{-1}(K)$ is Micro β -open.

Conversely, let K be any Micro open set in V containing $f(x)$, then $x \in f^{-1}(K)$ and by hypothesis $f^{-1}(K)$ is a Micro β -open set in U containing x , so $f(f^{-1}(K)) \subseteq K$. Thus, f is Micro β -continuous.

Theorem 3.24. For a function $f : U \rightarrow V$, the following statements are equivalent:

1. f is Micro β -continuous.
2. $f^{-1}(K)$ is a Micro β -open set in U , for each Micro open subset K of V .
3. $f^{-1}(F)$ is a Micro β -closed set in U , for each Micro closed subset F of V .
4. $f(\text{Mic-}\beta\text{cl}(A)) \subseteq \text{Mic-cl}(f(A))$, for each subset A of U .
5. $\text{Mic-}\beta\text{cl}(f^{-1}(B)) \subseteq f^{-1}(\text{Mic-cl}(B))$, for each subset B of V .
6. $f^{-1}(\text{Mic-int}(B)) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(B))$, for each subset B of V .

Proof. (1) $\Rightarrow$ (2): Directly from Theorem 3.23.

(2) $\Rightarrow$ (3): Let F be any Micro closed subset of V . Then, $V \setminus F$ is a Micro open subset of V . By (2), $f^{-1}(V \setminus F) = U \setminus f^{-1}(F)$ is a Micro β -open set in U and hence $f^{-1}(F)$ is a Micro β -closed set in U .

(3) $\Rightarrow$ (4): Let A be any subset of U . Then, $f(A) \subseteq \text{Mic-cl}(f(A))$ and $\text{Mic-cl}(f(A))$ is a Micro closed set in V . Hence, $A \subseteq f^{-1}(\text{Mic-cl}(f(A)))$. By (3), we have $f^{-1}(\text{Mic-cl}(f(A)))$ is a Micro β -closed set in U . Therefore, $\text{Mic-}\beta\text{cl}(A) \subseteq f^{-1}(\text{Mic-cl}(f(A)))$. Hence, $f(\text{Mic-}\beta\text{cl}(A)) \subseteq \text{Mic-cl}(f(A))$.

(4) $\Rightarrow$ (5): Let B be any subset of V . Then, $f^{-1}(B)$ is a subset of U . By (4), we have $f(\text{Mic-}\beta\text{cl}(f^{-1}(B))) \subseteq \text{Mic-cl}(f(f^{-1}(B))) \subseteq \text{Mic-cl}(B)$. Hence, $\text{Mic-}\beta\text{cl}(f^{-1}(B)) \subseteq f^{-1}(\text{Mic-cl}(B))$.

(5) $\Leftrightarrow$ (6): Let B be any subset of V . Then, apply (5) to $V \setminus B$ we obtain $\text{Mic-}\beta\text{cl}(f^{-1}(V \setminus B)) \subseteq f^{-1}(\text{Mic-cl}(V \setminus B)) \Leftrightarrow \text{Mic-}\beta\text{cl}(U \setminus f^{-1}(B)) \subseteq f^{-1}(V \setminus \text{Mic-int}(B)) \Leftrightarrow U \setminus \text{Mic-}\beta\text{int}(f^{-1}(B)) \subseteq U \setminus f^{-1}(\text{Mic-int}(B)) \Leftrightarrow f^{-1}(\text{Mic-int}(B)) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(B))$. Thus, $f^{-1}(\text{Mic-int}(B)) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(B))$.

(6) $\Rightarrow$ (1): Let $x \in U$ and K be any Micro open subset of V containing $f(x)$. By (6), we have $f^{-1}(\text{Mic-int}(K)) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(K))$ implies that $f^{-1}(K) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(K))$. Hence, $f^{-1}(K)$ is a Micro β -open set in U which contains x and clearly $f(f^{-1}(K)) \subseteq K$. Thus, f is Micro β -continuous.

Theorem 3.25. For a function $f : U \rightarrow V$, the following statements are equivalent:

1. f is Micro Wb -continuous.
2. $f^{-1}(K) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$, for every Micro open subset K of V .
3. $\text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(F))) \subseteq f^{-1}(F)$, for every Micro closed subset F of V .
4. $\text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(\text{Mic-cl}(B)))) \subseteq f^{-1}(\text{Mic-cl}(B))$, for every subset B of V .
5. $f^{-1}(\text{Mic-int}(B)) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(\text{Mic-int}(B))))$, for every subset B of V .
6. $\text{Mic-}\beta\text{cl}(f^{-1}(K)) \subseteq f^{-1}(\text{Mic-cl}(K))$, for every Micro open subset K of V .

Proof. (1) $\Rightarrow$ (2): Let K be a Micro open subset of V such that $x \in f^{-1}(K)$. Then, $f(x) \in K$ and there exists $L \in \text{Mic-}\beta\text{O}(U, x)$ containing x such that $f(L) \subseteq \text{Mic-cl}(K)$. Thus, we obtain $x \in L \subseteq f^{-1}(\text{Mic-cl}(K))$. This implies that $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$.

(2) $\Rightarrow$ (3): Let F be any Micro closed subset of V . Suppose that $x \notin f^{-1}(F)$, then $V \setminus F$ is Micro open in V and $x \in U \setminus f^{-1}(F) = f^{-1}(V \setminus F)$. By (2), we have $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(V \setminus F))) = \text{Mic-}\beta\text{int}(f^{-1}(V \setminus \text{Mic-int}(F))) = \text{Mic-}\beta\text{int}(U \setminus f^{-1}(\text{Mic-int}(F))) = U \setminus (\text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(F))))$. Therefore, we obtain $x \notin \text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(F)))$.

(3) $\Rightarrow$ (4): Let B be any subset of V . Then, $\text{Mic-cl}(B)$ is Micro closed in V and by (3), we have that if $x \in \text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(\text{Mic-cl}(B))))$, then $x \in f^{-1}(\text{Mic-cl}(B))$.

(4) $\Rightarrow$ (5): Let B be any subset of V and $x \in f^{-1}(\text{Mic-int}(B))$. Then, we have $x \in f^{-1}(\text{Mic-int}(B)) = U \setminus f^{-1}(\text{Mic-cl}(V \setminus B))$. Then, $x \notin f^{-1}(\text{Mic-cl}(V \setminus B))$ and by (4), $x \in U \setminus \text{Mic-}\beta\text{cl}(f^{-1}(\text{Mic-int}(\text{Mic-cl}(V \setminus B)))) = \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(\text{Mic-int}(B))))$.

(5) $\Rightarrow$ (6): Let K be any Micro open subset of V . Suppose that $x \notin f^{-1}(\text{Mic-cl}(K))$. Then, $f(x) \notin \text{Mic-cl}(K)$ and there exists a Micro open set W containing $f(x)$ such that $W \cap K = \emptyset$; hence $\text{Mic-cl}(W) \cap K = \emptyset$. By (5), we have $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(W)))$ and hence there exists $L \in \text{Mic-}\beta\text{O}(U, X)$ such that $x \in L \subseteq f^{-1}(\text{Mic-cl}(W))$. Since $\text{Mic-cl}(W) \cap K = \emptyset$, then $L \cap f^{-1}(K) = \emptyset$ and by Theorem 3.15, $x \notin \text{Mic-}\beta\text{cl}(f^{-1}(K))$. Therefore, if $x \in \text{Mic-}\beta\text{cl}(f^{-1}(K))$, then $x \in f^{-1}(\text{Mic-cl}(K))$.

(6) $\Rightarrow$ (1): Let $x \in U$ and K any Micro open subset of V containing $f(x)$. Then, we have $x \in f^{-1}(K) \subseteq f^{-1}(\text{Mic-int}(\text{Mic-cl}(K))) = U \setminus f^{-1}(\text{Mic-cl}(V \setminus \text{Mic-cl}(K)))$. By (6), $x \notin \text{Mic-}\beta\text{cl}(f^{-1}(V \setminus \text{Mic-cl}(K)))$ and hence $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$. Therefore, there exists $L \in \text{Mic-}\beta\text{O}(U, X)$ such that $x \in L \subseteq f^{-1}(\text{Mic-cl}(K))$ and hence $f(L) \subseteq \text{Mic-cl}(K)$. This shows that f is Micro Wb-continuous.

Theorem 3.26. For a function $f : U \rightarrow V$, the following statements are equivalent:

1. f is Micro Wb-continuous at $x \in U$.
2. $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$, for each Micro open set K containing $f(x)$.

Proof. (1) $\Rightarrow$ (2): Let K be any Micro open set containing $f(x)$. Then, there exists a Micro β -open set L containing x such that $f(L) \subseteq \text{Mic-cl}(K)$ and so $L \subseteq f^{-1}(\text{Mic-cl}(K))$. Since L is Micro β -open, then $x \in L = \text{Mic-}\beta\text{int}(L) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$.

(2) $\Rightarrow$ (1): Let $x \in \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$, for each Micro open set K containing $f(x)$. Take $L = \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$. Then, $f(L) \subseteq \text{Mic-cl}(K)$. Moreover, L is Micro β -open. Thus, f is Micro Wb-continuous at $x \in U$.

Theorem 3.27. For a function $f : U \rightarrow V$, the following statements are equivalent:

1. f is Micro Wb-continuous.
2. $\text{Mic-}\beta\text{cl}(f^{-1}(K)) \subseteq f^{-1}(\text{Mic-cl}(K))$, for every Micro pre-open K in V .
3. $f^{-1}(K) \subseteq \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$, for every Micro pre-open K in V .

Proof. (1) $\Rightarrow$ (2): Let K be any Micro pre-open subset of V such that $x \in \text{Mic-}\beta\text{cl}(f^{-1}(K))$. Suppose that $x \notin f^{-1}(\text{Mic-cl}(K))$. Then, there exists a Micro open set W containing $f(x)$ such that $W \cap K = \emptyset$. Thus, we have $W \cap \text{Mic-cl}(K) = \emptyset$ and hence $\text{Mic-cl}(W) \cap \text{Mic-int}(\text{Mic-cl}(K)) = \emptyset$. Since K is Micro pre-open, then $K \subseteq \text{Mic-int}(\text{Mic-cl}(K))$ and we have $K \cap \text{Mic-cl}(W) = \emptyset$. Since f is Micro Wb-continuous at $x \in U$ and W is a Micro open set containing $f(x)$, then there exists a Micro β -open subset L of U containing x such that $f(L) \subseteq \text{Mic-cl}(W)$. So, $f(L) \cap K = \emptyset$ and hence $L \cap f^{-1}(K) = \emptyset$. This shows that $x \notin \text{Mic-}\beta\text{cl}(f^{-1}(K))$. This is a contradiction. Therefore, $x \in f^{-1}(\text{Mic-cl}(K))$.

(2) $\Rightarrow$ (3): Let K be Micro pre-open in V and $x \in f^{-1}(K)$. Then, $f^{-1}(K) \subseteq f^{-1}(\text{Mic-int}(\text{Mic-cl}(K))) = U \setminus f^{-1}(\text{Mic-cl}(V \setminus \text{Mic-cl}(K)))$. Therefore, $x \notin f^{-1}(\text{Mic-cl}(V \setminus \text{Mic-cl}(K)))$. Then by (2), $x \notin \text{Mic-}\beta\text{cl}(f^{-1}(V \setminus \text{Mic-cl}(K)))$. Hence, $x \in U \setminus \text{Mic-}\beta\text{cl}(f^{-1}(V \setminus \text{Mic-cl}(K))) = \text{Mic-}\beta\text{int}(f^{-1}(\text{Mic-cl}(K)))$.

(3) $\Rightarrow$ (1): This follows from Theorem 3.25, since every Micro open set is Micro pre-open.

References

- [1] S. Chandrasekar, On micro topological spaces, Journal of New Theory, 26(2019), 23–31. [1](#), [2.5](#), [2.6](#), [2](#), [3](#), [2.8](#)
- [2] S. Chandrasekar and G. Swathi, Micro- α -open sets in micro topological spaces, International Journal of Research in Advent Technology, 6 (10)(2018),2633–2637. [1](#), [1](#)
- [3] H. Z. Ibrahim, Weak forms of γ -open sets and new separation axioms, Int. J. of Scientific and Engineering Research, 3 (4)(2012), 1–4. [1](#)
- [4] H. Z. Ibrahim, β -g-bicontinuous, β -g-compactness and β -g-stable in ditopological texture spaces, J. Korean Soc. Math. Educ. Ser. B: Pure Appl. Math., 26 (1) (2019), 35–47. [1](#)

- [5] H. Z. Ibrahim, β - γ -irresolute and β - γ -closed graph, Gen. Math. Notes, **15** (2) (2013), 32–44. [1](#)
- [6] H. Z. Ibrahim, On some separation axioms via β - γ -open sets, Gen. Math. Notes, **15** (2)(2013), 14–31. [1](#)
- [7] H. Z. Ibrahim, δ^* -open sets and relation between some weak and strong forms of δ^* -open sets in topological spaces, Journal of Advanced Studies in Topology, **6**(1)(2015), 20–27. [1](#)
- [8] H. Z. Ibrahim, τ^* -open sets and relation between some weak and strong forms of τ^* -open sets in topological spaces, International Journal of Engineering Sciences & Research Technology **1**, **4**(1)(2015), 672–677.
- [9] H. Z. Ibrahim, Strong forms of generalized closed sets in ditopological texture spaces, Journal of Advanced Studies in Topology, **6**(2)(2015), 61–68. <https://doi.org/10.20454/jast.2015.908> [1](#)
- [10] H. Z. Ibrahim, Applications of nano topology via nano operations, Submitted. [1](#)
- [11] H. Z. Ibrahim, On micro b -open sets, Submitted. [1](#), [4](#), [2.9](#), [2.10](#)
- [12] M. L. Thivagar and C. Richard, Note on nano topological spaces, Communicated. [1](#), [2.2](#), [2.3](#)
- [13] M. L. Thivagar, C. Richard and N. R. Paul, Mathematical Innovations of a Modern Topology in Medical Events, International Journal of Information Science, **2** (4) (2012), 33–36. <https://doi.org/10.5923/j.ijis.20120204.01> [2.2](#)
- [14] M. L. Thivagar and C. Richard, On nano forms of weakly open sets, International Journal of Mathematics and Statistics Invention, **1**(1)(2013), 31–37. [1](#), [1](#)
- [15] I. L. Reilly and M. K. Vamanamurthy, On α -sets in topological spaces, Tamkang J.Math., **16** (1985), 7–11. [2.1](#)
- [16] M. Parimala, C. Indiranti and S. Jafar, On nano β -open sets in nano topological spaces, Jordan Journal of Mathematics and Statistics, **9** (3)(2016), 173–184. [1](#), [2](#)
- [17] Z. Pawlak, Rough sets, International journal of information and computer sciences, **11** (1982), 341–356. [1](#)