IJCRR
Section: Healthcare
ISI Impact Factor
(2019-20): 1.628
IC Value (2019): 90.81
SJIF (2020) = 7.893

Copyright@IJCRR

Pyrimidine Chalcones as Thymidylate Kinase Inhibitors for treating Tuberculosis: A Letter to the Editor

Debarshi Kar Mahapatra

Department of Pharmaceutical Chemistry, Dadasaheb Balpande College of Pharmacy, Nagpur 440037, Maharashtra, India.

It is a pleasure writing a Letter to the Editor for the published article: **Rasapelly RK, Kannappan N, Jarpula D. A Novel Approach for Synthesis of Pyrimidine Chalcones Against Thymidylate Kinase Protein Targets And Evaluation of Antimycobacterial Activity. Int J Cur Res Rev. 2020;12(21):100-106.**

The World Health Organization (WHO) believes that nearly 2 billion individuals are infected with *Mycobacterium tuberculosis*, the bacterium that causes tuberculosis (TB). Last year, ten million people were sick with TB and 1.4 million died as a result of the disease. Many antibiotics are inherently resistant to the infection, reducing the number of drugs that may be used to treat it. Thick, waxy, hydrophobic cell membranes, as well as the presence of drug degrading and modifying enzymes, are all factors that contribute to intrinsic resistance. Despite significant attempts to adopt new pharmacotherapeutic methods for the treatment of tuberculosis, the illness continues to be a serious issue throughout the globe. As a result, new families of compounds with specific antimicrobial activity are urgently needed. The control of cellular pathways linked to pathogenesis is a well-known method for understanding a wide range of TB symptoms. As a result, identifying regulators and molecular target inhibitors to fight tuberculosis is a promising approach for discovering and developing anti-mycobacterial medicines.

Natural products have been found to have anti-infective properties. Chalcones, also known as 1,3-diphenyl-2E-propene-1-one, are a family of natural compounds that include the benzylideneacetophenone scaffold, which connects the two aromatic nuclei through a three-carbon α , β -unsaturated carbonyl bridge. Chalcones are an open-chain intermediate in the synthesis of flavones by aurones that may be found in various conjugated forms in nature as precursors to flavonoids and isoflavonoids. The pyrimidine chalcones are well known for exhibiting anti-infective activities such as anti-bacterial, anti-fungal, anti-mycobacterial, anti-malarial, anti-viral, and anti-parasitic. Chalcones having pyrimidine moieties as a part of the scaffold have recently been explored as promising anti-mycobacterial molecules against H₃₇Rv strain, mainly inhibiting protein tyrosine phosphatase A/B (Ptp-A/B) and thymidylate kinase, two key molecular targets in the pathogen.

This research paper represented the design, synthesis, characterization, molecular docking, and anti-tubercular screening of some pyrimidine chalcone inhibitors. Overall, the manuscript is really extremely interesting to read, really well written, the overall presentation is good, the produced data are quite adequate, the range of literature is relatively completely covered, and is factually imperative to the researchers / readers in the newer development of chalcone analogs and/or hybrid derivatives. While neglecting the minor errors of typographical and grammatical origin, however, few scientific expectations still remain unfulfilled to this excellently published paper:

1. ¹³C-NMR Spectral data and Mass data are lacking significantly.
2. *In silico* study with high-end software like Schrodinger (XP-Mode) would have been much better.
3. *In silico* pharmacokinetics is missing. It would have helped in identifying the most promising inhibitors.
4. Why only one target was selected in the *in silico* exploration? Multiple targets refer to multiple ways or more than one mechanism(s) or multiple approaches of therapy.
5. *In vitro* enzymatic screening of the most promising compound would have been better to practically identify the inhibitor affinity.
6. No structure-activity relationships (SARs) are established.

Although there are several positive points in this research, which must be appreciated:

1. Use of microwave irradiation technology.
2. Spectral characterization is completely and perfectly described.
3. The number of synthesized compounds is excellent.
4. The yield of the compounds is impressive.

On the whole, the study facilitated the exploration/development of the newer lead molecules against tuberculosis. Additional efforts must be taken up to optimize the lead structure and the results will positively encourage enthusiastic researches for future development of therapeutically privileged (semi)-synthetic anti-mycobacterial molecules.