SYNTHESIS AND CHARACTERIZATION OF ASTAXANTHIN-METAL IONS (Cu^{2+} and Zn^{2+}) COMPLEX

S. Wibowo^{1,✉}, B. D. Prakoso¹, S. Najihah¹, Z. H. Fajar¹, S. Widyarti¹,
A. Sabarudin², D. W. Soeatmadji³ and S. B. Sumitro¹

¹Department of Biology, Faculty of Mathematics and Natural Sciences, Brawijaya University Jl. Veteran, Malang 65145, East Java, Indonesia.

²Department of Chemistry, Faculty of Mathematics and Natural Sciences, Brawijaya University, Jl. Veteran, Malang 65145, East Java, Indonesia.

³Department of Internal, School of Medicine, Brawijaya University, Jl. Veteran, Malang 65145, East Java, Indonesia.

✉Corresponding Author: wibowo@student.ub.ac.id

ABSTRACT

This study aimed was to determine the optimal ratio, duration, and temperature reaction to form the astaxanthin-metal ions complex based on the UV-Vis spectrum profile and confirmation of complex formation using FTIR. The method used consists of the synthesis of the astaxanthin-metal ions (Cu^{2+} and Zn^{2+}) and the determination of complex formation based on the UV-Vis spectrum profile which is evaluated at a wavelength of 190–790 nm. Then the selected complexes were analyzed for their functional groups using FTIR at wavenumbers of 400–4000 cm^{-1} . Astaxanthin-metal ions complex was successfully synthesized and characterized by UV-Vis and FTIR. The optimal ratio, temperature, and stirrer time for astaxanthin- Cu^{2+} are (3:1), 78°C, and 5 minutes, while astaxanthin- Zn^{2+} is (1:1), 78°C, and 15 minutes, respectively.

Keywords: Astaxanthin, Cu^{2+} , FTIR, Metal Ions, UV-Vis, Zn^{2+} .

RASAYAN J. Chem., Vol. 14, No.2, 2021

INTRODUCTION

Oxidative stress is one of the contributing factors to the emergence of various problems in the cellular environment.¹ Oxidative stress occurs because free radicals number exceed the number of antioxidants that become cell defense capacity.² Free radicals are defined as compounds, molecules, or atoms that have unpaired electrons in their outermost orbitals.³ These electrons are unstable, tend to be reactive and attract electrons from nearby molecules.⁴ Antioxidants are necessary to balance the condition. One antioxidant that has high levels of radical scavenger is astaxanthin. Astaxanthin is a terpene lipophilic compound that has polyene chains capable of stabilizing free radicals.⁵ The radical scavenger mechanism performed by astaxanthin is carried out in three ways namely SET (single electron transfer), radical adduct formation, and HAT (hydrogen atom transfer).⁶ In addition to being a radical scavenger, astaxanthin has also been proven to play a role in the stability of human serum albumin protein and glycated human serum albumin by preventing protein unfolding.⁷ Astaxanthin bioinorganic complexes with metal cations were known to produce chelating metal ions as antioxidants due to the useful properties of metal cations in donors and electron acceptors referred to as Lewis acid-base.^{8,9} Metal ions play an essential role in many biological processes due to the specific role that only those metals have.¹⁰ It can improve the ability of astaxanthin as a radical scavenger of free radicals. Some metal ions that can be coordinated with astaxanthin are Cu^{2+} and Zn^{2+} . Cu is one of the essential metals required in various catalytic processes. Cu in the form of metal ions (Cu^{2+}) can form coordination bonds with oxygen atomic donors. This makes Cu^{2+} metal ions capable of binding with four oxygen atoms of astaxanthin.¹¹ Zn has a fully charged d^{10} valence atomic orbital making it more stable.¹² Also, Zn is an essential component involved in almost all functional aspects of the body, such as in the human proteome, Lewis acid-base with proteins, and maintaining the structure of catalysis enzymes.¹³

The aim of this study was to determine the optimal ratio, duration and temperature reaction to form the astaxanthin-essential metal ions (Cu^{2+} and Zn^{2+}) complex and determine the physicochemical characteristics of the astaxanthin-essential metal ions complex based on UV-Vis and FTIR analysis.

EXPERIMENTAL

Materials

Astaxanthin (SML0982, CAS No 472-61-7) from Sigma-Aldrich, Ethanol Absolute, Aquades, Zn^{2+} and Cu^{2+} .

Synthesis of Astaxanthin-Essential Metal Ions Complex

Astaxanthin stock solution, according to Polyakov, is made by weighting as much as 10 mg then dissolved in 100 mL ethanol (0.167 mM) and after that was diluted, which results in variations of astaxanthin concentration between Cu^{2+} (1 μM , 2 μM , and 3 μM) and Zn^{2+} (2 μM and 20 μM).¹⁴ These variations are based on the characteristics of each metal. For stock solution, Cu^{2+} is made with $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ (2.56 mg) in 15 ml aquades then used as a working solution with a concentration of 1 μM . As for Zn^{2+} made by dissolving ZnCl_2 (4,32 gr) in 100 ml aquades, then diluted into working solutions with concentrations of 20 μM and 40 μM . In the formation of complex, treatment given includes stirrer time (5 and 15 minutes), temperature (20°C, 37 °C and 78 °C) and complex ratio of (astaxanthin : metal ions) such as for Cu^{2+} (1:1, 2:1 and 3:1) then Zn^{2+} (1:1, 1:2, 0,1:1 and 0,1:2). Temperature conditioning is carried out on a water bath placed on top of magnetic stirrers (Cimarec Thermom SP-88850105) at a speed of 110 rpm.

UV-Vis Spectroscopic Characterization

Compound profiling using a UV-Vis Spectrophotometer (Genesys 10 UV). Each sample was reviewed for UV-Vis spectrum profile with a wavelength of 190-790 nm, using a turbo scan survey mode (accuracy of 5 nm). The spectrophotometer calibration was performed using absolute ethanol.

FTIR Spectroscopic Characterization

Confirmation is carried out by FTIR because it can determine changes in functional groups in a biomolecule.¹⁵⁻¹⁸ Functional groups and confirmation of complex formation using FTIR (Shimadzu 8400S). Samples were previously freeze-dried to increase the translucency of the samples for FTIR analysis.¹⁹ Freeze dry of the sample is done by placing the sample into the freezer at a temperature of -25°C. The frozen sample was then transferred to a freeze dryer at -50°C with a vacuum pressure of 5 mpa. Freeze-dried samples are then added to the KBr compound before being put into the FTIR device. The data read from FTIR is a wavenumber in the range of 400-4000 cm^{-1} . Confirmation of the complex formation based on peaks at the wavenumber 600-400 cm^{-1} , which indicates the presence of metal bond groups in astaxanthin.

RESULTS AND DISCUSSION

UV-Vis Profile of Astaxanthin

In this study, astaxanthin was divided into two groups, which are complexed with Cu^{2+} and Zn^{2+} . Astaxanthin profiling to be complexed with Cu^{2+} begins with duration treatment, and the reaction temperature is given at concentrations of 1, 2, 3 and 20 μM at temperatures of 20°C, 37°C, and 78°C for 5 and 15 minutes. A temperature of 20°C is used to create a complex to determine how complex conditions are at room temperature. A temperature of 37°C is the body temperature as a reaction condition when the complex has been consumed. An increased temperature of 78°C is applied to achieve sufficient activation energy.²⁰ All astaxanthin samples have a peak at wavelengths of 475-480 nm. The astaxanthin code begins with Ax for astaxanthin and followed by the sample concentration (Ax(1) for 1 μM , Ax(2) for 2 μM , Ax(3) for 3 μM and Ax(4) for 20 μM , then the amount of stirrer time (5 minutes and 15 minutes) and ends with an incubation temperature value of 20, 37 or 78 (°C). Astaxanthin (1 μM) with Ax(1).15.20, Ax(1).5.20, dan Ax(1).5.37 codes have absorbance in the range of 0,256-0,277 Å. Ax(1).15.37, Ax(1).5.78, dan Ax(1).15.78 samples have higher absorbance of 0.313-0.457 Å. The astaxanthin 2 μM sample at various durations and reaction temperatures had a slightly different absorbance peak of 0.370,

0.371, 0.391, 0.417, 0.425 Å and at Ax(2).15.78 had a high-rise absorbance peak of 0.772 Å. The Ax(3).5.78 sample had the highest absorbance peak of 0.705 Å and Ax(3).15.78 at 0.247 Å resulting in the lowest absorbance (Fig.-1).

Fig.-1: UV-Vis Spectrum Profile of Astaxanthin at Various Reaction Duration and Temperature

Astaxanthin 20 μM (Fig.-1) with a reaction duration of 5 and 15 minutes at 20 and 37°C has a peak at a wavelength of 475-480 nm with an absorbance of 1.018-1.238 Å. Different results were shown for the reaction duration of 5 and 15 minutes at 78°C. Astaxanthin 20 μM with a reaction duration of 5 minutes at 78°C (Ax(4).5.78) produced a peak at a wavelength of 485 nm with an absorbance of 2.586 Å while for a reaction duration of 15 minutes at the same temperature, the absorbance increased to 3.004 Å at the same wavelength. Ax(4).15.78 appears to peak in the UV spectrum with a wavelength of 220 nm absorbance of 1.321 Å. From Fig.-1, it can be concluded that astaxanthin with higher concentration has higher absorbance. Also, the reaction temperature of 20°C and 37°C produced different spectral profiles, but it was not significant at the reaction time of 5 minutes and 15 minutes at both temperatures. In contrast, the reaction temperature of 78°C resulted in a significant increase in absorbance compared to the reaction temperatures of 20°C and 37°C, especially at the reaction duration of 15 minutes. Thoen stated that astaxanthin's peak is at a wavelength of 480 nm. The UV-Vis spectrum profile of the astaxanthin sample shows peaks in both UV and visible areas.²¹ The peak astaxanthin in the UV spectrum shows cis-astaxanthin, while the peaks in the visible area show the structure of the trans-astaxanthin structure.²² The UV-Vis spectrum profile of the astaxanthin sample at a typical treatment temperature of 78°C is shown by the absorbance of the peak, which is much different from the treatment at 20°C and 37°C. It is possible because of the isomerization of the astaxanthin structure. At a temperature of 70 °C, astaxanthin undergoes isomerization from All-E-astaxanthin's structure to 9Z-astaxanthin and 13Z-astaxanthin. E-astaxanthin and Z-astaxanthin's structure is a structure based on the nomenclature E-Z notation.²³ E stands for entgegen, which means the opposite, while z stands for zusammen that means together. E notation is a trans structure, named after the astaxanthin structure, which has a pair of OH and O groups located on opposite sides of the polyene chain. In comparison, z notation is a cis structure, namely for astaxanthin with OH and O groups located on the same side.

UV-Vis Profile of Metal Ions (Cu^{2+} and Zn^{2+})

A UV-Vis spectrum profile characterized Cu^{2+} metal ion samples at various reaction durations and temperatures. The Cu^{2+} metal ion sample used had a concentration of 1 μM . Treatment of reaction duration and temperature used is the same as astaxanthin. The Cu^{2+} metal ion used is in the form of a $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ salt solution. All Cu^{2+} samples in various treatments of reaction duration and temperature had a peak at a wavelength of 205 nm (Fig.-2A). The absorbance peak of each Cu^{2+} sample also has a minimal difference, so that it can be said to be almost the same. The location of the peak and the absorbance is the

same because they have the same concentration. Besides, the temperature and reaction time does not affect the difference in the UV-Vis spectrum profile of Cu^{2+} metal ions. This metal ions sample peak lies at a wavelength of 205 nm, which is the UV region. The characterization of the UV-Vis spectrum profile of $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ was also carried out by Ong, $\text{CuCl}_2 \cdot 2\text{H}_2\text{O}$ which was stirred for 14 hours at room temperature had a peak at 235 nm.²⁴

Fig.-2: UV-Vis Spectrum Profile of (A) Cu^{2+} and (B) Zn^{2+} at Various Reaction Duration and Temperature

Furthermore, the Zn^{2+} metal ion was divided into two concentrations, which are code A for 20 μM and D for 40 μM (Fig.-2B). The UV-Vis Zn^{2+} spectrum profile with a concentration of 20 μM with a reaction duration of 5 and 15 minutes at 20, 37, or 78 ($^{\circ}\text{C}$) has similarities to the spectrum profile. It can be seen from the peak at a wavelength of 195 nm with an absorbance of 0.472-0.514 Å. The UV-Vis spectrum profile of Zn^{2+} with a concentration of 40 μM has the character of a peak at 195 nm for the reaction duration of 5 and 15 minutes at a temperature of 20 and 37 $^{\circ}\text{C}$ with an absorbance of 0.634-0.657 Å. The reaction temperature treatment of 78 $^{\circ}\text{C}$ showed different results with the treatment at 20 and 37 $^{\circ}\text{C}$. The peaks were at a wavelength of 200 nm for both reaction durations with absorbances for reaction durations of 5 and 15 minutes of 0.752 and 0.725 Å, respectively. Based on the UV-Vis profile of the single Zn^{2+} compound, it can be determined that the Zn^{2+} solution with a concentration of 20 μM has a peak at a wavelength of 195 nm at all duration and temperature of the reaction and the absorbance difference is insignificant. Meanwhile, a single Zn^{2+} compound with a concentration of 40 μM has a peak at a wavelength of 195 nm, which occurs only at a reaction temperature of 20 $^{\circ}\text{C}$ and 37 $^{\circ}\text{C}$ for both reaction durations. Whereas at a temperature of 78 $^{\circ}\text{C}$, it has a peak at a wavelength of 200 nm in both duration of

the reaction with an absorbance of 0.725-0.752 Å. *Trivedi* stated that the maximum absorbance of ZnCl_2 is in the UV region at a wavelength of 197.6 nm. The shifting of Zn^{2+} spectrum profile is possible due to the influence of temperature and different solvents from the literature.²⁵

UV-Vis Profile of Astaxanthin-Metal Ions (Cu^{2+} and Zn^{2+}) Complex

Astaxanthin- Cu^{2+} (Fig.-3), which has been reacted at various durations and temperatures is then analyzed by complex formation based on UV-vis spectrum profiles. The concentration of astaxanthin used is 1, 2, and 3 μM while Cu^{2+} with a concentration of 1 μM . The code for astaxanthin- Cu^{2+} complex begins with the ratio code (Astaxanthin : Cu^{2+}) i.e. K1 (1:1), K2 (2:1) and K3 (3:1). After that followed by a stirrer time of 5 or 15 minutes and incubation temperatures of 20, 37 or 78 ($^{\circ}\text{C}$). Complex profiles with code K1.5.20 have an absorbance peak of 0.109 Å, which is the highest at a ratio of 1:1, while K2.5.37 at 0.141 Å, and K3.5.78 at 0.153 Å.

Fig.-3: UV-Vis Profile of the Astaxanthin- Cu^{2+} Complex (A) Ratio (1:1) , (B) Ratio (2:1), (C) Ratio (3:1)

The absorbance peak of the complex with 3:1 ratio is 0.153 Å indicates the highest absorbance peak compared to the 1:1 and 2:1 ratio complexes, so the greater the concentration of astaxanthin at the complex ratio then the absorbance peak is also more significant. The temperature treatment affects the immense absorbance value of the UV-Vis spectrum profile, the higher the reaction temperature, the absorbance peak value is also higher. From these results, four samples were selected that had the highest peak of any ratio group to continue its analysis using FTIR. The selected complexes are K1.5.20, K2.5.37, K3.5.20 and K3.5.78. The concentrations of astaxanthin used in forming astaxanthin- Zn^{2+} complex were 2 and 20 μM which would be complexed with metal ions Zn^{2+} with concentrations of 20 and 40 μM . Astaxanthin- Zn^{2+} complex was given a code starting with the complex ratio (Astaxanthin : Zn^{2+}), namely A (1:1), B (1:2), C (0.1:1) dan D (0.1:2) (Fig.-4). Furthermore, the stirrer times are 5 and 15 minutes, and then the incubation temperatures are 20, 37 or 78 ($^{\circ}\text{C}$). Complexes A, B, C, and D generally show a similar pattern of UV-Vis spectrum profiles, although the ratio of astaxanthin-metal concentrations in each sample is different. The astaxanthin- Zn^{2+} complex which forms a bioinorganic complex based on the UV-Vis spectrum are A.15.78 ratio (1:1), B.15.78 ratio (1:2), C.15.78 ratio (0.1:1), and D.5.78 ratio (0.1:2). Samples were selected based on the highest absorption peak, and this selection is the same as the previous astaxanthin- Cu^{2+} . Overall, there is a peak shift when compared between single astaxanthin and complexes A, B, C or D. This condition is referred to as a blue shift. Hypsochromic shift or blue shift is a

shift in the maximum absorbance towards a shorter wavelength.²⁶ *Simonyi* stated that the occurrence of hypsochromic shift indicates tight association which is often referred to as H-type or card-pack aggregate.²⁷ *Shuaith* states that the H-type aggregate can be caused by interference from the absorption of copper chloride.²⁰ It is due to the formation of a complex between the metal chlorides molecule astaxanthin at high temperatures. From this explanation, the peak shift towards hypsochromic occurs in astaxanthin- Zn^{2+} complex due to the presence of ZnCl_2 which is a metal chlorides salt complexed with astaxanthin. The hypsochromic shift can also occur due to atomic conjugation removal and solvent polarity.²⁶ In general, if a molecule absorbs UV or Vis radiation energy, the molecule is called a chromophore.²⁸

Fig.-4: UV-Vis Profile of the Astaxanthin- Zn^{2+} Complex (A) Ratio (1:1), (B) Ratio (1:2), (C) Ratio (0.1:1), (D) Ratio (0.1:2)

Like other carotenoid groups, the astaxanthin molecule consists of polyene carbon chains, namely carbon chains that are single and double-bonded (conjugated) and arranged alternately. This backbone structure is the chromophore of astaxanthin.²⁹ During the reading of the UV-Vis spectrum, the sample is exposed using UV-Vis waves, and the chromophore receives the energy appropriate for electron excitation. Some of the energy will be absorbed as π -bonding or n- electrons to be transferred to higher energy orbitals. Electron transitions usually occur from bonding or nonbonding to antibonding which is marked with an asterisk symbol or (*) above the transition symbol.³⁰ *Yadav* stated that hypsochromic occurs when the electron transition occurs from $n \rightarrow \pi^*$ and $\pi \rightarrow \pi^*$. The $n \rightarrow \pi^*$ transition is an R-Band transition while the $\pi \rightarrow \pi^*$ transition is a K-Band transition.²⁶ The maximum absorbance of astaxanthin occurs at a wavelength of 478 nm in ethanol solvent, at that wavelength, there is a transition from the conjugated molecule orbital $\pi \rightarrow \pi^*$ or called the K-Band.²⁰

FTIR Profile of Astaxanthin-Metal Ions (Cu^{2+} and Zn^{2+}) Complex

FTIR profile shows the functional groups and bonds present in a compound. The bond vibration between molecules produces peaks. The FTIR profile of a single Cu^{2+} has 4 peaks at wavenumber 452.07; 548.51; 1613.14; and 3364.39 cm^{-1} , while astaxanthin which shows a peak at wave number 1601.57 cm^{-1} shows group vibrations, namely (C=O), 718.23 cm^{-1} (=C-H), 3426.1 cm^{-1} (O-H), 1464.63 cm^{-1} (C=C) (Fig.-5A).

The FTIR profile of astaxanthin in this study is similar to that conducted by *Shuaith*.²⁰ The FTIR profile of a complex compound has a different shape from its single compound and has a form that is a combination of a single astaxanthin profile and a single Cu. The profile of astaxanthin-Cu²⁺ based on FTIR results has shown the formation of a complex because there are functional groups that comprise it, namely the group (C = O), (OH), (C = C), and (= CH) at wavenumber 1830-1530 cm⁻¹, 3550-3300 cm⁻¹, 1600-1430 cm⁻¹, 3100-3000 cm⁻¹ respectively (Table-1).

Fig.-5: FTIR Profile of selected (A) Astaxanthin-Cu²⁺ and (B) Astaxanthin-Zn²⁺ Complex

The Cu bond with astaxanthin is marked with a peak at the wave number 600-400 cm⁻¹ indicating the group (Cu-O). The most optimal formulation of astaxanthin-Cu²⁺ complex is a 3: 1 ratio at 78°C for 5 minutes because it has a UV-Vis spectrum profile which shows a peak in the visible area with the highest absorbance compared to other samples, namely 0.153.

FTIR also confirmed this complex that it had indeed been formed due to the presence of Cu²⁺ metal bonds with astaxanthin's oxygen, which was marked by a peak at wavenumber 517.65 cm⁻¹. The functional group characters of astaxanthin are (C-H), (C=C), (C=O), and (O-H).³² The functional groups (C-H) in astaxanthin are in wave numbers 666.16 and 1464.63 cm⁻¹. The functional groups (C=C) are in the wavenumbers 970.89 and 1601.57 cm⁻¹. The functional groups (C=O) are at wavenumbers 1653.64 cm⁻¹ dan (O-H) and (O-H) are at wavenumbers 3032.65 cm⁻¹ and 3491.68 cm⁻¹. Also, bonds (C-O) and (C-N) were found in the astaxanthin sample. The functional group (C-O) is in wave number 1073.54; 1129,33;

1200,40; 1281.41 and 1391.34 cm^{-1} . The functional group (C-N) is found at wave number 1034.54 cm^{-1} . Zn^{2+} has the characters of functional groups (M-X), (M-H), (M = O), and (O-H) (Fig.-5B).

Table-1: Identification of Functional Groups in Single Compounds and Selected Astaxanthin-metal Ions Complexes

Functional Group	Literature*	Astaxanthin	Zn^{2+}	A.15.78	B.15.78	C.15.78	D.5.78	Cu^{2+}	K1.5.20	K2.5.37	K3.5.20	K3.5.78
C-H	700±20	666.16		685.45								
	1465	1464.63		1468.49			1468.49					
	3000-2840						2934.29					
=C-H	1000-600**								895.67	988.25	835.88	685.45
C=O	1654*** 1830-1530**	1653.64							1626.64	1630.50	1624.71	1626.64
C=C	980-960	970.89										
	1650-1600	1601.57		1622.79	1622.79	1626.64	1622.79	1613.14				
	1600-1430**								1551.42	1454.99	1516.71	1516.71
C-O	1085-1050	1073.54										
	1150-1085	1129.33										
	1275-1200	1200.40		1213.9								
	1310-1250	1281.41		1289.12								
	1420-1330	1391.34										
	1210-1163			1177.26								
O-H	1420-1330					1391.34						
	3200-2700	3032.65										
	3550-3200	3491.68	3518.68	3547.61	3545.68	3526.4	3545.68	3364.39				
	3700-3584		3590.04									
	3550-3300**								3449.25	3441.53	3451.18	3451.18
M=O**	1010-850**		1003.68									
C-N	1250-1020	1034.54		1038.3								
N-O	1550-1500			1522.49		1520.56						
M-X**	750-100**		504.15	523.44			471.36					
M-H**	2250-1700**		1609.28 1935.23									
Cu-O**	600-400**							452.07 548.51	546.58	550.44	475.22	517.65

*Merck, **Stuart, ***Shuaith.^{31, 32, 20}

Stuart stated that metals coordinated with atoms such as halogens, H, N, or O form infrared characteristics in the fingerprint region.³² The functional group (M-X) or metal-halogen is in the wavenumber 504.15 cm^{-1} . The functional groups (M-H) are in wave numbers 1609.28 and 1935.23 cm^{-1} . The functional groups (M=O) are at wave number 1003.68 cm^{-1} and (O-H) are at wavenumbers 3518.68 and 3590.04 cm^{-1} . Metal bonds with oxygen and hydrogen because zinc chloride has deliquescent properties (becomes liquid) when there is H_2O in its anhydrous form.³³ It is also supported because the group IIIB atoms have a higher cation potential, so they have a higher hydration tendency. The ability to hydrate is influenced by the ionization energy, affinity, and electronegativity of the atom. In the periodic table, the smallest atomic number in a group has the strongest properties, so that group IIIB produces hydration conditions ($\text{Zn}^{2+} > \text{Cd}^{2+} > \text{Hg}^{2+}$). When ZnCl_2 is exposed to water molecules, the molecular formula changes to $(\text{ZnCl}_2 [\text{H}_2\text{O}]_n)$; where $n = 1, 1.5, 2.5, 3, \text{ dan } 4$. The n shows the number of water molecules bound by 1 ZnCl_2 molecule.²²

A.15.78 complex with ratio (1:1) has the most atomic functional groups compared to complex samples B.15.78, C.15.78, and D.5.78. A.15.78 complex has an atomic functional group (C-H) at wavenumber 685.45 cm^{-1} , (C=C) at 1622.79 cm^{-1} , (C-O) at 1213.9 ; 1289.12 ; and 1177.26 cm^{-1} , (O-H) at 3547 cm^{-1} , (C-N) at 1038.3 cm^{-1} , (N-O) at 1522.49 cm^{-1} , the last is (M-X) at 523.44 cm^{-1} (Table-1). B.15.78 sample has (C=C), (C-H), and (O-H) atomic functional groups, which are respectively in wavenumbers 1622.79 cm^{-1} , 1468.49 cm^{-1} , and 3545.68 cm^{-1} , respectively. C.15.78 sample has functional groups (C=C), (O-H), and (N-O) with wavenumbers 1626.64 cm^{-1} , 1391.34 cm^{-1} , 5326.4 cm^{-1} and 1520 cm^{-1} . D.5.78 complex has functional groups such as (C-H), (C=C), (O-H), and (M-X) at 1468.49 cm^{-1} and 2934.29 cm^{-1} , 162.79 cm^{-1} , 3545.68 cm^{-1} , and 471.36 cm^{-1} , respectively. From the atomic functional groups above, it is clear that the tendency for all complex samples to be similar to the ZnCl_2 FTIR profile, but the complex A ratio sample (1:1) still has astaxanthin profile because of the large number of functional groups present in the fingerprint area. *Alfanar & Notario* stated that the interaction between astaxanthin- Zn^{2+} is in the fingerprint area at 533 cm^{-1} .³⁴ A.15.78 complex with ratio (1:1) contained a functional group between the metal-oxygen at wavenumber 523.44 cm^{-1} so that it was confirmed to form the bioinorganic chemical complex of astaxanthin- Zn^{2+} .

CONCLUSION

The most optimal complex formulation to form astaxanthin- Cu^{2+} complex is a 3: 1 ratio with a reaction duration of 5 minutes at 78°C . This formulation is based on the UV-Vis spectrum profile, which has a high peak absorbance of 0.153 at a wavelength of 460-470 nm and is a solid bond so that this complex becomes the optimal formulation in forming a complex; besides that, it has been confirmed that the complex is formed by FTIR test. The physicochemical character of the complex tested using FTIR has a profile that shows the functional groups of the astaxanthin structure in the form of (C=O), (C-H), (O-H), (C=C), and the presence of Cu^{2+} bonds with astaxanthin oxygen to form groups (Cu-O) at wavenumber $600\text{-}400\text{ cm}^{-1}$. Meanwhile, in the astaxanthin- Zn^{2+} complex, it is known that the temperature and duration of the reaction affect complex formation. The astaxanthin- Zn^{2+} complex was formed at a reaction temperature of 78°C with a reaction duration of 15 minutes, marked by a peak as the astaxanthin character at a wavelength of about 480 nm. Based on the FTIR profile, the astaxanthin- Zn^{2+} complex's formation is characterized by the vibration of metal and oxygen bonds from astaxanthin as a ligand in the fingerprint area at 523.44 cm^{-1} . Further laboratory experiments are now undergoing in our laboratory.

ACKNOWLEDGEMENT

This research was supported by the Ministry of Research, Technology and Higher Education of Indonesia through the scheme of PMDSU.

REFERENCES

1. G. Pizzino, N. Irrera, M. Cucinotta, G. Pallio, F. Mannino, V. Arcoraci, F. Squadrito, D. Altavilla and A. Bitto, *Oxidative Medicine and Cellular Longevity*, **8416763** (2017), DOI:10.1155/2017/8416763
2. G. Hussein, U. Sankawa, H. Goto, K. Matsumoto, H. Watanabe, *Journal of Natural Products*, **3**, 443(2006), DOI:10.1021/np050354+
3. A. Phaniendra, D.B. Jestadi, L. Periyasamy, *Indian Journal of Clinical Biochemistry*, **1**, 11(2015), DOI:10.1007/s12291-014-0446-0
4. V. Lobo, A. Patil, A. Phatak, & N. Chandra, *Pharmacognosy Reviews*, **8**, 118(2010), DOI:10.4103/0973-7847.70902
5. L. Ekpe, K. Inaku & V. Ekpe, *Journal of Molecular Pathophysiology*, **1**, 1(2018), DOI:10.5455/jmp.20180627120817
6. J. Zhang, Z. Sun, P. Sun, T. Chena and F. Chen, *Food & Function*, **5**, 413(2014), DOI:10.1039/c3fo60607d
7. S. Wibowo, S. Widyarti, A. Sabarudin, D.W. Soeatmadji and S.B. Sumitro, *Asian Journal of Pharmaceutical and Clinical Research*, **12**, 276(2019), DOI:10.22159/ajpcr.2019.v12i9.34617
8. E. Hernández-Marin, A. Barbosa and A. Martínez, *Molecules*, **17**, 1039(2012), DOI:10.3390/molecules17011039

9. S. Wibowo, S. Widyarti, A. Sabarudin, D.W. Soeatmadji and S.B. Sumitro, *Heliyon*, **7**, e06548 (2021), [DOI:10.1016/j.heliyon.2021.e06548](https://doi.org/10.1016/j.heliyon.2021.e06548)
10. K.H. Reddy, Bioinorganic Chemistry, New Age International, Ltd., New Delhi (2003)
11. L. Zhao, F. Chen, G. Zhao, Z. Wang, X. Liao and X. Hu, *Journal of Agricultural and Food Chemistry*, **24**, 9620(2005), [DOI:10.1021/jf0517750](https://doi.org/10.1021/jf0517750)
12. Z. Liu, Y. M. Ng, P. J. Tiong, R. A. A. Talip, N. Jasin, V. Y. M. Jong and M. G. Tay, *International Journal of Inorganic Chemistry*, **7520640**, 1(2017), [DOI:10.1155/2017/7520640](https://doi.org/10.1155/2017/7520640)
13. P. Chellan and P. J. Sadler, *Philosophical Transactions of the Royal Society A.*, **2037**, 20140182 (2015), [DOI:10.1098/rsta.2014.0182](https://doi.org/10.1098/rsta.2014.0182)
14. N.E. Polyakov, A. L. Focsan, M. K. Bowman and L. D. Kispert, *The Journal of Physical Chemistry. B.* **114**, 16968(2010), [DOI:10.1021/jp109039v](https://doi.org/10.1021/jp109039v)
15. F. Faghihzadeh, N. M. Anaya, L. A. Schiffman and V. Oyaneded-Craver, *Nanotechnology for Environmental Engineering*, **1**, 1(2016), [DOI:10.1007/s41204-016-0001-8](https://doi.org/10.1007/s41204-016-0001-8)
16. K. Sivaram, M.C. Rao, G. Giridhar, M. Tejaswi, B.T.P. Madhav, V.G.K.M. Pisipati and R.K.N.R. Manepalli, *Rasayan Journal of Chemistry*, **10**, 16(2017), [DOI:10.7324/RJC.2017.1011539](https://doi.org/10.7324/RJC.2017.1011539)
17. N. Karikalan, *Rasayan Journal of Chemistry*, **11**, 1451(2018), [DOI:10.31788/RJC.2018.1143068](https://doi.org/10.31788/RJC.2018.1143068)
18. M.V.L. Kumari, A. Kaviarasi, A.R. Prabakaran and A. Anandavadeivel, *Rasayan Journal of Chemistry*, **12**, 1688(2019), [DOI:10.31788/RJC.2019.1245384](https://doi.org/10.31788/RJC.2019.1245384)
19. W.F. Wolkers and H. Oldenhof, *Spectroscopy*, **19**, 89(2005), [DOI:10.1155/2005/216509](https://doi.org/10.1155/2005/216509)
20. N. Shuaith, Thesis, Department of Science, Synthesis and Characterisation of Novel Astaxanthin Metal Complexes, Letterkenney Institute of Technology, Irlandia (2015)
21. H.H. Thoen, G. Johnsen and J. Berge, *Polar Biology*, **34**, 83(2011), [DOI:10.1007/s00300-010-0861-5](https://doi.org/10.1007/s00300-010-0861-5)
22. J.P. Yuan, & F. Chen, *Journal of Agricultural and Food Chemistry*, **47**, 3656(1999), [DOI:10.1021/jf981319u](https://doi.org/10.1021/jf981319u)
23. C. Yang, L. Zhang, H. Zhang, Q. Sun, R. Liu, J. Li, L. Wu and R. Tsao, *Journal of Agricultural and Food Chemistry*, **4**, 818(2017), [DOI:10.1021/acs.jafc.6b04962](https://doi.org/10.1021/acs.jafc.6b04962)
24. H.R. Ong, M.M.R. Khan, R. Ramli, Y. Du, S. Xi and R.M. Yunus, *RSC Advances*, **31**, 24544(2015), [DOI:10.1039/C4RA16919K](https://doi.org/10.1039/C4RA16919K)
25. M.K. Trivedi, K. K. Sethi, P. Panada and S. Jana, *International Journal of Pharmaceutical Investigation*, **1**, 33(2017), [DOI:10.4103/jphi.JPHI217](https://doi.org/10.4103/jphi.JPHI217)
26. L.D.S. Yadav, Organic Spectroscopy, Springer Science, India (2005)
27. S. Simonyi, Z. Bikadi, F. Zsila and J. Deli, *Chirality*, **15**, 680(2003), [DOI:10.1002/chir.10282](https://doi.org/10.1002/chir.10282)
28. A.K. Pahari, & B. S. Chauhan, Engineering Chemistry, Laxmi Pub., New Delhi (2006)
29. F. Shahidi, & C. Barrow, Marine Nutraceuticals and Functional Foods, CRC Press, New York (2008)
30. B.M. Ham, & A. Maham, Analytical Chemistry, John Wiley & Sons Inc., New Jersey (2016)
31. Merck. IR Spectrum Table & Chart, <https://www.sigmaaldrich.com>
32. B.H. Stuart, Infrared Spectroscopy: Fundamentals and Applications, Wiley, New Jersey, (2004)
33. J.J. Zuckerman, Inorganic Reactions and Methods, Wiley-VCH, New Jersey (1991)
34. R. Alfenaar and D. Notario, *Jurnal Kimia dan Kemasan*, **2**, 88(2019), [DOI:10.24817/jkk.v4i1i2.3366](https://doi.org/10.24817/jkk.v4i1i2.3366)
[RJC-6209/2020]