ISOLATION, CHARACTERIZATION AND ANTIDIABETIC ACTIVITY OF PHENOLIC COMPOUNDS ISOLATED FROM *Macaranga magna* TURRILL

Minarti^{1,2}, A.H. Cahyana¹ and A. Darmawan^{2,✉}

¹Department of Chemistry, University of Indonesia, Depok, West Java, Indonesia, 16424

²Research Center for Chemistry, Indonesian Institute of Sciences, Tangerang Selatan, Banten, Indonesia, 15314

✉Corresponding Author: ahmaddarmawan2013@gmail.com

ABSTRACT

Southeast Sulawesi is an Indonesian region that has a large amount of biodiversity resources. As part of our continuous drug discovery project, three phenolic compounds, namely (1) scopoletin, (2) apigenin, and (3) gallic acid were successfully isolated from the leaf extract of *Macaranga magna* Turrill. The isolation and purification processes were conducted by serial chromatography, and their chemical structures were elucidated on the basis of spectroscopic data (FT-IR, FT-NMR and LC-ESI-MS), as well as through data comparison with the related body of literature. Their potential antidiabetic activity was evaluated against α -glucosidase enzymes. Scopoletin showed the highest antidiabetic activity compared to apigenin and gallic acid, with IC₅₀ values of 91.7, 28.3, and 198.5 μ g/mL, respectively.

Keywords: Scopoletin, Apigenin, Gallic Acid, *Macaranga magna* Turrill, α -glucosidase, Antidiabetic

RASAYAN J. Chem., Vol. 14, No.4, 2021

INTRODUCTION

As a megabiodiversity country, Indonesia has diverse biodiversity, especially in terms of plant species. However, there remain many plant species that have not yet been explored and utilized.¹ *Macaranga* (Euphorbiaceae) known in Indonesia as 'mahang-mahangan,' is a large genus of plant with over 300 species and is widely distributed to Africa, Madagascar, Asia, Australia, and the Pacific islands.²⁻⁵ *Macaranga* species have an important role in traditional medicine systems to treat certain diseases due to their bioactive compounds such as *Macaranga gigantea* and *M. triloba*, which have been used to treat fungal infections, leaf decoction, stomachaches^{2,6}, *Macaranga hypoleuca* can be used as a febrifuge, expectorant and antispasmodic, and has antioxidant and anti-bacterial properties.⁶ *M. denticulata* has been used to treat icteric hepatitis⁷; *M. indica* has been used to treat anemia and tumors in India and Bangladesh^{2,6}; *M. peltata* has been used to treat relief from venereal sores and has antioxidant, antimicrobial and cytotoxicity activity⁷; *M. hurifolia* Beille has been used to treat cough and diabetes, to relieve oedema in pregnant women and as a purgative⁸; and *M. barteri* Müll.-Arg has been used as an antioxidant, anti-inflammatory and anti-hyperalgesic, and has also been reported to relieve persistent cough, bronchitis, stomatitis, gastric ulcers, dysentery, swelling and arthritis.⁹ Previous studies showed that the *Macaranga* genus is a source for various phenolic compounds such as flavonoids^{2,3,9-20,25} and stilbenes.^{8,15,21-23}

Previous phytochemical studies on the *Macaranga* genus have shown that this genus is a source for various and unique bioactive compounds such as apigenin, apigenin-8-C-glycoside and scopoletin from *M. gigantifolia*^{2,7,16}, 6,8-diprenyl-4'-methyl-naringenin, 4'-deprenyl-4-methoxymappain, 4'-methyl-8-prenyltaxifolin from *M. balansae*²⁴, macatanarin D, schweinfurthin H, vedelianin, schweinfurthin F, schweinfurthin, 4'-deprenyl-mappain from *M. tanarius*^{5,22} as a cytotoxic active compound, glyasperin A from *M. gigantea* and *M. pruinosa* as an antioxidant.²⁵⁻²⁶ Nymphaeol C, solophenol D, nymphaeol A, and

nymphaeol B were isolated from the ethyl acetate fraction of *M. tanarius* leaves as antiplasmodial active compounds. Varyophyllene, neophytadiene, α -humulene, phytol, ethyl-ester hexadecanoic acid¹⁸, and nerolidol from *M. barteri* were isolated as antimicrobes. Macatrichocarpins A–D, oxymacatrichocarpin C, and isomacatrichocarpin from *M. tricocharpa* were isolated as antibacterial agents.²⁷

Diabetes mellitus is a metabolic disease caused by a decrease in insulin secretion or insulin sensitivity. Indonesia has become one of the highest diabetes prevalent countries in the world, and it has been estimated that 21.3 million people will have diabetes by 2030.²⁷ Inhibiting the activity of the enzyme will delay the absorption of the glucose, and the levels of blood glucose will decrease as a result.²⁹ Due to the prevalence of diabetes, the exploration and utilization of natural products as resources to treat diabetes is the primary aim of this research.

Macaranga magna Turrill belongs to the *Macaranga* genus, which has not been explored intensively. As far as our knowledge, the phytochemical study of *M. magna* was only reported by Minarti et al.²⁹ As part of our continued studies of drug discovery projects from Indonesian natural products resources, we explored the secondary metabolites compound content of the leaves of *M. magna* Turrill and evaluated their biological activity as α -glucosidase inhibitory activity.

EXPERIMENTAL

Material and Methods

Macaranga magna Turrill. Leaves were collected from Mekongga Forest, District Kolaka, Southeast Sulawesi. Plant species identification was performed by Herbarium Bogoriense, Research Center for Biology, Indonesian Institute of Sciences, Indonesia. Distilled technical grade organic solvents of *n*-hexane, ethyl acetate, and methanol were used for extraction, fractionation, isolation and purification. Silica gel 60 (0.063-0.200 mm) (Merck, 1.07734.1000) and TLC plate (Silica gel 60 F254) (Merck, 1.05554.0007) were used for isolation and chemical compounds spot identification. Instrumentation analysis of the isolated compounds used chloroform-*d* (Merck, 1.02450.0100), methanol-*d*₄ (Cambridge Isotope Laboratories, Inc., DLM-24-50), and acetone-*d*₆ (Cambridge Isotope Laboratories, Inc., DLM-9-25) for NMR (JEOL NMR ECZ500R 500 MHz) analysis; and methanol hypergrade (Merck, 1.06035.2500) and acetonitrile hypergrade (Merck, 1.00039.2500) for LC-MS/MS (Waters, Xevo G2-XS QTOF) analysis. A spectrophotometer (Agilent Technologies, Cary 60 UV-Vis) was used for maximum wavelength analysis, FTIR (Shimadzu, IR Prestige-21) for the determination of chemical groups, and a melting point meter (Fisher Scientific and KRUSS, M5000). *p*-NPG (Wako 351-17473, Osaka, Japan) and α -glucosidase enzyme from yeast (Wako 46511003, Osaka, Japan) were used for the determination of α -Glucosidase inhibitory activity.

Extraction and Isolation

Dried and powdered *M. magna* Turrill leaves were macerated with distilled methanol to obtain methanol extract. The methanol extract was further fractionated with *n*-hexane, ethyl acetate, acetone and *n*-butanol. About 4.2 g of ethyl acetate fraction was subjected to silica gel column chromatography and eluted using mobile phase gradient system solvent, which started from *n*-hexane, ethyl acetate and methanol, to obtain 11 sub-fractions. Sub-fraction 4 (SFea-4) and SFea-7 were subjected to silica gel column chromatography using a gradient system solvent from *n*-hexane and ethyl acetate to methanol, which led to the isolation of compounds **1** and **2**. The isolation and purification process of 9.9 g of acetone fraction used silica gel column chromatography with the same system solvent of the mobile phase, which led to the isolation of compound **3**, obtained from the recrystallization process using chloroform: methanol (1:1) of sub-fraction 7 (SFac-7).

Compound 1: pale yellow powder, mp. 202-204 °C, UV λ_{max}^{MeOH} 298, 345. FTIR (KBr) ν_{max} (cm⁻¹) 3228, 2924, 1711, 1610, 1176. LC-ESI-MS *m/z* 193 [M+H]⁺. ¹H-NMR (500 MHz, in Acetone-*d*₆) (δ ppm) 6.17 (1H, *d*, *J* = 9.5 Hz, H-2), 6.78 (1H, *s*, H-8), 7.18 (1H, *s*, H-5), 7.83 (1H, *d*, *J* = 9.5, H-3), 3.89 (3H, *s*, -OCH₃). ¹³C-NMR (125 MHz, in Acetone-*d*₆) (δ ppm) 56.7 (-OCH₃), 112.1 (C-10), 109.9 (C-5), 146.0 (C-6), 151.9 (C-7), 103.7 (C-8), 151.2 (C-9), 113.3 (C-3), 144.7 (C-4).

Compound 2: yellowish-green powder, mp. 345-347 °C, UV λ_{max}^{MeOH} 273 & 338. FTIR (KBr) V_{max} (cm⁻¹) 3246, 1654, 1570. LC-ESI-MS m/z 271.2 [M+H]⁺. ¹H-NMR (DMSO, 500 MHz) (δ ppm) δ_H 6,75 (1H, s); 6,15 (1H, d, J = 1,95 Hz); 6,44 (1H, d, J = 1,95 Hz); 7,91 (2H, d, J = 9,05 Hz); 6,90 (2H, d, J = 9,05 Hz); 12,94 (-OH). ¹³C-NMR (125 MHz, in DMSO,) δ_C 163,7; 102,8; 181,7; 161,3; 99,0; 164,6; 94,1; 157,4; 103,6; 121,2; 128,5 (2C); 116,0 (2C), 161,5.

Compound 3: white powder, mp. 250-252 °C, UV λ_{max}^{MeOH} 272. FTIR (KBr) V_{max} (cm⁻¹) 3282, 1701, 1618, 1440, 1315, 1031, 862, 711. LC-ESI-MS m/z 170, ¹H-NMR (500 MHz, in methanol-*d*₄) 7.05 (2H, s). ¹³C-NMR (125 MHz, in methanol-*d*₄) 122.0 (C-1), 110.4 (C-2/C-6), 146.5 (C-3/C-5), 139.7 (C-4), and 170.5 (C-7).

α -Glucosidase Inhibitory Activity

α -glucosidase inhibitory activity of isolated compounds was conducted using α -Glucosidase inhibitory activity methods developed by Dewi³⁰. About 250 μ L of α -Glucosidase (0.124 unit/mL), 495 μ L of 0.1 M phosphate buffer (pH 7.0), and 5 μ L of various concentrations of samples in DMSO (50-200 μ g/mL) were pre-incubated at 37°C for 5 minutes. The addition of 250 μ L of 5 mM *p*-NPG was used for the initiation reaction. The mixtures were further incubated at 37 °C for 15 minutes. The reaction was stopped by the addition of 1 mL of 0.1 M Na₂CO₃. α -Glucosidase inhibitory activity was determined by the measuring of the *p*-NPG released using a spectrophotometer at λ 410 nm.

RESULTS AND DISCUSSION

Macaranga plants are known as a resource for unique biologically active compounds, especially phenolic compounds. In this study, we have successfully isolated three phenolic compounds from ethyl acetate fraction and acetone fraction, and one sterol compound from ethyl acetate fraction.

Compound **1** was obtained as a pale-yellow powder with a melting point of 202-204 °C. The UV-Visible spectrum of compound **1** showed typical maximum absorption bands of scopoletin at 298 and 345 nm, indicating that compound **1** has unsaturated constituents which form a conjugated bond to an aromatic group, as well as the presence of a phenolic group. FTIR spectroscopy data showed the presence of hydroxyl (O-H) in the phenol group at 3320 cm⁻¹, in the carbonyl group (-C=O) at 1,711 cm⁻¹, and in the aromatic group (-CH=CH) group at 1,610 cm⁻¹. The H-NMR spectrum showed the presence of four two aromatic protons at δ_H 6.78 (1H, *s*, H-8) and 7.18 (1H, *s*, H-5), two alkene protons at δ_H 6.17 (1H, *d*, J = 9.5 Hz, H-2) and 7.83 (1H, *d*, J = 9.5, H-3), and one methoxy group attached to the aromatic group at δ_H 3.89 ppm. ¹³C-NMR data combined with HMQC data in Table 1 shows that compound **1** has four quaternary carbons at δ_C 56.7 112.1 (C-10), 146.0 (C-6), 151.9 (C-7), and 151.2 (C-9); one carbonyl group (-C=O) at δ_C 161.3 (C-2); and one methoxy group at δ_C 56.7. Based on the HMQC/HMBC correlation data (Table-1 and Fig.-1), supported with LC-ESI-MS data, it can be confirmed that Compound **1** is scopoletin.

Table-1: 1D- and 2D-NMR Data of Compound **1**

No.	Compound 1		
	δ_H (ΣH , m, J Hz)	HMQC	HMBC
2.		161.3	
3.	6.17 (1H, <i>d</i> , 9.5)	113.3	C-10
4.	7.83 (1H, <i>d</i> , 9.5)	144.7	C-2, C-5, C-9
5.		112.1	
6.	7.18 (1H, <i>s</i>)	109.9	C-4, C-7, C-9
7.		146.0	
8.		151.9	
9.	6.78 (1H, <i>s</i>)	103.8	C-6, C-9, C-10
10.		151.2	
11.	3.89 (OCH ₃)	56.7

Fig.-1: (a) HMQC Correlations, (b) HMBC Correlations, (c) Chemical Structure of Compound 1 (scopoletin)

Compound **2** was obtained as a yellowish-green crystal with a melting point of 345-347 °C. UV-visible absorption spectra of Compound **2** in methanol exhibited two maximum peaks at λ 273 and 338 nm. Markham (1988) indicated that Compound **2** is a flavon type of flavonoid compound. The FTIR spectra showed broadband at 3246 cm^{-1} , indicating the presence of an intermolecular hydroxyl ($-\text{OH}$) stretch vibration. A vibration band at 1,654 cm^{-1} indicates the existence of a conjugation system between the carbonyl group ($-\text{C}=\text{O}$) with $-\text{C}=\text{C}-$ in C-2 and C-3 positions and a vibration band at 1,570 cm^{-1} indicates the presence of an aromatic double bond ($\text{C}=\text{C}$). ^1H -NMR spectra (Table-2) of Compound **2** indicates the existence of six aromatic protons at δ_{H} 6.15 (1H, d, $J = 1.95$ Hz, H-6), 6.44 (1H, d, $J = 1.95$ Hz, H-8), 7.91 (2H, d, $J = 9.05$ Hz, H-2'/H-6'), and 6.90 (2H, d, $J = 9.05$ Hz, H-3'/H-5'); one proton of heterocyclic pyrane at δ_{H} 6.75 (1H, s, H-3); and one proton of a hydrogen bond at δ_{H} 12.94 (1H, s). ^{13}C -NMR spectra (Table-2) indicates the existence of seven carbon methin at δ_{C} 116.0 (2C, C3'/C-5'), 128.5 (2C, C-2'/C-6'), 102.8 (C-3), and 99.0 (C-6), 94.1 (C-8); one carbonyl ($-\text{C}=\text{O}$) group at δ_{C} 181.7; and seven quaternary carbon at δ_{C} 103.6 (C-10), 121.2 (C-1'), 157.4 (C-9), 161.3 (C-5), 161.5 (C-4'), 163.7 (C-2), and 164.6 (C-7). The correlation between proton peak at δ_{H} 6.75 with a carbon atom at δ_{C} 163.7, 103.6, and 121.2 indicate that the proton is located at the H-3 position. This indicates that Compound **2** is a flavon. Based on proton-carbon correlations obtained from HMQC and HMBC spectrum data (Table-2 and Fig.-2), supported with UV-visible, FTIR, and LC-ESI-MS data, Compound **2** is the flavon compound, apigenin.

Table-2: 1D- and 2D-NMR Data of Compound **2**

No.	Compound 2		
	δ_{H} (ΣH , m, J Hz)	HMQC	HMBC
2.		163.7	
3.	6.75 (1H, s)	102.8	C-2, C-10, C-1'
4.		181.7 ($-\text{C}=\text{O}$)	
5.		161.3	
6.	6.15 (1H, d, 1.95)	99.0	C-5, C-7, C-8, C-10
7.		164.6	
8.	6.44 (1H, d, 1.95)	94.1	C-6, C-7, C-9, C-10
9.		157.4	
10.		103.6	
1'.		121.2	
2'/6'	7.91 (2H, d, 9.05)	128.5 (2C)	C-4'
3'/5'	6.90 (2H, d, 9.05)	116.0 (2C)	C-1', C-4'
4'		161.5	
-OH	12.94 (1H, s)		C-5

Fig.-2: (a) HMQC Correlations, (b) HMBC Correlations, (c) Chemical Structure of Compound 2 (apigenin)

Compound 3 was obtained as a white powder with a melting point of 250-252 °C. UV-visible absorption spectra of Compound 3 in methanol exhibited a maximum peak at λ 272, indicating the presence of a phenolic group. The IR spectra of Compound 3 indicate the existence of hydroxyl (-O-H) group at 3282 cm^{-1} , -C-H aromatic rings at 3024, 862, 711 cm^{-1} , -C=C-aromatic at 1618, 1539 cm^{-1} , carbonyl ester group (-COOH) at 1701 cm^{-1} , and -C-O ester group at 1315, 1031 cm^{-1} .

^1H -NMR spectra (Table 3) of Compound 3 indicate the existence of two symmetrical aromatic protons at δ_{H} 7.05 (2H, s). ^{13}C -NMR spectra (Table 3) showed the existence of two symmetrical carbon methin at δ_{C} 110.4 (2C, C3/C-7), three phenolic carbon atoms at δ_{C} 139.7 (C-5) and 146.5 (2C, C-4/C-6), one quaternary carbon atom at δ_{C} 122.0, and carbonyl (-C=O) at δ_{C} 170.5. Based on the proton-carbon correlations obtained from HMQC and HMBC spectrum data (Table-3 and Fig.-3), supported with UV-visible, FTIR, and LC-ESI-MS data, indicated Compound 3 is gallic acid.

Table-3: 1D- and 2D-NMR Data of Compound 3

No.	Compound 3		
	δ_{H} (ΣH , m, J Hz)	HMQC	HMBC
1.		170.5 (-C=O)	
2.		122.0	
3/7.	7.06 (2H, s)	110.4 (2C)	C-5, C-4/C-6, C-1
4/6.		146.5 (2C)	
5		139.7	

The anti-diabetic activity of isolated compounds was evaluated against α -glucosidase enzyme using the methods developed by Dewi³⁰, where quercetin, a plant polyphenol compound that has many biological activities such as anti-inflammatory, anticancer, antidiabetic, neuroprotective, anti-allergic and antioxidant, was used as positive control.³¹⁻³² The results show that all the isolated compounds have lower inhibitory activity against α -glucosidase enzyme, compared to quercetin (IC_{50} 5.05 $\mu\text{g/ml}$), with the inhibitory activity (IC_{50}) of Compounds 1-3 at 91.73, 28.28, and 197.22 $\mu\text{g/ml}$, respectively (Table-4).

CONCLUSION

Our continuous project of drug discovery from Indonesian natural product resources has led to the isolation of three secondary metabolite compounds, namely, (1) scopoletin, (2) apigenin, and (3) gallic acid, from the leaf of *Macaranga magna* Turrill. Their biological activity of α -glucosidase inhibitory activity was lower compared to quercetin as a positive control (IC_{50} 5.05 $\mu\text{g/ml}$). Apigenin, a flavonoid

compound, and quercetin, showed the highest α -glucosidase inhibitory activity with an IC_{50} value of 28.28 $\mu\text{g/mL}$, compared to scopoletin and gallic acid with IC_{50} values of 91.73 and 197.22 $\mu\text{g/mL}$, respectively. This is first report of isolation of scopoletin, apigenin, and gallic acid from *Macaranga magna* Turrill.

Fig.-3: (a) HMQC Correlations, (b) HMBC Correlations, (c) Chemical Structure of Compound 3 (Gallic Acid)

Table-4: α -glucosidase Inhibitory Activity

No.	Sample	Absorbance	Concentrations ($\mu\text{g/mL}$)	Inhibition (%)	IC_{50} ($\mu\text{g/mL}$)
	Blank	0.1195			
1	Quercetin	0.0717 0.3501 0.6112 0.8540	10 7.5 5 2.5	94.00 70.69 48.83 28.51	5.05
2	Compound 1	0.1058 0.5105 0.8045 0.9300	200 100 50 25	91.14 57.26 32.65 22.14	91.73
3	Compound 2	0.1218 0.5940 1.0157 1.0472	50 25 12.5 10	89.80 50.27 14.97 12.33	28.28
4	Compound 3	0.5909 0.8525 0.9342 1.0169 1.0899	200 100 50 25 10	50.53 28.63 21.79 14.87 8.76	197.22

ACKNOWLEDGEMENT

The authors would like to thank the Indonesian Institute of Sciences (LIPI) for their 'by-research program'.

REFERENCES

- Yuliet, E.Y Sukandar, K. Budipramana, I.K. Adnyana, *Rasayan Journal of Chemistry*, **13(2)**, 826 (2020), <https://doi.org/10.31788/RJC.2020.1325607>
- S. Fajriah, Megawati, A. Darmawan, *The Journal of Tropical Life Science*, **6(1)**, 7(2016)

3. A. Khatun, M. Rahman, S. Kabir, *World Journal of Pharmaceutical Research*, **3(10)**, 172(2014)
4. J.J. Magadula, *Journal of Medicinal Plant Research*, **8(12)**, 489(2014), <https://doi.org/10.5897/JMPR2014.5396>
5. T.M.H. Doan, T.L. Nguyen, T.T.V. Trinh, V.N. Vu, T.D. Phi, M. Litaudon, F. Roussi, V.M. Chau, V.C. Pham, *Journal of Analytical Methods in Chemistry*, **2019**, 1 (2019), <https://doi.org/10.1155/2019/2917032>
6. T.Y. Lim, Y.Y. Lim, C.M. Yule, *Journal of Tropical Forest Science*, **26(1)**, 134 (2014).
7. G. Primahana, A. Darmawan, *The Journal of Pure and Applied Chemistry Research*, **6(1)**, 22 (2017), <https://doi.org/10.21776/ub.jpacr.2017.006.01.298>
8. P. Segun, M. Gbadebo, M. Adebawale, C. Olufolabo, A.F. Jaiyesimi, *Acta Pharmaceutica Scientia*, **57(4)**, 93 (2019), <https://doi.org/10.23893/1307-2080.APS.05726>
9. E. Asante-Kwatia, Y. Jibira, A.Y. Mensah, D. Osei-Sarfoh, *Discovery Phytomedicine*, **6(3)**, 130 (2019), <https://doi.org/10.15562/phytomedicine.2019.104>
10. L.T.N. Vu, L.T. Anh, N.T. Cuc, N.X. Nhiem, B.H. Tai, P.V. Kiem, M. Litaudon, T.D. Thach, C.V. Minh, H.D.T. Mai, P.V. Cuong, *Natural Product Research*, **35(13)**, 2123(2019), <https://doi.org/10.1080/14786419.2019.1662007>
11. I. Zakaria, N. Ahmat, R. Ahmad, F.M. Jaafar, N. Ab. Ghani, S. Khamis, *World Applied Sciences Journal*, **9(9)**, 1003 (2010).
12. C.Y. Ragasa, J. de Jesus, *Research Journal of Pharmaceutical, Biological and Chemical Sciences*, **5(3)**, 701 (2014)
13. M. Tanjung, E.H. Hakim, D. Mujahidin, M. Hanafi, Y.M. Syah, *Journal of Asian Natural Products Research*, **11(11)**, 929 (2009), <https://doi.org/10.1080/10286020903302315>
14. D.S. Yang, J.G. Wei, W.B. Peng, S.M. Wang, C. Sun, Y.P. Yang, K.C. Liu, X.L. Li, *Fitoterapia*, **99**, 261(2014), <https://doi.org/10.1016/j.fitote.2014.10.003>
15. D.S. Yang, Z.L. Li, W.B. Peng, Y.P. Yang, X. Wang, K.C. Liu, X.L. Li, W.L. Xiao, *Fitoterapia*, **103**, 165(2015), <https://doi.org/10.1016/j.fitote.2015.04.001>
16. A. Darmawan, S. Kosela, L. B.S. Kardono, Y.M. Syah, *Journal of Applied Pharmaceutical Science*, **2(12)**, 175(2012), <https://doi.org/10.7324/JAPS.2012.21231>
17. M.S. Fareza, Y.M. Syah, D. Mujahidin, L.D. Juliawaty, I. Kurniasih, *Z. Naturforsch*, **69c**, 375 (2014), <https://doi.org/10.5560/ZNC.2014-0066>
18. E. Marlina, W. Astuti, K. Kosala, R. Hairani, T.S. Tjahjandarie, M. Tanjung, *Asian Journal of Chemistry*, **30(4)**, 795 (2018), <https://doi.org/10.14233/ajchem.2018.21004>
19. E. Marlina, R. Ruga, R. Hairani, T.S. Tjahjandarie, M. Tanjung, *Journal of Physics: Conference Series*, Conf. Series, **1277** (2019), 012014(2019), <https://doi.org/10.1088/1742-6596/1277/1/012014>
20. J.H. Lee, Y.G. Kim, S.K. Khadke, A. Yamano, J.T. Woo, J. Lee, *Phytomedicine*, **63**, (2019), <https://doi.org/10.1016/j.phymed.2019.153033>
21. B.J. Yoder, S. Cao, A. Norris, J.S. Miller, F. Ratovoson, J. Razafitsalama, R. Andriantsiferana, V.E. Rasamison, G.I. Kingston, *Journal of Natural Products*, **70(3)**, 342(2007), <https://doi.org/10.1021/np060484y>
22. T. Péresse, G. Jézéquel, P.M. Allard, V.C. Pham, T.M.H. Doan, F. Blanchard, J. Bignon, H. Lévaïque, J.L. Wolfender, M. Litaudon, F. Roussi. *Journal of Natural Products*, **80(10)**, 2684 (2017), <https://doi.org/10.1021/acs.jnatprod.7b00409>
23. J.A. Beutler, J. Jato, G.M. Cragg, M.R. Boyd. *Natural Product Letters*, **14(5)**, 399(2000), <https://doi.org/10.1080/10575630008043774>
24. H.D.T. Mai, T.P. Toan, G.T. Huu, T.N. Le, V.T.K. Oanh, N.T.M. Hang, H.T. Thu, V.M. Chau, M. Litaudon, V.C. Pham, *Natural Product Research*, (2019), <https://doi.org/10.1080/14786419.2019.1587425>
25. N.S. Aminah, A.N. Kristanti, M. Tanjung, *Journal of Chemical and Pharmaceutical Research*, **6(6)**, 688(2014).

26. E.T. Arung, J.R. Sinamabela, E. Rosamah, I.W. Kusuma, H. Kuspradini, A.E. Alam, Y. Amen, H. Tanaka, D. Satria, K. Shimizu, H. Ishikawa, *Natural Product Communications*, **July 2019**, 1 (2019), <https://doi.org/10.1177/1934578X19867192>
27. A. Ogundajo, B. Okeleye, A.O. Ashafa, *Asian Pacific Journal of Tropical Biomedicine*, **7(7)**, 654 (2017), <https://doi.org/10.1016/j.apjtb.2017.06.014>
28. W. Aligita, S. Muhsinin, E. Susilawati, Dahlia, D.S. Pratiwi, D. Aprilliani, A. Artarini, I.K. Adnyana, *Rasayan Journal of Chemistry*, **12(1)**, 157(2019), <https://doi.org/10.31788/RJC.2019.1215059>
29. Minarti, A.H. Cahyana, A. Darmawan, Proceedings of the 5th International Symposium on Applied Chemistry 2019, AIP Conf. Proc. 2175, (2019), Jakarta, Indonesia, 020029-1-020029-9, <https://doi.org/10.1063/1.5134593>
30. R.T. Dewi, A. Darmawan, H. Mulyani., P.D.N. Lotulung, Minarti, Megawati, *Malaysia Journal of Science*, **37(1)**, 70 (2018), <https://doi.org/10.22452/mjs.vol37no1.5>
31. D. Sukandar, S. Nurbayti, T. Rudiana, T.W. Husna, *Indonesian Journal of Applied Chemistry*, **19(1)**, 11(2017).
32. M. Bule, A. Abdurahman, S. Nikfar, M. Abdollahi, M. Amini, *Food and Chemical Toxicology*, **125**, 496(2019), <https://doi.org/10.1016/j.fct.2019.01.037>

[RJC-6290/2020]