Biomarkers in Acute Myocardial Infarction

G. Anand Kumar MD DNB (Cardio)¹

¹Consultant Cardiologist, VIMS, Salem.

ABSTRACT

Acute myocardial infarction, the major cause of death in developed countries is becoming more prevalent in developing countries due to rapid urbanization and change to western life style. Myocardial infarction occurs due to occlusion of the coronary artery leading to myocardial necrosis, heart failure and arrhythmias, causing an increase in morbidity. Early recognition of acute myocardial infarction (AMI) leads to early treatment by means of thrombolysis, percutaneous intervention or coronary artery bypass surgery leading to decrease in the size of infarct and thereby reducing the complications of AMI. This article would focus on the currently used cardiac biomarkers for acute myocardial infarction and emerging role of high sensitive troponin in future and clinical usefulness of creatine kinase MB isoform.

Keywords : Cardiac biomarker, Troponin, High sensitive troponin, Creatine kinase MB (CK MB), Multi marker testing.

INTRODUCTION:

The Myocardial infarction which was a major cause for mortality and morbidity in western world has been catching up as an important cause of death in developing countries where communicable disease was the major cause of death. The reason for increase in coronary artery disease in the developing countries is due to westernization of the lifestyle. The traditional theory for causation of coronary artery disease centers on complex interplay between genetic and environmental, modifiable and non modifiable risk factors setting into motion an inflammatory cascade of monocyte migration, lipid oxidation and atheromatous plaque formation.¹

The cascade of thrombotic events following atherosclerotic plaque rupture causes occlusion of the coronary artery, interrupting blood supply and oxygen to myocardium resulting in infarction. Myocardial necrosis results in heart failure, myocardial rupture or arrhythmias. Early treatment

of acute myocardial infarction with treatment as fibrinolysis, percutaneous coronary intervention and coronary artery bypass grafting has improved outcome.²

For early diagnosis and initiation of treatment, cardiac biomarkers play a major role. This article would review the new biomarkers available and the advantage over the previously used biomarkers.

CURRENT CARDIAC BIOMARKERS:

Troponin:

The widely used biomarker for myocardial injury is cardiac troponin cTn. The troponin is made up of 3 subunits namely C, I and T which together control calcium mediated interaction of actin and myosin leading to contraction and relaxation of striated muscle. Troponin I and troponin T are expressed only in cardiac muscle; therefore these biomarkers have a high specificity for myocardial damage. CTn are detectable in peripheral circulation when damage to cardiac myocyte first leads to release of

Address for correspondence:

Dr.G. Anand Kumar MD DNB (Cardio), Consultant Cardiologist, VIMS, Salem.

Email id: ganandkumar@gmail.com Mobile no: 9843469245

DOI : <http://dx.doi.org/10.7324/NJBMS.2018.8303>

cytoplasmic cTn which accounts for 3% to 5% of cTnI and 7% cTn T.^{3,4} After infarction cTn is detectable in blood after a delay of 2 to 4 hrs and cTn remain detectable for 4 to 7 days for cTn I and 10 to 14 days for cTn T, cleared from circulation by reticuloendothelial system and fragmented into molecules that are cleared renally.

The sensitivity of cTn I and cTn T at initial presentation for acute myocardial infarction was 3.7% and 33.3% respectively which improved to 82% and 89% after 6 hrs and to 89% and 96% after 12 hrs,⁵ leading to development of new generation of high sensitive assays called hsc Tn ((high sensitive troponin) with a 10 to 100 fold lower limit of detection. The hscTn assays have allowed the diagnostic cut off to be lowered to the level of 99th percentile or lower.^{6,7}

Creatine Kinase (CK) and Creatine Kinase MB (CK MB)

Creatine kinase and its MB isoform CK-MB were the most commonly used serologic test for the diagnosis of myocardial infarction prior to widespread adoption of troponin. With the development of high sensitive troponin the use of creatine kinase and its isoform creatine kinase MB utility has markedly diminished over time. Increases in CK usually begin 4–6 h after the onset of MI, and peaks at around 18–24 h, while returning to baseline levels by 36–48 h.⁸ A total CK elevation of twofold above normal, with a simultaneously elevated CK-MB is required for the diagnosis of an MI. However, the important drawback of this is a false positive result, which can be due to skeletal muscle damage from trauma, cardiopulmonary resuscitation, defibrillation, cardiac and noncardiac surgical procedures, cocaine abuse, intramuscular injections, convulsions, alcohol intoxication, hypothyroidism, renal failure and pulmonary embolism.⁹ In the diagnosis of Acute coronary syndrome, CK is comparatively less

sensitive and specific than troponin. Several trials in the past have compared troponin with CK-MB. A review of around 30,000 patients in the multicenter, 'Can Rapid Risk Stratification of Unstable Angina Patients Suppress Adverse Outcomes with Early Implementation of the American College of Cardiology (ACC)/American Heart Association (AHA) (CRUSADE)' trial, showed that troponin was more sensitive and has an enhanced prognostic value than CK-MB.¹⁰ Clinical scenario where CK-MB measurement scores over troponin is in the setting of periprocedural PCI or CABG myocardial infarction. Also some clinicians prefer the use of CK-MB for the detection of early reinfarction, since CK-MB levels return to baseline in 36 to 48 hrs after infarction. Troponin does not normalize that rapidly and is elevated in plasma for 10 to 14 days.

Troponin VS CK-MB

A number of studies have shown that troponin have enhanced prognostic value compared with CK-MB measurements. The GEACE registry which studied more than 10,000 ACS patients, the in hospital mortality was highest when troponin was elevated¹¹ as shown in table 1.

Table 1. Troponin Vs CK-MB

Troponin	CK-MB	Mortality
Positive	Positive	7.7%
Positive	Negative	3.9%
Negative	Positive	2.3%
Negative	Negative	1.7%

Myoglobin

Myoglobin is a small cytoplasmic heme protein found in all muscles. Myoglobin increases within 1 to 3 hr in the setting of myocardial necrosis, peaks within 6 to 9 hr and become normal in 24 hr.¹² Myoglobin is the earliest marker to rise after acute myocardial infarction (2 hr from the onset of chest pain) because of its relatively small size and high cytoplasmic content. A single myoglobin measurement has been shown to have a sensitivity of 70 % and a negative predictive value of 97.4% for predicting acute myocardial infarction in patients suspected to have ACS. The disadvantage of myoglobin as novel biomarker is the rapid increase and normalization of myoglobin after AMI may lead to normal values for patient who presents 24 hrs after symptoms. Also myoglobin has limited specificity for myocardial necrosis in patients who have renal insufficiency and skeletal muscle trauma.¹³

Heart fatty acid binding protein (HFABP)

HFABP is a small low molecular weight (15kda), 132 amino acid, soluble protein with general characteristic resembling myoglobin, involved in fatty acid transport and metabolism. Although majority of HFABP is expressed in myocardium, a small quantity can be found in brain, kidney and skeletal muscle.¹⁴ HFABP has an early rise after an AMI, detected as soon as 30 minutes after onset of ischemia episode, peaks in blood after 6-8 hrs and returns to baseline value after nearly 24 to 30 hrs. Disadvantage of HFABP is that it is unsuitable as a diagnostic tool for patients presenting greater than 6 hrs from onset of symptoms due to rapid renal clearance. When HFABP was added to troponin for risk stratification, patient who has elevated troponin and HFABP has high mortality¹⁵, as shown in Table 2.

Table 2. HFABPs Vs Troponin

HFABP	Troponin	Mortality
>5.8 mg/l	Positive	7.93%
>5.8 mg/l	Negative	4.93%
<5.8 mg/l	Negative	0

HFABP is a promising biomarker to be used in combination with conventional troponin test readily available as a rapid bedside diagnostic tool.

Ischemia modified albumin(IMA)

IMA is a biomarker for acute ischemia that is approved by U.S. FDA. When exposed to ischemic condition, the N terminal of albumin is damaged, which makes it unable to bind metal and capable of being measured by albumin cobalt binding test.¹⁶ Its level in the blood increases within minutes of ischemia and return to normal within 6 to 12 hrs. IMA has been implicated in the detection of acute ischemia prior to necrosis.

Lactate dehydrogenase (LDH)

Lactate dehydrogenase was commonly used in past in combination with aspartate aminotransferase and creatine kinase in diagnosis of acute myocardial infarction. LDH activity rises to abnormal levels approximately 10 hrs after the onset of MI, peaks at 24 to 48 hrs and remain elevated for 6 to 8 days. Since troponin are more specific than LDH current recommendation no longer suggest LDH for the diagnosis of AMI.

MULTIMARKER TESTING:

The future of biomarker in the detection of myocardial injury call for a multimarker approach for diagnosis and prognosis. Current AHA guidelines for cTn measurement recommend testing on presentation and again at 8 to 12 hr post symptom onset. National Academy of clinical Biochemistry recommends an early marker at 0 to 6 hr and definitive marker 6 to 9 hr post presentation. Multiple markers analyzed on presentation and or repeated within a shorter time interval to comply with emergency department guideline may be sufficient for appropriate risk stratification. sensitivity for AMI 80 to 100% for CK MB /myoglobin up to 4 hr post presentation, 94 to 96.7% for cTn I/myoglobin upto 1.5 hr post presentation, 92.6% for cTnI/ CK MB upto 8 hr post presentation and 100% for cTn I, CK MB , myoglobin upto 2 hr post presentation compared with sensitivities for cTn alone of 86 to 92.3%.¹⁷

The emerging generation of high sensitive troponin assays obviate the need of other assays particularly myoglobin and CK MB as elevated high sensitive troponin may precede clear cut myocardial necrosis making it a single biomarker for acute myocardial infarction in the future.

REFERENCES:

1. Berenson GS, Srinivasan SR, Bao W, Newman WP 3rd, Tracy RE, et al. Association between multiple cardiovascular risk factors and atherosclerosis in children and young adults. The Bogalusa Heart Study. *N Engl J Med*.1998; 338: 1650-1656.
2. Tunstall-Pedoe H, Vanuzzo D, Hobbs M, Mahonen M, Cepaitis Z, Kuulasmaa K, et al: Estimation of contribution of changes in coronary care to improving survival, event rates, and coronary heart disease mortality across the WHO MONICA Project populations. *Lancet* 2000; 355: 688-700.
3. Higgins JP, Higgins JA. Elevation of cardiac troponin I indicates more than myocardial ischemia. *Clin Invest Med*. 2003; 26: 133-147.
4. Katus HA, Remppis A, Scheffold T, Diederich K W, Kuebler W. Intracellular compartmentation of cardiac troponin T and its release kinetics in patients with reperfused and nonreperfused myocardial infarction. *Am J Cardiol*.1991;67: 1360-1367.
5. Tucker JF, Collins RA, Anderson AJ, Hauser J, Kalas J, et al. Early diagnostic efficiency of cardiac troponin I and Troponin T for acute myocardial infarction. *Acad Emerg Med*.1997; 4: 13-21.
6. Jaffe AS, Lindahl B, Katus HA. Sensitive troponin I assay in patients with suspected acute coronary syndrome. *JAMA*. 2011; 306: 488-489
7. Thygesen K, Mair J, Katus H, Plebani M, Venge P, et al. Recommendations for the use of cardiac troponin measurement in acute cardiac care. *Eur Heart J*.2010; 31: 2197-2204.
8. Newby LK, Roe MT, Chen AY, et al. Frequency and clinical implications of discordant creatine kinase-MB and troponin measurements in acute coronary syndromes. *J. Am. Coll. Cardiol*. 2006; 47(2): 312–318.
9. Wolf PL. Abnormalities in serum enzymes in skeletal muscle diseases. *Am. J. Clin. Pathol*. 1991; 95(3): 293–296.
10. Patel MR, Chen AY, Peterson ED, et al. Prevalence, predictors, and outcomes of patients with non-ST-segment elevation myocardial infarction and insignificant

- coronary artery disease: results from the Can Rapid risk stratification of Unstable angina patients Suppress ADverse outcomes with Early implementation of the ACC/AHA Guidelines (CRUSADE) initiative. *Am. Heart J.* 2006;152(4): 641–647.
11. Santopinto JJ, Fox KA, Goldberg RJ, et al. Creatinine clearance and adverse hospital outcomes in patients with acute coronary syndromes: findings from the global registry of acute coronary events (GRACE) *Heart Br. Card. Soc.* 2003; 89(9): 1003–1008.
 12. Mair J, Artner-Dworzak E, Lechleitner P, Morass B, Smidt J, et al. Early diagnosis of acute myocardial infarction by a newly developed rapid immunoturbidimetric assay for myoglobin. *Br Heart J.* 1992; 68: 462-468.
 13. de Winter RJ, Koster RW, Sturk A, Sanders GT. Value of myoglobin, troponin T, and CK-MBmass in ruling out an acute myocardial infarction in the emergency room. *Circulation.* 1995; 92: 3401-3407.
 14. Chan D, Ng LL. Biomarkers in acute myocardial infarction. *BMC.* 2010; Med 8: 34.
 15. Kilcullen N, Viswanathan K, Das R, Morrell C, Farrin A, Barth JH, Hall AS: Heart-type fatty acid-binding protein predicts long-term mortality after acute coronary syndrome and identifies high-risk patients across the range of troponin values. *J Am Coll Cardiol* 2007; 50: 2061-2067.
 16. Sharma R, Gaze DC, Pellerin D, Mehta RL, Gregson H, et al. Evaluation of ischaemia-modified albumin as a marker of myocardial ischaemia in end-stage renal disease. *Clin Sci (Lond).* 2007; 113: 25-32.
 17. Macdonald SP, Nagree Y. Rapid risk stratification in suspected acute coronary syndrome using serial multiple cardiac biomarkers: a pilot study. *Emerg Med Australas.* 2008; 20: 403-409.

Received on 26/02/2018 Revised on 20/03/2018 Accepted on 02/04/2018