II. SPACE TECHNOLOGY AND ENVIRONMENTAL MONITORING

ASPECTS AND PERSPECTIVE OF INTERIM ECOLOGICAL MONITORING APPLICATION ON ECOSYSTEMS BY MEANS OF REMOTE SENSING

Kameliya Radeva

Abstract: Identifying and detailing the biophysical characteristics of species and habitats, habitat mapping, predicting the distribution of species and spatial variability in species richness and detecting the influence of natural and human-caused disasters on biodiversity urgently implies interim ecological monitoring implementation as part of regular ecological monitoring. The term ‘interim ecological monitoring’ implies the development of models for ecological monitoring, while integrating remotely sensed data. This paper explores how recent advances in spatial and spectral resolutions of satellite sensors can contribute to studying aspects of biodiversity, as well as discussing advantages and challenges of adopting such technologies for biodiversity conservation. The application of interim ecological monitoring methodology contributes to completion of the available data base for the actual condition of ecosystems considering different influence indicators. This paper demonstrates that interim remote sensing data can offer valuable information about diversity patterns and has the potential to become an effective tool in biodiversity evaluation and conservation. The review highlights the aspects and perspectives for interim ecological monitoring application in view of remote sensing potential for investigating the environmental and ecological processes, analyzing the condition of the landscape features and species diversity. This paper emphasizes on the possibilities of interim ecological monitoring to provide decision-makers with high quality and consistent data on ecosystems services potential.

Keywords: interim ecological monitoring, remote sensing, ecosystem services, biodiversity

INTRODUCTION

Monitoring is a special information system for repetitive observations and analyzes of one or more elements on the environment condition which is intended to register, assess and predict the influence of different factors on ecosystems. The system which monitors, analyzes and forecasts the biosphere changes resulting from different impacts is defined as ecological monitoring [1]. The purpose of the present report is to consider aspects and perspectives of interim ecological monitoring as part of ecological monitoring by means of aerospace data with different spectral and space resolution and in relevance of object type to be monitored. Current research illustrates how ecological problems ranging from biodiversity loss to land-use change have benefited greatly from advances in geospatial technologies such as GIS and remote sensing, both in the provision of data and access to spatial data analysis tools. The integration of GIS and remote sensing for ecological mapping and monitoring, while addressed in earlier investigations from Stoms and Estes, Franklin and Goodchild, has become more important as these data and technologies continue to evolve, and as ecological issues become more critical [2]. Reconciling differing stakeholder perspectives and needs will boost the timely provisioning of reliable information on the current and changing distribution of biodiversity to enable effective conservation management. Interim

ecological monitoring will enhance the process of developing tools and models for consistent multi-annual monitoring of protected areas and their surroundings through integrated use of remote-sensing and in-field data [3]. The application of interim ecological monitoring data and results for designing flexible measurements and undertaking adequate actions to maintain and improve ecosystems condition is just one aspect of issue to be discussed in this report.

The investigation made by Christine Schleupner in “Estimation of spatial wetland distribution potentials in Europe“ [4] refers to the role of remote sensing for monitoring application of wetlands. The author designed a model that provides an important prerequisite for the further development of a cost-effective wetland site-selection model of an EU-wide ecological network of wetland sites. The basis of this model is the optimal combination of existing spatial datasets to obtain the spatial distribution of wetlands by definition of flexible knowledge rules [5]. A promising approach is the one developed by Costanza et al. [6] to model the effects of wetlands for hurricane protection in the US. In principle, this approach could be used to model several ecosystem types simultaneously (e.g., mangroves, corals and seagrasses) and their interactions. Costanza et al. considered extending their approach to the global scale, but were prevented from doing so by the lack of good wetland maps [6, 7]. Even CORINE

doi.org/10.32006/eeep.2018.2.2634

biotopes data [8] that are based on reported NATURA2000 sites do not represent the existing wetlands completely. Most studies concentrate on nonforested peatland only [9]. The most detailed information about wetland habitats in Europe offers the EUNIS (European Nature Information System). However, interim ecological monitoring can generate data for wetlands condition and helps to update the existing or develop new and more precise maps of wetlands.

Accurate ecological monitoring has been implemented in Kruger National Park (KNP), Southern Africa to inform management practices, thereby maintaining biodiversity. The park operates as a preserve for endemic flora and fauna, and is a major visitor attraction. It is believed that management practices throughout the twentieth century have fundamentally altered vegetation distribution throughout the park. In particular, numerous artificial water resources were created to attract wildlife to specific locations for viewing by tourists. The increased water resources may have led to an overall increase in vegetation abundance, and may be contributing to structural homogenisation of the park's vegetation. Related processes include infestation of alien plant species and dramatic growth in certain animal populations, notably elephant and rhinoceros. A rich remote sensing (and associated geographical information system) data resource is available for the park, including images from Landsat, SPOT, IKONOS and QuickBird instruments, and numerous airborne images and photographs. Given the difficulties of allocating pixels to specific land cover classes in mixed environments like KNP, it can be useful to use continuous rather than thematic representations. Research remains ongoing to exploit fine spatial resolution satellite sensor imagery, in combination with coarser spatial resolution imagery, for ecological investigation in KNP. In particular, vegetation information derived from remote sensing is integrated with animal population data to further our understanding of the dynamic relationship between flora and fauna [10, 11, 12].

METHODS AND MATERIALS

A key element in the environmental monitoring process is to achieve reliable, timely and effective environmental monitoring of the state and trends in biodiversity that will enable the establishment and implementation of conservation strategies. Interim ecological monitoring gives such an opportunity as results obtained by its application will contribute to

data availability for ecosystem status assessment upon the time of its implementation and ability to convey information to the policy making and implementation processes of indicators for integrated ecosystems and their services assessment. Remote-sensing technologies can deliver data on habitat quantity (amount, configuration) and quality (e.g., structure, distribution of individual plant species, habitat types and/or communities, and persistence) across a range of spatial resolutions and temporal frequencies [13, 14].

In the present report two objectives on which the interim ecological monitoring has been applied are analyzed: Wetland in the area of quarry lakes, Negovan village, Sofia city and Nature Reserve SINI BRYAG near Bozhevtzi village, Sliven region, that is a century-old virgin beech forest. For the implementation of interim ecological monitoring on described sites remote sensing images with high resolutions have been used. This type of images serve as measure biophysical properties of surface environments (e.g., vegetation, water and soil) of relevance to conservation, changes in the state of land cover or habitat quantity/quality and generate categorical products (2D/3D thematic maps and spatial distributions) [15]. Images with high resolution have been applied for monitoring of small-sized habitats and/or areas with difficult access. Interim ecological monitoring on both objects has been applied with aim to establish and track the ecosystem condition prior to regular monitoring.

The structure of interim ecological monitoring application is designed based on three main criteria [16]. The first criteria can be determined when the temporal interval of the natural hazard occurrence falls within the temporal interval accepted in the methodologies for regular monitoring implementation:

$$\Delta t_{RM}^i = t_{LRM}^i - t_{PRM}^i \quad (1),$$

where Δt_{RM}^i – temporal interval of regular monitoring for a certain ecosystem type (i),

t_{LRM}^i – last temporal point of regular monitoring for a certain ecosystem type (i),

t_{PRM}^i – previous temporal point of regular monitoring for a certain ecosystem type (i).

The second criterion refers to establishing the necessity of interim ecological monitoring implementation. We can define two cases. In the first one a check whether t_{EPHENj}^i belongs or not to Δt_{RM}^i , where t_{EPHENj}^i is temporal interval of the occurrence of natural hazard or manmade influence for a certain ecosystem type (i) and Δt_{RM}^i is

temporal interval of regular monitoring for a certain ecosystem type (i).

The third criteria demonstrates the dependence between temporal interval of regular monitoring Δt_{RM}^i for a certain ecosystem type (i) and temporal

interval for interim ecological methodology Δt_M^i , i.e.

$$\Delta t_{RM}^i \in \Delta t_M^i \quad (2)$$

The structure of interim ecological monitoring methodology has been shown on Fig. 1 below.

Fig.1. Structure on Interim Ecological Monitoring Methodology

During interim ecological monitoring complicated processes can be analyzed that take place in ecosystem and thus implies biotic and abiotic factors to be included in multicomponent analysis [17, 18].

The essential moment in remote sensing application is the choice of satellite data consistent with temporal criteria of interim ecological monitoring. On the other hand the data choice should be consistent with the ecosystems type and their functional and geometrical characteristics. Interim monitoring on environment survey methodology includes the application of Vegetation Indices (VI), ranging from evapotranspiration estimates to forest structure. The Normalized Difference Vegetation Index (NDVI) and Normalized Difference Greenness Index (NDGI) have become the most extensively used VI in ecological remote sensing.

Most wetland studies deal with water quality and quantity assessments at local, regional or watershed scale, others concentrate on wetland functions and values or on vegetation and biodiversity patterns. [19–31]. Interim ecological monitoring data can help to distinguish between existing wetland habitats and potential convertible sites for restoration of wetland biotopes by considering actual land use options. The wetland in the area of quarry lakes in Negovan village that has been analyzed in the present article is such an example.

RESULTS AND DISCUSSION

The results in the present paper have been obtained by interim ecological monitoring implementation on two different ecosystems: wetland in the area of quarry lakes in Negovan village and protected area (Nature Reserve SINI BRYAG). The research (interim ecological monitoring) has been fulfilled for two different cases considering the lack of natural hazards for the Nature reserve and abiotic factor influence for the wetland.

For the wetland high resolution data and multispectral satellite data from sensor Sentinel 2 before and after the application of restoration measures and for the nature reserve multispectral satellite data from sensor Sentinel 2 for two dates have been used. The temporal intervals are selected in a way that they coincide with the temporal interval of regular monitoring. Terrestrial data from precedent temporal intervals have been used for the verification of the obtained results from interim ecological monitoring application.

For the observed ecosystems methods of satellite data processing that include orthogonalization of satellite images, NDVI, NDGI, correlation analysis and multicomponent analysis have been used. [32, 33].

A common means of characterising the presence of vegetation, for instance, is through NDVI analysis

(Fig. 3, 4). This has the benefit that pixels are not forced into (sometimes inappropriate) classes, but instead provide a proportional measure of vegetation. Further, analyses such as NDVI can contribute to the derivation of biophysical variables such as LAI and NPP. Habitat interim monitoring can be performed by comparing analyses such as land cover classification and NDVI over time. Research remains ongoing to exploit fine spatial resolution satellite sensor imagery, in combination

with coarser spatial resolution imagery for ecological investigation in different ecosystems types while interim ecological monitoring has been implemented.

In Fig. 2 images from Google Earth with different dates and for a four-year period, as well as composite satellite images for 06/04/2016 (Ch 432,482,1384) from Multispectral Instrument (MSI) from Sentinel 2 for the Wetland in the area of quarry lakes in Negovan village have been presented.

Fig. 2. Wetland Negovan (Sofia region, Bulgaria) from Google Earth for different temporal dates and Sentinel 2 (g, h, i)

Having in regard the information submitted in Fig. 3 a) 3D-distribution of W/G 06/04/2016, b) 3D-distribution of Wetness for 09/04/2017 and 23/07/2017, c) 3D-distribution of W/G 09/04/2017, e) 3D-distribution of W/G 23/07/2017, it can be concluded that the water level has started to decrease and vegetation cover to increase after 06/04/2016 till the last image for 11/08/2017. Analysis shows that the decrease of water surface is

due to low rain and demonstrates there is a process of fouling in which case adequate measures for area restoration of water surface for the wetland should be completed. In order to test the hypotheses relating to biodiversity change (e.g., on species-habitat relationships) it is necessary that upon interim ecological monitoring application data with high resolution can be used in combination with discretely collected field data [34].

In Fig.4 composite satellite images (Ch 432) from MSI from Sentinel 2 for the protected area Nature Reserve SINI BRYAG for three different years 2010, 2015 and 2017 and similar dates have been presented. Results for the value of Greenness

and spectral reflectance characteristic diagram for the period 19-24/08/2017 have been obtained based on Tasseled Cup Transformation (TCT). The results show that no change in the vegetation dynamics has been detected.

Fig. 3. Wetland Negovan (Sofia region, Bulgaria): a) 3D-distribution of W/G 06/04/2016, b) 3D-distribution of Wetness for 09/04/2017 and 23/07/2017, c) 3D- distribution of G/W 09/04/2017, e) 3D- distribution of G/W 23/07/2017

Referring to the obtained results from the interim ecological monitoring on two different ecosystem types (wetland in the area of quarry and protected area) the following conclusions can be generated:

- The satellite data from remote sensing are applicable and present objective results in interim ecological monitoring implementation;

- The results from the wetland research demonstrate that interim monitoring should be implemented since even only by abiotic factors monitoring in case of lack of natural hazard occurrence, substantial changes that lead to misbalance in the sustainable development of the ecosystem have been observed;

- In the case of the Nature reserve SINI BRYAG substantial modifications during interim monitoring are not identified. This fact confirms the sustainable development of the ecosystem during the period considered;

- Interim ecological monitoring methodology includes estimation of vegetation structural and biophysical parameters from remotely-sensed data. Therefore interim ecological monitoring uses empirical methods to relate spectral data and various image derivatives to vegetation characteristics. If these parameters are strongly correlated with remotely sensed data, they can be used to predict those biophysical attributes over large extents;

- Data derived from interim ecological monitoring can be used to improve the Species Distribution Models by its updating and high quality of input data;

- Interim ecological monitoring data can be valuable for conducting specific actions for wetland/protected area restoration since river regulations, peat extraction or urbanization on former wetland areas often lead to changed potentials in wetland/protected area restoration.

Fig. 4. Nature Reserve SINI BRYAG: a) and b) – Airphoto images; c) and d) – Composite Satellite images (Ch 432)

Fig. 5. Nature Reserve SINI BRYAG: a) 3D- distribution of NDVI / NDGI/ dNDVI for 19-24/08/2017. b) Spectral reflectance characteristic for 19-24/08/2017

ASPECT AND PERSPECTIVES

Maintaining biodiversity in the face of increasing anthropogenic pressures has become a major concern for governmental agencies and institutions that manage protected areas, and is one of the major focal areas of the European Union's Biodiversity strategy to 2020 [35]. Many researches have been made about anthropogenic impacts on land-cover change and the extent of remaining habitats [36] but the magnitude and rate of change in biodiversity within habitats are typically poorly quantified [37]. Therefore, interim ecological monitoring can be accepted as a way for providing data necessary for establishing magnitude and rate of change in biodiversity within habitats models.

Improving a methodology for interim environmental monitoring that serves as a tool to complement the available ecosystem current database is a perspective for ecosystem assessment and analysis using remote sensing data. Analyzing and discussing issues on data reliability and techniques for actual data for the condition of different environmental factors turned out to be a challenge for the application of interim ecological monitoring methodology. Such issues can be referred to image corrections and registration of image data (for biophysical retrieval and change detection), along with proper registration of multi-date, multi-scale and multi-source datasets; adopting new advances in image interpretation, as currently

exploited in the remote-sensing area, both integrating object oriented image analysis and knowledge-driven information mining; defining habitat type, extent and quality; assessing habitat quality and biodiversity from remote-sensing data; incorporating uncertainty into mapping habitats using remote sensing data; integrating expert knowledge in ecology and remote-sensing and including appropriate ground-truthing; selecting appropriate space-time-related scales of application; analysing the interrelationships between habitat quality, distribution and structure.

Using continuous or gradual properties of spectral data in Species Distribution Models is a key challenge in near future. More attention should be placed on the ability to consistently characterize biological diversity at multiple scales, using integrated remotely sensed and GIS data. In addition to using the reflectance values, the unique synoptic perspective of remote sensing for quantifying landscape characteristics should also be more fully studied. Change detection models should explore more flexible statistical methods and include ancillary GIS data. In particular, vegetation information derived from remote sensing can be integrated with animal population data to further understanding of the dynamic relationship between flora and fauna.

The information used for scientifically valid ecological mapping and monitoring needs to be frequently updated, sufficiently detailed and spatially continuous. Although the traditional inconsistency in spatial scale between remotely sensed data (indirect) and ecological field observations (direct) has been a major obstacle to more extensive integration of remote sensing in ecological research, access to increasingly fine spatial resolution data has resulted in great progress in this area [38, 39]. Further, advances in remote sensing theory, data and technology over the past 35 years have led to general and robust methods of large area data collection that, for many ecological attributes, can provide more reliable estimates than field methods. Remote sensing facilitates data collection in difficult- or impossible-to-reach areas and provides an important synoptic and multitemporal perspective. There is also a growing demand of policy makers and researchers for high-accuracy landscape information at the European level [40, 41].

CONCLUSIONS

Interim ecological monitoring based on remote sensing helps identification of biodiversity elements that are underrepresented in conservation areas and thus it is useful for modelling disturbed landscapes

However, aggregating statistical and spatial data from many sources into one database, having in regard that sources differ in spatial accuracy, reliability, acquisition data and class definition yet remains a challenge for future common cooperation among environmentalists and remote sense experts.

ACKNOWLEDGEMENT: *This study was carried out with the support of Project BG161PO003-1.2.04-0053-C0001 from Operational Programme "Development of the Competitiveness of the Bulgarian Economy 2007–2013" and Programme "COPERNICUS" of European Union (European Space Agency).*

REFERENCES

1. Radeva, K., Environmental monitoring on wetlands restoration in the area of quarry lakes, *Ecological Engineering and Environment Protection*, 4, 2015, 15-22, ISSN:1311-8668.
2. Miller J. , Rogan J., Using GIS and remote sensing for ecological mapping and monitoring, In: *Integration of GIS and Remote Sensing*, John Wiley & Sons Ltd., The Atrium, Southern Gate, Chichester, West Sussex , PO19 8SQ, England
3. Lucas, R.M., Blonda, P., Bunting, P., Jones, G., Inglada, J., Arias, M., Kosmidou, V., Manakos, I., Adamo, M., Charmock, R., Tarantino, C., Mùcher, C.A., Jongman, R.H.G., Kramer, H., Arvor, D., Honrado, J.P., Mairota, P., The Earth Observation Data for Habitat Monitoring (EODHaM) system. *Int. J. Appl. Earth Obs. Geoinf.* 37, 2015, 1–6.
4. Schleupner, Chr., Estimation of spatial wetland distribution potentials in Europe, http://epub.sub.uni-hamburg.de/epub/volltexte/2012/16805/pdf/Wetland_I_schleupner135.pdf (Retrieved on April 25, 2018)
5. Mùcher C.A., Hennekens S.M., Bunce R.G.H. & Schaminée J.H.J., Spatial identification of European Habitats to support the design and implementation of a pan-European ecological network. In: D McCollin & JI Jackson (eds.) *Planning, people and practice; the landscape ecology of sustainable landscapes*. IALE, UK, 2004, pp. 217–225.
6. Costanza, R., Pérez-Maqueo, O., Martinez, M. L., Sutton, P., Anderson, S. J., & Mulder, K., The value of coastal wetlands for hurricane protection. *Ambio* 37, 2008, 241–248.
7. Balmford A., Rodrigues Ana S.L., *The Economics of Ecosystems and Biodiversity: Scoping the Science* Env/070307/2007/486089/Etu/B2 Final Report, University of Cambridge
8. European Environment Agency, CORINE biotopes data, <https://www.eea.europa.eu/data-and-maps/data/corine-biotopes> (Retrieved on April 25, 2018).

9. Montanarella L., Jones R.J.A. & Hiederer R., The distribution of peatland in Europe, In: International Mire Conservation Group and International Peat Society (eds.) *Mires and Peat*, Volume 1 (Article 1), 2006, Available on http://www.mires_and_peat.net.
10. Van Wilgen, B.W., Biggs, H.C. and Potgieter, A.L.F., Fire management and research in the Kruger National Park, with suggestions on the detection of thresholds of potential concern, *Koedoe*, 41, 1998, 69-87.
11. Eckhardt, H., Van Wilgen, B.W. and Biggs, H.C., Trends in woody vegetation cover in relation to fire in the Kruger National park, South Africa, between 1944 and 1993, *African Journal of Ecology*, 38, 2000, 108-115.
12. Alpin, P., Remote sensing as a means of ecological investigation, January 2004, www.researchgate.net/publication/228599659_Remote_sensing_as_a_means_of_ecological_investigation
13. He, K.S., Rocchini, D., Neteler, M., Nagendra, H., Benefits of hyperspectral remote sensing for tracking plant invasions. *Divers. Distrib.* 17, 2011, 381-392.
14. Wulder, M.A., Hall, R.J., Coops, N.C., Franklin, S.E., High spatial resolution remotely sensed data for ecosystem characterization. *Bioscience* 54, 2004, 511-521.
15. Nagendra, H., Lucas, R.M., Honrado, J.P., Jongman, R.H.G., Tarantino, C., Adamo, M., Mairota, P., Remote-sensing for conservation monitoring: assessing protected areas, habitat extent, habitat condition, species diversity and threats. *Ecol. Indic.* 33, 2013, 45-59
16. Radeva K., Ivanova I., Borisova D. Application of remote sensing for ecosystems monitoring and risk assessment, Sixth international Conference on Remote sensing and GeoInformation of the Environment Proc SPIE 26-29 March, 2018, Paphos, Cyprus.
17. Nedkov R., Velizarova E., Molla I., Zaharinova M., NDVI changes and their relationships with climatic parameters for the South-eastern part of BULGARIA, Third European SCGIS Conference of Geoinformation technologies for natural and cultural heritage conservation, 2016, 32-41.
18. Aapo Hyvarinen, Juha Karhunen, Erkki Oja, Independent Component Analysis, A Wiley Interscience Publication John Wiley & Sons INC., 2001, 141-143.
19. Brinson M.M. & Rheinhardt R., The role of reference wetlands in functional assessment and mitigation, *Ecological Applications* 6 (1), 1996, 69-76.
20. Russel G.D., Hawkins C.P. & O'Neill M.P., The role of GIS in selecting sites for riparian restoration based on hydrology and land use, *Restoration Ecology* 5 (4S), 1997, 56-68.
21. Bhaduri B. Harbor J., Engel B. & Grove M., Assessing watershed-scale, long-term hydrologic impacts of land-use change using a GIS-NPS model. *Environmental Management* 26 (6), 2000, 643-658.
22. Richardson, C.J. & Nunnery, K., Ecological Functional Assessment (EFA): A new approach to determining wetland health. *Transformations of Nutrients in Natural and Constructed Wetlands*, 2001, 95-111.
23. Winter T.C. & Labaugh J.W., Hydrologic considerations in defining isolated wetlands. *Wetlands* 23 (3), 2003, 531-540.
24. Vepraskas M.J., White J.G., Huffman, R.L., Fernandez G.P., Skaggs R.W., Lees B., Zanner C.W., Stucky J.M., Gregory J.D., Broome W.S., Ewing J.M., Szuch R.P., Luginbuhl S. & Kreiser G.S., Methodology to assess soil, hydrologic, and site parameters that affect wetland restoration. The Center for Transportation and the Environment, N.C.DOT, Report No. FHWA/NC/2003-06, 2005.
25. Brinson M.M., Hydrogeomorphic Classification for Wetlands. US Army Corps of Engineers, Wetland Research Program. TR: WRP-DE-4, Washington DC, 1993.
26. Bayless E.R., A study of natural and restored wetland hydrology. U.S. Geological Survey Fact Sheet FS-104-99. U.S. Geological Survey, Department of the Interior, 1999.
27. Leibowitz S.G. Isolated wetlands and their functions: An ecological perspective. *Wetlands*, 23 (3), 2003, 517-531.
28. Mitsch W.J. & Gosselink J.G. The value of wetlands: The importance of landscape setting and scale. *Ecological Economics* 35, 2000, 25-33.
29. Miller R.C. & Zedler J.B. Responses of native and invasive wetland plants to hydroperiod and water depth. *Plant ecology* 167, 2003, 57-69.
30. Kiviat E. & MacDonald K., Biodiversity Patterns and conservation in the Hackensack Meadowlands, New Jersey. *Urban Habitats* 2 (1), 2004, 28-61.
31. Mahaney W.M, Wardrop D.H. & Brooks R.P., Impacts of sedimentation and nutrient enrichment on wetland plant community development. *Plant Ecology* 175, 2004, 227-243.
32. Nedkov, R., Orthogonal transformation of segmented images from the satellite SENTINEL-2, *Comptes rendus de l'Académie bulgare des Sciences*, 70, 5, Prof. Marin Drinov Publishing House of Bulgarian Academy of Sciences, 2017, 687-692.

33. Nedkov, R., Normalized Differential Greenness Index for Vegetation Dynamics Assessment, Comptes rendus de l'Académie bulgare des Sciences, 70, 8, Prof. Marin Drinov Publishing House of Bulgarian Academy of Sciences, 2017, 1143-1146, ISSN:1310-1331.
34. Newton, A.C., Hill, R.A., Echeverría, C., Golicher, D., Benayas, J.M.R., Cayuela, L., Hinsley, S.A., Remote-sensing and the future of landscape ecology. Prog. Phys. Geogr. 33, 2009, 528-546.
35. MAES Mapping and Assessment of Ecosystems and their Services. An analytical framework for ecosystem assessments under action 5 of the EU biodiversity strategy to 2020. Technical Report 2014 – 080, European Commission, 2014.
36. Blaschke, T., Towards a framework for change detection based on image objects. Göttinger Geographische Abh. 113, 2005, 1-9.
37. Lengyel, S., Dri, E., Varga, Z., Horvat, R., Tothmeresz, B., Henry, P., Kobler, A., Kutnar, L., Babij, V., Seliskar, A., Christia, C., Papastergiadou, E., Gruber, B., Henle, K., Habitat monitoring in Europe: a description of current practices. Biodivers. Conserv. 17, 2008, 3327-3339.
38. Turner, W., Spector, S., Gardiner, N., Fladeland, M., Sterling, E. and Steininger, M., Remote sensing for biodiversity science and conservation. Trends in Ecology and Evolution, 18, 2003, 306-314.
39. Kerr, J.T. and Ostrovsky, M., From space to species: ecological applications of remote sensing. Trends in Ecology and Evolution, 18, 2003, 299-305.
40. Wascher D.M., Agri-environmental indicators for sustainable agriculture in Europe. European Centre for Nature Conservation, Tilburg. ECNC Technical Report Series, Wageningen, 2000.
41. Klijn J.A., Driving forces behind landscape transformation in Europe, from a conceptual approach to policy options. In: RHG Jongmann (ed.) The new dimensions of the European landscape. Wageningen UR Frontis Series, Springer, Berlin, 2002, 201-218.

АСПЕКТИ И ПЕРСПЕКТИВИ НА ПРИЛАГАНЕТО НА МЕЖДИНЕН ЕКОЛОГИЧЕН МОНИТОРИНГ ЧРЕЗ ИЗПОЛЗВАНЕ НА ДИСТАНЦИОННИ ИЗСЛЕДВАНИЯ

Камелия Радева

Резюме: Идентифицирането и детайлизирането на характеристиките на видовете и местообитанията, картографирането на местообитанията, прогнозирането на разпространението на видовете и пространствената променливост на видовото богатство, установяването на влиянието на природни и причинени от човека бедствия върху биоразнообразието налагат извършването на междинен екологичен мониторинг като част от регулярния екологичен мониторинг. Терминът "междинен екологичен мониторинг" предполага разработването на модели за екологичен мониторинг, чрез използване на данни от дистанционни изследвания. Тази статия изследва как последните постижения на спътникови сензори с висока пространствена и спектрална разделителна способност могат да допринесат за изследване на аспектите на биоразнообразието, както и за дебатирание на предимства и предизвикателства, свързани с използване на такива технологии за опазване на биологичното разнообразие. Прилагането на методика за междинен екологичен мониторинг допринася за допълване на наличната база данни за актуалното състояние на екосистемите, като се имат предвид различни показатели на влияние. Статията показва, че данните от междинния екологичен мониторинг на базата на дистанционни изследвания могат да предоставят ценна информация за модели на екологичното разнообразие и имат потенциал да се превърнат в ефективен инструмент за оценка и опазване на биологичното разнообразие. Статията акцентира върху аспектите и перспективите от прилагането на междинен екологичен мониторинг от гледна точка на възможностите на дистанционните изследвания за проучване на екологичните процеси и анализиране на състоянието на характеристиките на ландшафта и видовото разнообразие. Тази статия подчертава възможностите на междинния екологичен мониторинг за осигуряване на отговорните институции с висококачествени данни относно потенциала на екосистемните услуги.

Ключови думи: междинен екологичен мониторинг, дистанционни изследвания, екосистемни услуги, биоразнообразие

Kameliya Radeva

Space Research and Technology Institute
at the Bulgarian Academy of Science
Acad. Georgi Bonchev Str., bl. 1
1113 Sofia, Bulgaria
Tel.: +359 879 379 023
E-mail: kameliya.radeva@space.bas.bg

Камелия Радева

Институт за космически изследвания и технологии
– БАН
ул. „Акад. Георги Бончев“, бл.1
София 1113
Tel.: +359 879 379 023
E-mail: kameliya.radeva@space.bas.bg