ISSN 2518-1726 (Online),
ISSN 1991-346X (Print)

ҚАЗАҚСТАН РЕСПУБЛИКАСЫ
ҰЛТТЫҚ ҒЫЛЫМ АКАДЕМИЯСЫНЫҢ
әл-Фараби атындағы Қазақ ұлттық университетінің

Х А Б А Р Л А Р Ы

ИЗВЕСТИЯ

НАЦИОНАЛЬНОЙ АКАДЕМИИ НАУК
РЕСПУБЛИКИ КАЗАХСТАН
Қазақстан Ұлттық академиясының
им. әл-Фараби

N E W S

OF THE NATIONAL ACADEMY OF SCIENCES
OF THE REPUBLIC OF KAZAKHSTAN
Al-Farabi
Kazakh National University

SERIES PHYSICO-MATHEMATICAL

1 (335)

JANUARY – FEBRUARY 2021

PUBLISHED SINCE JANUARY 1963

PUBLISHED 6 TIMES A YEAR

ALMATY, NAS RK

NAS RK is pleased to announce that News of NAS RK. Series physico-mathematical journal has been accepted for indexing in the Emerging Sources Citation Index, a new edition of Web of Science. Content in this index is under consideration by Clarivate Analytics to be accepted in the Science Citation Index Expanded, the Social Sciences Citation Index, and the Arts & Humanities Citation Index. The quality and depth of content Web of Science offers to researchers, authors, publishers, and institutions sets it apart from other research databases. The inclusion of News of NAS RK. Series of chemistry and technologies in the Emerging Sources Citation Index demonstrates our dedication to providing the most relevant and influential content of chemical sciences to our community.

Қазақстан Республикасы Ұлттық ғылым академиясы "ҚР ҰҒА Хабарлары. Физикалық-математикалық сериясы" ғылыми журналының Web of Science-тің жаңаланған нұсқасы Emerging Sources Citation Index-те индекстелуге қабылданғанын хабарлайды. Бұл индекстелу барысында Clarivate Analytics компаниясы журналды одан әрі the Science Citation Index Expanded, the Social Sciences Citation Index және the Arts & Humanities Citation Index-ке қабылдау мәселесін қарастыруда. Web of Science зерттеушілер, авторлар, баспашылар мен мекемелерге контент тереңдігі мен сапасын ұсынады. ҚР ҰҒА Хабарлары. Химия және технология сериясы Emerging Sources Citation Index-ке енуі біздің қоғамдастық үшін ең өзекті және беделді химиялық ғылымдар бойынша контентке адалдығымызды білдіреді.

НАН РК сообщает, что научный журнал «Известия НАН РК. Серия физико-математическая» был принят для индексирования в Emerging Sources Citation Index, обновленной версии Web of Science. Содержание в этом индексировании находится в стадии рассмотрения компанией Clarivate Analytics для дальнейшего принятия журнала в the Science Citation Index Expanded, the Social Sciences Citation Index и the Arts & Humanities Citation Index. Web of Science предлагает качество и глубину контента для исследователей, авторов, издателей и учреждений. Включение Известия НАН РК в Emerging Sources Citation Index демонстрирует нашу приверженность к наиболее актуальному и влиятельному контенту по химическим наукам для нашего сообщества.

Б а с р е д а к т о р ы
ф.-м.ғ.д., проф., ҚР ҰҒА академигі
Ғ.М. Мұтанов

Р е д а к ц и я а л қ а с ы:

Асанова А.Т. проф. (Қазақстан)
Бошкаев К.А. PhD докторы (Қазақстан)
Байгунчечков Ж.Ж. проф., академик (Қазақстан)
Quevedo Hernando проф. (Мексика)
Жүсіпов М.А. проф. (Қазақстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Қазақстан)
Михалевич А.А. проф., академик (Белорусь)
Мырзакулов Р. проф., академик (Қазақстан)
Рамазанов Т.С. проф., академик (Қазақстан)
Такибаев Н.Ж. проф., академик (Қазақстан), бас ред. орынбасары
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Қазақстан)
Харин С.Н. проф., академик (Қазақстан)

«ҚР ҰҒА Хабарлары. Физика-математикалық сериясы».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Меншіктенуші: «Қазақстан Республикасының Ұлттық ғылым академиясы» РҚБ (Алматы қ.).

Қазақстан Республикасының Ақпарат және коммуникациялар министрлігінің Ақпарат комитетінде
14.02.2018 ж. берілген № **16906-Ж** мерзімдік басылым тіркеуіне қойылу туралы куәлік.

Тақырыптық бағыты: *физика-математика ғылымдары және ақпараттық
технологиялар саласындағы басым ғылыми зерттеулерді
жариялау.*

Мерзімділігі: жылына 6 рет.

Тиражы: 300 дана.

Редакцияның мекенжайы: 050010, Алматы қ., Шевченко көш., 28; 219, 220 бөл.;

тел.: 272-13-19; 272-13-18,

<http://physics-mathematics.kz/index.php/en/archive>

© Қазақстан Республикасының Ұлттық ғылым академиясы, 2021

Типографияның мекенжайы: «NurNaz GRACE», Алматы қ., Рысқұлов көш., 103.

Главный редактор
д.ф.-м.н., проф. академик НАН РК
Г.М. Мутанов

Редакционная коллегия:

Асанова А.Т. проф. (Казахстан)
Бошкаев К.А. доктор PhD (Казахстан)
Байгунчеков Ж.Ж. проф., академик (Казахстан)
Quevedo Hernando проф. (Мексика)
Жусупов М.А. проф. (Казахстан)
Ковалев А.М. проф., академик (Украина)
Калимолдаев М.Н. проф., академик (Казахстан)
Михалевич А.А. проф., академик (Беларусь)
Мырзакулов Р. проф., академик (Казахстан)
Рамазанов Т.С. проф., академик (Казахстан)
Такибаев Н.Ж. проф., академик (Казахстан), зам. гл. ред.
Тигиняну И. проф., академик (Молдова)
Уалиев З.Г. проф., чл.-корр. (Казахстан)
Харин С.Н. проф., академик (Казахстан)

«Известия НАН РК. Серия физика-математическая».

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Собственник: РОО «Национальная академия наук Республики Казахстан» (г. Алматы).

Свидетельство о постановке на учет периодического печатного издания в Комитете информации Министерства информации и коммуникаций Республики Казахстан № 16906-Ж, выданное 14.02.2018 г.

Тематическая направленность: *публикация приоритетных научных исследований в области физико-математических наук и информационных технологий.*

Периодичность: 6 раз в год.

Тираж: 300 экземпляров.

Адрес редакции: 050010, г. Алматы, ул. Шевченко, 28; ком. 219, 220; тел.: 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© Национальная академия наук Республики Казахстан, 2021

Адрес типографии: «NurNaz GRACE», г. Алматы, ул. Рыскулова, 103.

Editor in chief
doctor of physics and mathematics, professor, academician of NAS RK
G.M. Mutanov

Editorial board:

Asanova A.T. prof. (Kazakhstan)
Boshkayev K.A. PhD (Kazakhstan)
Baigunchekov Zh.Zh. prof., akademik (Kazakhstan)
Quevedo Hemando prof. (Mexico)
Zhusupov M.A. prof. (Kazakhstan)
Kovalev A.M. prof., academician (Ukraine)
Kalimoldaev M.N. prof., akademik (Kazakhstan)
Mikhalevich A.A. prof., academician (Belarus)
Myrzakulov R. prof., akademik (Kazakhstan)
Ramazanov T.S. prof., akademik (Kazakhstan)
Takibayev N.Zh. prof., academician (Kazakhstan), deputy editor in chief.
Tiginyanu I. prof., academician (Moldova)
Ualiev Z.G. prof., chl.-korr. (Kazakhstan)
Kharin S.N. prof., academician (Kazakhstan)

News of the National Academy of Sciences of the Republic of Kazakhstan. Physical-mathematical series.

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Owner: RPA "National Academy of Sciences of the Republic of Kazakhstan" (Almaty).

The certificate of registration of a periodical printed publication in the Committee of information of the Ministry of Information and Communications of the Republic of Kazakhstan **No. 16906-Ж**, issued on 14.02.2018.

Thematic scope: *publication of priority research in the field of physical and mathematical sciences and information technology.*

Periodicity: 6 times a year.

Circulation: 300 copies.

Editorial address: 28, Shevchenko str., of. 219, 220, Almaty, 050010, tel. 272-13-19; 272-13-18,
<http://physics-mathematics.kz/index.php/en/archive>

© National Academy of Sciences of the Republic of Kazakhstan, 2021

Address of printing house: «NurNaz GRACE», 103, Ryskulov str, Almaty.

NEWS

OF THE NATIONAL ACADEMY OF SCIENCES OF THE REPUBLIC OF KAZAKHSTAN
PHYSICO-MATHEMATICAL SERIES

ISSN 1991-346X

Volume 1, Number 335 (2021), 6 – 13

<https://doi.org/10.32014/2021.2518-1726.1>

UDC 531.31; 519.21

MRNTI 27.29.17, 27.35.30

M.I. Tleubergenov^{1,2}, G.K. Vassilina^{1,3}, A.T. Sarypbek^{1,2}

¹Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan;

²Al-Farabi Kazakh National University, Almaty, Kazakhstan;

³Almaty University of Energy and Communications, Almaty, Kazakhstan.

E-mail: marat207@mail.ru, v_gulmira@mail.ru, alua.sarypbek@mail.ru

ON INVERSE STOCHASTIC RECONSTRUCTION PROBLEM

Abstract. In this paper, general reconstruction problem in the class of second-order stochastic differential equations of the Ito type is considered for given properties of motion, when the control is included in the drift coefficient. And the form of control parameters is determined by the quasi-inversion method, which provides necessary and sufficient conditions for existence of a given integral manifold. Further, the solution of the Meshchersky's stochastic problem is given, which is one of the inverse problems of dynamics and, according to the well-known Galiullin's classification, refers to the restoration problem.

It is assumed that random perturbations belong to the class of processes with independent increments. To solve the posed problem an equation of perturbed motion is drawn up by the Ito rule of stochastic differentiation. And, further, the Erugin method in combination with the quasi-inversion method is used to construct: 1) a set of control vector functions and 2) a set of diffusion matrices that provide necessary and sufficient conditions for a given second-order differential equation of Ito type to have a given integral manifold.

The linear case of a stochastic problem with drift control is considered separately. In the linear setting, in contrast to the nonlinear formulation, the conditions of solvability in the presence of random perturbations from the class of processes with independent increments coincide with the conditions of solvability in a similar linear case in the presence of random perturbations from the class of independent Wiener processes. Also considered is the scalar case of the recovery problem with drift controls.

Key Words: Ito stochastic differential equation, reconstruction problem, Meshchersky's problem, integral manifold, quasi-inversion method.

Introduction The theory of inverse problems of the dynamics of systems described by ordinary differential equations [1-6, etc.] goes back to the fundamental work of Erugin [7]. In 7, a set of ordinary differential equations is constructed that have a given integral curve. The inverse problems of constructing automatic control systems for program motion are studied in [8-11]. It should be noted that one of the general methods for solving inverse problems of dynamics in the class of ordinary differential equations, namely, the quasi-inversion method was proposed in [3].

Inverse problems of dynamics in the class of partial differential equations are studied in [12-14], and in the class of stochastic differential equations in [15-19].

1 Stochastic reconstruction problem

Let us consider the general reconstruction problem in the class of second-order Ito stochastic differential equations by the given properties of motion, when the control is included in the drift coefficient. And by the quasi-inversion method we define the form of control parameters that provides necessary and sufficient conditions for the existence of a given integral manifold. Further, the solution of Meshchersky's stochastic problem is given, which is one of the inverse problems of dynamics and it belongs to reconstruction problem according to the well-known Galiullin's classification of [1].

1.1 Problem statement. Let us consider the second-order Ito stochastic differential equation

$$\ddot{x} = f(x, \dot{x}, t) + D(x, \dot{x}, t)u + \sigma(x, \dot{x}, t)\dot{\xi}. \quad (1.1)$$

It is required to determine the vector-function $u = u(x, \dot{x}, t) \in R^r$ included in the drift coefficient for the given integral manifold

$$\Lambda(t): \lambda(x, \dot{x}, t) = 0, \quad \lambda = \lambda(x, \dot{x}, t) \in C_{x\dot{x}t}^{121}, \quad \lambda \in R^m, \quad (1.2)$$

here $C_{x\dot{x}t}^{121}$ is set of $\gamma(x, \dot{x}, t)$ functions, which are continuously differentiable with respect to x and t and twice continuously differentiable with respect to $\dot{x}$.

In other words, for the given f , D , σ and λ , the control u should be defined so that the set (1.2) is the integral set of equation (1.1).

It is assumed that the functions $f(x, \dot{x}, t)$, $D(x, \dot{x}, t)$, $\sigma(x, \dot{x}, t)$, included in the above equation, have the smoothness necessary for further reasoning and satisfy the existence and uniqueness theorem up to stochastic equivalence of the solution $(x(t)^T, \dot{x}(t)^T)^T$ of (1.1) with the initial condition $(x(t_0)^T, \dot{x}(t_0)^T)^T = (x_0^T, \dot{x}_0^T)^T$, which is continuous strictly Markov process with probability 1 [20].

Here $\{\xi_1(t, \omega), \dots, \xi_k(t, \omega)\}$ is a system of random processes with independent increments, which, following [19], can be represented as a sum $\xi = \xi_0 + \int c(y)P^0(t, dy)$ of processes Wiener process ξ_0 and Poisson process P^0 . $P^0(t, dy)$ is the number of process P^0 jumps in the interval $[0, t]$ that fall on the set dy ; $c(y)$ is a vector function that maps space R^{2n} into the space R^k of values of process $\xi(t)$ for any t .

This problem is one of the inverse problems of dynamics, and in the absence of random perturbations ($\sigma \equiv 0$) it has been sufficiently fully investigated in [1-6], and the case $\sigma \neq 0$ and $\{\xi_1(t, \omega), \dots, \xi_k(t, \omega)\}$ is a system of independent Wiener processes, as a particular form of processes with independent increments, is considered in [21].

In this paper, the quasi-inversion method is used to solve the stochastic recovery problem [3, p. 12].

By Ito rule of stochastic differentiation [20, p.204] for solving the posed problem the equation of perturbed motion

$$\dot{\lambda} = \frac{\partial \lambda}{\partial t} + \frac{\partial \lambda}{\partial x} \dot{x} + \frac{\partial \lambda}{\partial \dot{x}} f + \frac{\partial \lambda}{\partial \dot{x}} Du + \frac{\partial \lambda}{\partial \dot{x}} \sigma \dot{\xi}_0 + S_1 + S_2 + S_3, \quad (1.4)$$

is compiled. Here $S_1 = \frac{1}{2} \frac{\partial^2 \lambda}{\partial \dot{x}^2} : \sigma \sigma^T$; $S_2 = \int [\lambda(x, \dot{x} + \sigma c(y), t) - \lambda(x, \dot{x}, t) - \frac{\partial \lambda}{\partial \dot{x}} \sigma \dot{x} c(y)] dy$;

$S_3 = \int [\lambda(x, \dot{x} + \sigma c(y), t) - \lambda(x, \dot{x}, t)] \dot{P}_0(t, dy)$. Following [20], $\frac{1}{2} \frac{\partial^2 \lambda}{\partial \dot{x}^2} : D$ is a vector whose elements are the traces of the products of the matrices of the second derivatives of the corresponding elements $\lambda_\mu(x, \dot{x}, t)$ of the vector $\lambda(x, \dot{x}, t)$ with respect to components $\dot{x}$ by the matrix D

$$\frac{\partial^2 \lambda}{\partial \dot{x}^2} : D = \begin{bmatrix} tr \left(\frac{\partial^2 \lambda_1}{\partial \dot{x}^2} D \right) \\ \vdots \\ tr \left(\frac{\partial^2 \lambda_m}{\partial \dot{x}^2} D \right) \end{bmatrix}, \quad D = \sigma \sigma^T.$$

We introduce arbitrary Erugin functions [1]: an m -dimensional vector function A and a $(m \times k)$ -matrix B , with the properties $A(0, x, \dot{x}, t) \equiv 0$, $B(0, x, \dot{x}, t) \equiv 0$, such that

$$\dot{\lambda} = A(\lambda, x, \dot{x}, t) + B(\lambda, x, \dot{x}, t)\dot{\xi}, \quad (1.5)$$

takes place. Here ξ is the same process with independent increments included in (1.1) and represented as a sum of Wiener process and Poisson process [20]:

$$\xi = \xi_0 + \int c(y)P^0(t, dy) \text{ or } \dot{\xi} = \dot{\xi}_0 + \int c(y)\dot{P}^0(t, dy).$$

Based on equations (1.4) and (1.5), we obtain the relations

$$\frac{\partial \lambda}{\partial \dot{x}} Du = A - \frac{\partial \lambda}{\partial t} - \frac{\partial \lambda}{\partial x} \dot{x} - \frac{\partial \lambda}{\partial \dot{x}} f - S_1 - S_2 - S_3, \quad (1.6)$$

$$\frac{\partial \lambda}{\partial \dot{x}} \sigma = B, \quad (1.7)$$

from which you need to determine the control u and the matrix σ . To solve the problem, you need the following lemma.

Lemma 1 [3, p.12-13]. The set of all solutions of a linear system

$$H \mathcal{G} = g, \quad H = (h_{\mu\nu}), \quad \mathcal{G} = (\mathcal{G}_\nu), \quad g = (g_\mu), \quad \mu = \overline{1, m}, \quad \nu = \overline{1, n}, \quad m \leq n, \quad (1.8)$$

is determined by the expression

$$\mathcal{G} = s[HC] + H^+ g. \quad (1.9)$$

Here H is matrix has rank m . s is arbitrary scalar, $[HC] = [h_1 \dots h_m c_{m+1} \dots c_{n-1}]$ is the cross product of vectors $h_\mu = (h_{\mu\nu})$ and $c_\rho = (c_{\rho\nu})$, $\rho = \overline{m+1, n-1}$; $H^+ = H^T (HH^T)^{-1}$, H^T is the matrix transposed to H .

Denoting, by formula (1.9) of Lemma 1 from relations (1.6), (1.7), we define the required vector - function and columns, matrices in the form:

$$\text{Denoting } \tilde{D} = \frac{\partial \lambda}{\partial \dot{x}} D, \text{ by formula (1.9) from (1.6), (1.7) we define the required vector - function } u$$

and columns σ_i , $i = \overline{1, k}$ of σ in the form :

$$u = s_1 [\tilde{D}C] + (\tilde{D})^+ \left(A - \frac{\partial \lambda}{\partial t} - \frac{\partial \lambda}{\partial x} \dot{x} - \frac{\partial \lambda}{\partial \dot{x}} f - S_1 - S_2 - S_3 \right), \quad (1.10)$$

$$\sigma_i = s_2 \left[\frac{\partial \lambda}{\partial \dot{x}} C \right] + \left(\frac{\partial \lambda}{\partial \dot{x}} \right)^+ B_i, \quad i = \overline{1, k}. \quad (1.11)$$

Therefore, the following theorem is true.

Theorem 1.1 A necessary and sufficient condition that second-order Ito differential equation (1.1) has a given integral manifold (1.2) is that the control function u has the form (1.10) and the columns σ_i of diffusion matrix σ have the form (1.11).

Remark 1.1 If $c(y) \equiv 0$, then $S_2 \equiv S_3 \equiv 0$ and a solution of this problem coincides with the solution of the reconstruction problem previously considered in [21] in the presence of random perturbations from the class of independent Wiener processes.

Remark 1.2 For $m = n$, formula (1.9) takes the form $\mathcal{G} = H^{-1}g$, since in this case, the first term of the formula as a cross product of n vectors in n -dimensional space identically equals to zero $[HC] \equiv 0$, and the second term $H^+ g$ takes the form $H^{-1}g$, since for $m = n$, the rectangular matrix H becomes square matrix, and under the assumption $\det H \neq 0$ we have $H^+ = H^T (HH^T)^{-1} = H^T (H^T)^{-1} H^{-1} = H^{-1}$.

1.2 The linear case of a stochastic problem with drift control. Let a second-order Ito stochastic differential equation, linear in drift

$$\ddot{x} = E_1(t)x + E_2(t)\dot{x} + D(t)u + l_1(t) + T(t)\dot{\xi} \quad (1.12)$$

be given. It is required the control vector-function $u = u(x, \dot{x}, t) \in R^r$ by given integral manifold

$$\Lambda(t) : \lambda \equiv G_1x + G_2\dot{x} + l_2(t) = 0. \quad (1.13)$$

That is, by the given $(m \times n)$ -matrices $G_1(t), G_2(t)$ of the m -dimensional function $l(t)$, also by the given $(n \times n)$ -matrices $E_1(t), E_2(t)$, $(n \times r)$ -matrix $D(t)$ and n -dimensional function $l_1(t)$, it is required to determine the vector function $u = u(x, \dot{x}, t) \in R^r$ and $(n \times k)$ -matrix $T(t)$ so that for the constructed equation (1.12) the given properties (1.13) are an integral manifold.

In this problem the equation of perturbed motion (1.4) has the form

$$\dot{\lambda} = \dot{G}_1x + \dot{G}_2\dot{x} + \dot{l}_2(t) + \dot{G}_1\dot{x} + G_2(E_1(t)x + E_2(t)\dot{x} + D(t)u + l_1(t)) + G_2T\dot{\xi}_0. \quad (1.14)$$

On the other hand, following Erugin's method with the help of an arbitrary vector-function $A = A_1(t)\lambda$ and a matrix-function B_1 with the property $B_1(0, x, \dot{x}, t) \equiv 0$, we have

$$\dot{\lambda} = A_1(t)\lambda + B_1(\lambda, x, \dot{x}, t)\dot{\xi}. \quad (1.15)$$

Hence, relations (1.14) and (1.15) imply the equalities

$$\begin{cases} G_2Du = [A_1G_1 - \dot{G}_1 - G_2E_1]x + [A_1G_2 - G_2 - G_1 - G_2E_2]\dot{x} + A_1l_2 - G_2l_1 - \dot{l}_2, \\ G_2T = B_1. \end{cases} \quad (1.16)$$

Further, from (1.16) using Lemma 1, we have

$$u = s_1[D_1C] + (D_1)^+ g_1, \quad (1.17)$$

$$T_i = s_2[G_2C] + (G_2)^+ B_i, \quad i = \overline{1, k}, \quad (1.18)$$

here $D_1 = G_2D$, $g_1 = [A_1G_1 - \dot{G}_1 - G_2E_1]x + [A_1G_2 - G_2 - G_1 - G_2E_2]\dot{x} + A_1l_2 - G_2l_1 - \dot{l}_2$, T_i, B_i are i -th columns of matrices T and B respectively. s_1, s_2 are arbitrary scalar values. This proves the following theorem.

Theorem 1.2. A necessary and sufficient condition that second-order Ito stochastic differential equation (1.12) linear in drift, has a given linear integral manifold (1.2) is that the control parameter has the form (1.17) and the diffusion matrix has the form (1.18).

Remark 1.2. In the linear case, in contrast to the nonlinear one $S_1 \equiv S_2 \equiv S_3 \equiv 0$, the conditions of solvability in Theorem 1.2 in the presence of random perturbations from the class of processes with independent increments coincide with the conditions of solvability in a similar linear case in the presence of random perturbations from the class of independent Wiener processes [21].

1.3 Scalar case of the reconstruction problem with drift controls. Let a second-order Ito stochastic differential equation

$$\ddot{x} = f_2(x, \dot{x}, t) + \gamma_1(x, \dot{x}, t)u_1 + \gamma(x, \dot{x}, t)\dot{\xi}. \quad (1.20)$$

be given. It is required to determine the scalar function $u_1 = u_1(x, \dot{x}, t)$ for a given integral manifold

$$\lambda_2(x, \dot{x}, t) = 0, \quad \lambda_2 \in R^1. \quad (1.21)$$

In other words, for given f_2, γ, γ_1 and λ_2 define the control parameter u_1 in such a way that the set (1.21) be an integral set of equation (1.20).

According to the rule of stochastic differentiation, we compose the equation

$$\dot{\lambda}_2 = \frac{\partial \lambda_2}{\partial t} + \frac{\partial \lambda_2}{\partial x} \dot{x} + \frac{\partial \lambda_2}{\partial \dot{x}} f_2 + \frac{\partial \lambda_2}{\partial \dot{x}} \gamma_1 u_1 + \tilde{S}_1 + \tilde{S}_2 + \tilde{S}_3 + \frac{\partial \lambda_2}{\partial \ddot{x}} \gamma \dot{\xi}_0, \quad (1.22)$$

of perturbed motion; here

$$\tilde{S}_1 = \frac{1}{2} \frac{\partial^2 \lambda_2}{\partial \dot{x}^2} \gamma^2, \quad \tilde{S}_3 = \int [\lambda_2(x, \dot{x} + \gamma c(y), t) - \lambda_1(x, \dot{x}, t)] dP^0(t, dy),$$

$$\tilde{S}_2 = \int [\lambda_2(x, \dot{x} + \gamma c(y), t) - \lambda_2(x, \dot{x}, t) - \frac{\partial \lambda_2}{\partial \dot{x}} \gamma c(y)] dy.$$

We introduce arbitrary Erugin's scalar functions $a = a(\lambda_2, x, \dot{x}, t)$ and $b = b(\lambda_2, x, \dot{x}, t)$, with the properties $a(0, x, \dot{x}, t) \equiv b(0, x, \dot{x}, t) \equiv 0$ and such that

$$\dot{\lambda}_2 = a(\lambda_2, x, \dot{x}, t) + b(\lambda_2, x, \dot{x}, t) \dot{\xi}, \quad (1.23)$$

In view of (5.3) and (5.4), we arrive at the relations

$$\frac{\partial \lambda_2}{\partial x} \gamma_1 u_1 = a - \frac{\partial \lambda_2}{\partial t} - \frac{\partial \lambda_2}{\partial x} \dot{x} - \frac{\partial \lambda_2}{\partial \dot{x}} f_2 - \tilde{S}_1 - \tilde{S}_2 - \tilde{S}_3, \quad (1.24)$$

$$\frac{\partial \lambda_2}{\partial x} \gamma = b. \quad (1.25)$$

Then, by virtue of equalities (1.24) and (1.25), the control parameter u_1 and the diffusion coefficient are defined in the form

$$u_1 = \left(\frac{\partial \lambda_2}{\partial \dot{x}} \gamma_1 \right)^{-1} \left[a - \frac{\partial \lambda_2}{\partial t} - \frac{\partial \lambda_2}{\partial x} \dot{x} - \frac{\partial \lambda_2}{\partial \dot{x}} f_2 - \frac{1}{2} \frac{\partial^2 \lambda_2}{\partial \dot{x}^2} \gamma^2 - \tilde{S}_2 - \tilde{S}_3 \right], \quad (1.26)$$

$$\gamma = \left(\frac{\partial \lambda_2}{\partial \dot{x}} \right)^{-1} b. \quad (1.27)$$

Consequently, the following theorem holds.

Theorem 1.3 A necessary and sufficient condition that second-order scalar differential equation of Ito type (1.20) has a given integral manifold (1.21) is that the control parameter u_1 has the form (1.26) and diffusion coefficient has the form (1.27).

Thus, in Section 1, we obtained necessary and sufficient conditions for the solvability of the reconstruction problem with drift control in the presence of random perturbations from the class of processes with independent increments in general nonlinear case, linear case and scalar nonlinear case. The considered setting generalizes the reconstruction problem in the presence of random perturbations from the class of independent Wiener processes, previously studied in [21].

2 Meshchersky's stochastic problem

Problem statement. Find the law of change in the mass of a point, at which it describes a given trajectory under the action of given external forces [22, p.19].

Let us consider the problem of realizing the motion of a heavy point of variable mass $m(t)$ in a homogeneous gravity field, namely, vertical ascent according to the laws of change in the range y and height z

$$\Lambda(t): \begin{cases} \lambda_1(t) \equiv y - \varphi(t) = 0, \\ \lambda_2(t) \equiv z - \psi(t) = 0, \end{cases} \quad (2.1)$$

The equations of point motion [1, c.16-17] taking into account the action of random perturbing forces have the following form:

$$\begin{cases} m\ddot{y} = \dot{m}(\mu - 1)\dot{y} - mf(z, v) \frac{\dot{y}}{v} - \sigma_1(y, z, t) \dot{\xi}, \\ m\ddot{z} = \dot{m}(\eta - 1)\dot{z} - mf(z, v) \frac{\dot{z}}{v} - mg - \sigma_2(y, z, t) \dot{\xi}, \end{cases} \quad (2.2)$$

here $f(z, v)$ is medium resistance per unit mass; $v = \sqrt{\dot{y}^2 + \dot{z}^2}$ is point speed; $\mu = \mu(t)$, $v = v(t)$ are the ratio of the projections of the velocities of the changing mass and the mass of the point itself on the coordinate axis y, z .

It is required to restore the equations of motion (2.2) (that is, to determine the laws of variation of quantities μ , v and m) so that they admit a given particular motion (2.1).

The perturbed motion equations have the form

$$\begin{cases} \ddot{\lambda}_1 \equiv \ddot{y} - \ddot{\varphi}(t) = \frac{\dot{m}}{m}(\mu - 1)\dot{y} - f(z, v)\frac{\dot{y}}{v} - \frac{\sigma_1}{m}\dot{\xi} - \ddot{\varphi}(t), \\ \ddot{\lambda}_2 \equiv \ddot{z} - \ddot{\psi}(t) = \frac{\dot{m}}{m}(\eta - 1)\dot{z} - f(z, v)\frac{\dot{z}}{v} - g - \frac{\sigma_2}{m}\dot{\xi} - \ddot{\psi}(t). \end{cases} \quad (2.3)$$

Further, following Erugin's method [7], we introduce functions $A_1 = A_1(\lambda_1, \dot{\lambda}_1, \lambda_2, \dot{\lambda}_2, y, z, t)$,

$$A_2 = A_2(\lambda_1, \dot{\lambda}_1, \lambda_2, \dot{\lambda}_2, y, z, t), \quad B_1 = B_1(\lambda_1, \dot{\lambda}_1, \lambda_2, \dot{\lambda}_2, y, z, t), \quad B_2 = B_2(\lambda_1, \dot{\lambda}_1, \lambda_2, \dot{\lambda}_2, y, z, t),$$

With the properties $A_1(0, 0, 0, 0, y, z, t) \equiv A_2(0, 0, 0, 0, y, z, t) \equiv B_1(0, 0, 0, 0, y, z, t) \equiv$

$\equiv B_2(0, 0, 0, 0, y, z, t) \equiv 0$, and such that

$$\begin{cases} \ddot{\lambda}_1 = A_1 + B_1\dot{\xi}, \\ \ddot{\lambda}_2 = A_2 + B_2\dot{\xi}. \end{cases} \quad (2.4)$$

Comparison of the systems of equations (2.3) and (2.4) leads, if we exclude strictly vertical and strictly horizontal motions (i.e., $\dot{\varphi}$ and $\dot{\psi}$ are not identically zero), to relations that solve the posed Meshchersky stochastic problem

$$\begin{cases} \mu = 1 + \frac{m}{\dot{m}} \left[\frac{A_1}{\dot{y}} + \frac{f}{v} + \frac{\ddot{\varphi}}{\dot{y}} \right], \\ \eta = 1 + \frac{m}{\dot{m}} \left[\frac{A_2}{\dot{z}} + \frac{f}{v} + \frac{g}{\dot{z}} + \frac{\ddot{\psi}}{\dot{y}} \right], \\ \sigma_{1j} = mB_{1j}, \\ \sigma_{2j} = mB_{2j}. \end{cases} \quad (2.5)$$

In particular, for $\sigma_{ij} \equiv 0$ ($i, j = 1, 2$) and $A_1 \equiv A_2 \equiv B_1 \equiv B_2 \equiv 0$ conditions (2.5) coincide with conditions in the class of second-order ordinary differential equations [1, p. 17].

This research has been funded by the Science Committee of the Ministry of Education and Science of the Republic of Kazakhstan (Grant No. AP 08955847).

М.Ы. Тілеубергенов^{1,2}, Г.Қ. Василина^{1,3}, А.Т. Сарыпбек^{1,2}

¹Математика және математикалық модельдеу институты, Алматы, Қазақстан;

²Әл-Фараби атындағы Қазақ ұлттық университеті, Алматы, Қазақстан;

³Алматы энергетика және байланыс университеті, Алматы, Қазақстан

КЕРІ СТОХАСТИКАЛЫҚ ҚАЛПЫНА КЕЛТІРУ ЕСЕБІ ТУРАЛЫ

Аннотация. Аталмыш жұмыста жалпы қалпына келтіру есебі қозғалыс қасиеттері бойынша басқару қирату коэффициентіне кіргенде екінші ретті Ито түріндегі берілген стохастикалық дифференциалдық теңдеулер класында қарастырылады және квазиқайтару әдісімен интегралдық көпбейненің қажетті әрі жеткілікті шарттарын қамтамасыз ететін басқарушы параметрлер түрі анықталады. Төменде Мещерскийдің

стохастикалық есебінің шешімі келтірілген, ол динамиканың кері есептерінің бірі болып табылады және Галиуллиннің белгілі классификациясына сәйкес қалпына келтіру міндетіне жатады.

Кездейсоқ бұзылулар тәуелсіз өсуі бар процестер класына жатады деп болжанады. Берілген мәселені шешу үшін ИТО стохастикалық саралау ережесі бойынша бұзылған қозғалыс теңдеуі жасалады. Бұдан әрі еругин әдісі квазикайтару әдісімен біріктіріліп құрылады: 1) басқарушы вектор-функциялар жиыны; 2) Ито типінің екінші ретті берілген дифференциалдық теңдеудің берілген интегралдық көпбейне ие болуы үшін қажетті және жеткілікті жағдайларды қамтамасыз ететін диффузия матрицаларының жиынтығы.

Бұзу басқармасы бар стохастикалық мәселенің сызықтық жағдайы бөлек қарастырылады. Сызықтық қойылымда сызықтық емес рұқсат ету жағдайынан айырмашылығы, тәуелсіз өсуі бар процестер класынан кездейсоқ бұзылулар болған кезде, ұқсас сызықтық жағдайда, тәуелсіз винер процестері класынан кездейсоқ бұзылулар болған кезде рұқсат ету шарттарына сәйкес келеді. Сондай-ақ, бұзу басқармаларымен қалпына келтіру мәселесінің скалярлық жағдайы қарастырылады.

Түйін сөздер: ИТО-ның стохастикалық дифференциалдық теңдеуі, қалпына келтіру есебі, Мещерский есебі, интегралдық көпбейне, квазикайтару әдісі.

М. И. Тлеубергенов^{1,2}, Г.К. Василина^{1,3}, А.Т. Сарыпбек^{1,2}

¹Институт математики и математического моделирования, Алматы, Казахстан;

²Казахский национальный университет имени аль-Фараби, Алматы, Казахстан;

³Алматинский Университет энергетики и связи, Алматы, Казахстан

ОБ ОБРАТНОЙ СТОХАСТИЧЕСКОЙ ЗАДАЧЕ ВОССТАНОВЛЕНИЯ

Аннотация. В данной работе рассматривается общая задача восстановления в классе стохастических дифференциальных уравнений второго порядка типа Ито по заданным свойствам движения, когда управление входит в коэффициент сноса и методом квазиобращения определяется вид управляющих параметров, обеспечивающий необходимые и достаточные условия существования заданного интегрального многообразия. Далее приводится решение стохастической задачи Мещерского, которая является одной из обратных задач динамики и по известной классификации Галиуллина относится к задаче восстановления.

Предполагается, что случайные возмущения относятся к классу процессов с независимыми приращениями. Для решения поставленной задачи по правилу стохастического дифференцирования Ито составляется уравнение возмущенного движения. И далее методом Еругина в сочетании с методом квазиобращения строятся: 1) множество управляющих вектор-функций и 2) множество матриц диффузий, которые обеспечивают необходимые и достаточные условия того, чтобы заданное дифференциальное уравнение второго порядка типа Ито имело заданное интегральное многообразие.

Отдельно рассматривается линейный случай стохастической задачи с управлением по сносу. В линейной постановке в отличие от нелинейной условия разрешимости при наличии случайных возмущений из класса процессов с независимыми приращениями совпадают с условиями разрешимости в аналогичном линейном случае при наличии случайных возмущений из класса независимых винеровских процессов. Также рассмотрен скалярный случай задачи восстановления с управлениями по сносу.

Ключевые слова: стохастическое дифференциальное уравнение Ито, задача восстановления, задача Мещерского, интегральное многообразие, метод квазиобращения.

Information about authors:

Tleubergenov M.I., Doctor of Physics and Mathematics, Senior Researcher, Institute of Mathematics and Mathematical Modeling, Almaty, Kazakhstan, Professor, Al-Farabi Kazakh National University, Almaty, Kazakhstan, <https://orcid.org/0000-0003-1947-6714>, marat207@mail.ru;

Vassilina G.K., PhD, Senior Researcher, Institute of Mathematics and Mathematical Modeling, Senior Lecturer, Almaty University of Energy and Communications, Almaty, Kazakhstan, <https://orcid.org/0000-0003-2504-9620>, v_gulmira@mail.ru;

Sarypbek A.T., master student, Al-Farabi Kazakh National University, Almaty, Kazakhstan, Mathematics and Mathematical Modeling, Almaty, Kazakhstan, <https://orcid.org/0000-0002-5504-1972>, alua.sarypbek@mail.ru

REFERENCES

- [1] Galiullin A.S. Methods for solving inverse problems of dynamics. Moscow: Nauka, 1986. 224 p.
- [2] Galiullin A.S. Selected works in two volumes. Moscow: RUDN, 2009. Vol. I, II. 462 p.
- [3] Mukhametzyanov I.A., Mukharlyamov R.G. Equations of program motions. Moscow: Publishing House of the Peoples' Friendship University, 1986. 88 p.
- [4] Mukharlyamov R.G. Differential-algebraic equations of programmed motions of Lagrangian dynamical systems // Mechanics of Solids. 2011. T. 46. № 4. C. 534-543.

- [5] Mukharlyamov R.G., Tleubergenov M.I. Control of system dynamics and constrains stabilization, Communications in Computer and Information Science. 2017. 700. Pp. 431-442.
- [6] Llibre J., Ramirez R. Inverse Problems in Ordinary Differential Equations and Applications. Springer International Publishing Switzerland, 2016. 266 pp.
- [7] Yerugin N.P. Construction of the entire set of systems of differential equations having a given integral curve // Applied Mathematics and Mechanics. Moscow, 1952. Vol.10, vol. 6. P. 659-670.
- [8] Zhumatov S.S. Asymptotic Stability of Implicit Differential Systems in the Vicinity of Program Manifold // Ukrainian Mathematical Journal. 2014. Vol. 66(4). Pp. 625-632. DOI: 10.1007/s11253-014-0959-y
- [9] Zhumatov S.S. Exponential Stability of a Program Manifold of Indirect Control Systems // Ukrainian Mathematical Journal. Vol. 62, 6 (2010). Pp. 907-915. DOI: 10.1007/s11253-010-0399-2
- [10] Zhumatov S.S. Stability of a Program Manifold of Control Systems with Locally Quadratic Relations, Ukrainian Mathematical Journal. Vol. 61, 3 (2009). Pp. 500-509. DOI: 10.1007/s11253-009-0224-y
- [11] Zhumatov S.S. Absolute stability of a program manifold of non-autonomous basic control systems, News of the national academy of sciences of the republic of Kazakhstan. Series physico-mathematical. 2018. Vol. 6, № 322. P. 120-125. <https://doi.org/10.32014/2018.2518-1726.15>
- [12] S.A. Budochkina, V.M. Savchin, An Operator Equation with the Second Time Derivative and Hamilton-admissible Equations // Doklady Mathematics. 2016. Vol.94 (2). Pp. 487-489.
- [13] Savchin V.M., Budochkina S.A. Nonclassical Hamilton's Actions and the Numerical Performance of Variational Methodssfor Some Dissipative Problems. Springer, Cham, Communications in Computer and Information Science. 2016. Vol.678. Pp. 624-634.
- [14] Savchin V.M., Budochkina S.A. Invariance of functionals and related Euler-Lagrange equations. Russian Mathematics. 2017. Vol. 61(2). Pp. 49-54.
- [15] Tleubergenov M.I., Ibraeva G.T. Main Inverse Problem for Differential System with Generate Diffusion // Ukrainian Mathematical Journal. 2013. Vol. 65 (5). Pp. 787-792. DOI: 10.1007/s11253-013-0815-5
- [16] Tleubergenov M.I. On the inverse stochastic reconstruction problem // Differential Equations. 2014. Vol. 50 (2). Pp. 274-278. DOI: 10.1134/S0012266114020165
- [17] Tleubergenov M.I., Ibraeva G.T. Stochastic inverse problem with indirect control // Differential Equations. 2017. Vol. 53 (10). Pp. 1387-1391. DOI: 10.1134/S0012266117100172
- [18] Vassilina G.K., Tleubergenov M.I. Solution of the Problem of Stochastic Stability of an Integral Manifold by the Second Lyapunov Method // Ukrainian Mathematical Journal. 2016. Vol. 68(1). Pp. 14-28. DOI: 10.1007/s11253-016-1205-6
- [19] Tleubergenov M.I., Ibraeva G.T. On the Solvability of the Main Inverse Problem for Stochastic Differential Systems // Ukrainian Mathematical Journal. 2019. Vol.71 (1). Pp. 157-165. DOI: 10.1007/s11253-019-01631-w
- [20] Pugachev V.S., Sinitsyn I.N. Stochastic differential systems. Analysis and filtering. Moscow: Nauka, 1990. 632 p.
- [21] Tleubergenov M.I. On the inverse problem of reconstruction stochastic differential systems // Differential Equations. M., 2001. Vol. 37, No 5. P. 714-716.
- [22] Meshchersky I.V. Works on the mechanics of bodies of variable mass. Moscow, Leningrad: Gostekhizdat, 1952. 280 p.

Publication Ethics and Publication Malpractice
in the journals of the National Academy of Sciences of the Republic of Kazakhstan

For information on Ethics in publishing and Ethical guidelines for journal publication see <http://www.elsevier.com/publishingethics> and <http://www.elsevier.com/journal-authors/ethics>.

Submission of an article to the National Academy of Sciences of the Republic of Kazakhstan implies that the described work has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <http://www.elsevier.com/postingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere in the same form, in English or in any other language, including electronically without the written consent of the copyright-holder. In particular, translations into English of papers already published in another language are not accepted.

No other forms of scientific misconduct are allowed, such as plagiarism, falsification, fraudulent data, incorrect interpretation of other works, incorrect citations, etc. The National Academy of Sciences of the Republic of Kazakhstan follows the Code of Conduct of the Committee on Publication Ethics (COPE), and follows the COPE Flowcharts for Resolving Cases of Suspected Misconduct (http://publicationethics.org/files/u2/New_Code.pdf). To verify originality, your article may be checked by the Cross Check originality detection service <http://www.elsevier.com/editors/plagdetect>.

The authors are obliged to participate in peer review process and be ready to provide corrections, clarifications, retractions and apologies when needed. All authors of a paper should have significantly contributed to the research.

The reviewers should provide objective judgments and should point out relevant published works which are not yet cited. Reviewed articles should be treated confidentially. The reviewers will be chosen in such a way that there is no conflict of interests with respect to the research, the authors and/or the research funders.

The editors have complete responsibility and authority to reject or accept a paper, and they will only accept a paper when reasonably certain. They will preserve anonymity of reviewers and promote publication of corrections, clarifications, retractions and apologies when needed. The acceptance of a paper automatically implies the copyright transfer to the National Academy of Sciences of the Republic of Kazakhstan.

The Editorial Board of the National Academy of Sciences of the Republic of Kazakhstan will monitor and safeguard publishing ethics.

(Правила оформления статьи для публикации в журнале смотреть на сайтах:

www.nauka-nanrk.kz

<http://physics-mathematics.kz/index.php/en/archive>

ISSN 2518-1726 (Online), ISSN 1991-346X (Print)

Редакторы: *М. С. Ахметова, Д. С. Аленов, А. Ахметова*
Верстка на компьютере *А.М. Кульгинбаевой*

Подписано в печать 08.02.2021.
Формат 60x88¹/₈. Бумага офсетная. Печать – ризограф.
6,75 п.л. Тираж 300. Заказ 1.