On Reverse Super Edge Magic Total Labeling of Subdivided Trees

Abdul Raheem

Department of Mathematics,

National University of Singapore, Singapore 119076

rahimciit7@gmail.com

Abstract

A reverse edge magic total labeling of a graph G is a one-to-one map $\lambda : V(G) \cup E(G) \rightarrow \{1, 2, \dots, |V(G) \cup E(G)|\} = [1, |V(G) \cup E(G)|]$ with the property that there is an integer constant k such that $\{\lambda(xy) - (\lambda(x) + \lambda(y))/xy \in V(G)\} = k$. If $(V(G)) = [1, |V(G)|]$ then the reverse edge magic labeling is called reverse super edge-magic labeling. In this paper we will formulate the reverse edge magic labeling of two subclasses of trees.

Keywords: star, subdivision of star, reverse super edge magic labeling

Introduction

All graphs in this paper are finite, undirected, simple and planar. The graph G has the vertex set $V(G)$ and the edge set $E(G)$. A (v, e) -graph G is a graph such that $|V(G)| = v$ and $|E(G)| = e$. A general reference for graph-theoretic ideas can be seen in [1,2]. A labeling (or valuation) of a graph is a map that converts graph elements into numbers (usually to positive or non-negative integers). In this paper, the domain will be the set of all vertices and edges and such a labeling is known as total labeling. Some labeling alternatively uses either the vertex-set or the edge-set and we shall address them as vertex-labeling or edge-labeling, respectively.

Definition 1.1. A reverse edge magic total labeling of a graph G is a one-to-one map $\lambda : V(G) \cup E(G) \rightarrow \{1, 2, \dots, |V(G) \cup E(G)|\} = [1, |V(G) \cup E(G)|]$ with the property that there is an integer constant k such that $\{\lambda(xy) - (\lambda(x) + \lambda(y))/xy \in V(G)\} = k$.

Definition 1.2. λ is called the reverse super edge magic total labeling and G is known as a reverse super edge magic total graph. Enomoto et al. [3] proposed the following conjecture,

Conjecture 1.1. *Every tree admits a super edge magic total labeling.*

In favor of this conjecture, many authors have considered a super edge magic total labeling for different classes of trees. For detailed studies the reader can see [4–14].

Definition 1.3. Let $n_i \geq 1, 1 \leq i \leq r$ and $r \geq 2$. A subdivided star $Sb(n_1, n_2, \dots, n_r)$ is a tree obtained by inserting $n_i - 1$ nodes to each of the i th edge of the star $K_{1,r}$. Let us define the set of nodes and edges as follows,

$V(G) = \{c\} \cup \{x_i^{l_i} | 1 \leq i \leq r; 1 \leq l_i \leq n_i\}$ and $E(G) = \{cx_i^{l_i} | 1 \leq i \leq r\} \cup \{x_i^{l_i}x_i^{l_i+1} | 1 \leq i \leq r; 1 \leq l_i \leq n_i - 1\}$.

However, the investigation of different results related to a reverse super edge magic total labeling of the subdivided star $Sb(n_1, n_2, \dots, n_r)$ for $n_1 \neq n_2, \dots, \neq n_r$ is still an open problem. In this paper, we will formulate a reverse super edge magic total labeling of the subclasses of

subdivided stars denoted by $Sb(mn, mn, mn, 2mn, n_6, n_r, \dots, n_r)$ and $Sb(mn, mn, 2n, 2n + 2, 4n + 3, n_6, \dots, n_r)$ under certain conditions.

Let us consider the following Lemma which we will use frequently in the main theorems.

Lemma 1.1. A graph with vertices v and e edges is reverse super edge magic total labeling if and only if there exists a bijective function $\lambda: V(G) \rightarrow [1, v]$ such that the set consists of consecutive integers. In such a case, λ extends to a reverse super edge magic total labeling of G with magic constant $k = 2v - s - 1$, where $s = \max(S)$.

2. Main Results

In this section, we will prove the main results related to a reverse super edge magic total labeling of the more generalized subclasses of subdivided trees.

Theorem 2.1. The graph $G \cong Sb(n, n, n, n, 2n, n_6, \dots, n_r)$ admits the reverse edge magic total labeling for any odd $n \geq 3$, $r \geq 6$, $n_p = 2^{p-4}n - 2p + 11$ and $6 \leq p \leq r$

Proof Let $v = |V(G)|$ and $e = |E(G)|$ then

$$v = 6n + 1 + \sum_{t=6}^r [2^{t-4}n - 2t] \text{ and } e = v - 1$$

Let us define $\lambda: V(G) \rightarrow [1, v]$ as follows,

$$\lambda(c) = (4n + 2) + \sum_{t=6}^r [2^{t-5}n - t + 6]$$

For odd $1 \leq l \leq n_i$, where $1 \leq i \leq 5$ and for $1 \leq i \leq r$:

$$\lambda(w) = \begin{cases} \frac{l_1 + 1}{2}, & \text{if } w = x_1^{l_1} \\ n + 1 - \frac{l_2 - 1}{2}, & \text{if } w = x_2^{l_2} \\ (n + 2) + \frac{l_3 - 1}{2}, & \text{if } w = x_3^{l_3} \\ 2(n + 1) + \frac{l_4 - 1}{2}, & \text{if } w = x_4^{l_4} \\ (3n + 2) - \frac{l_5 - 1}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$$\lambda(x_i^{l_i}) = (3n + 2) + \sum_{t=6}^i [2^{t-5}n - t + 6] - \frac{l_i - 1}{2} \text{ Accordingly.}$$

For even $1 \leq l \leq n_i$ and $\gamma = (3n + 2) + \sum_{t=6}^r [2^{t-6}n + 1]$, where $1 \leq i \leq 5$ and for $6 \leq i \leq r$:

$$\lambda(w) = \begin{cases} (\gamma + 1) \frac{l_1 - 2}{2}, & \text{if } w = x_1^{l_1} \\ (\gamma + n - 1) - \frac{l_2 - 2}{2}, & \text{if } w = x_2^{l_2} \\ (\gamma + n + 1) + \frac{l_3 - 2}{2}, & \text{if } w = x_3^{l_3} \\ (\gamma + 2n - 1) + \frac{l_4 - 2}{2}, & \text{if } w = x_4^{l_4} \\ (\gamma + 3n - 1) - \frac{l_5 - 2}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$\lambda(x_i^{l_i}) = (\gamma + 3n - 1) + \sum_{t=6}^i [2^{t-5}(3n) - 2t + 11] - \frac{l_i - 2}{2}$ Accordingly.

By using the above scheme of labeling, we get the set of edge-sums consecutive integer sequence $S = [\gamma + 2, \gamma + 1 + e]$. Therefore, by Lemma 1.1 can be extended to a reverse edge magic total labeling with magic constant $k = (3n - 2) + \sum_{t=6}^i [2^{t-6}(3n) - 2t + 10]$.

Theorem 2.2. The graph $G \cong T(3n, 3n, 3n, 3n, 6n, n_6, \dots, n_r)$ admits the reverse edge magic total labeling with $k = 2v - s - 1$ for any odd $n \geq 3, r \geq 6, n_p = 2^{t-6}(3n) - 2t + 11$ and $6 \leq p \leq r$

Proof

Let $v = |V(G)|$ and $e = |E(G)|$ then

$$v = (18n + 1) + \sum_{t=6}^r [2^{t-4}(3n) - 2t + 11] \text{ and } e = v - 1$$

Let us define $\lambda: V(G) \rightarrow [1, v]$ as follows,

$$\lambda(c) = (12n + 2) + \sum_{t=6}^r [2^{t-5}(3n) - t + 6]$$

For odd $1 \leq l \leq n_i$, where $1 \leq i \leq 5$ and for $1 \leq i \leq r$:

$$\lambda(w) = \begin{cases} \frac{l_1 + 1}{2}, & \text{if } w = x_1^{l_1} \\ 3n + 1 - \frac{l_2 - 1}{2}, & \text{if } w = x_2^{l_2} \\ (3n + 2) + \frac{l_3 - 1}{2}, & \text{if } w = x_3^{l_3} \\ (6n + 2) - \frac{l_4 - 1}{2}, & \text{if } w = x_4^{l_4} \\ (9n + 2) - \frac{l_5 - 1}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$\lambda(x_i^{l_i}) = (9n + 2) + \sum_{t=6}^i [2^{t-5}(3n) - t + 6] - \frac{l_i - 1}{2}$ Accordingly.

For even $1 \leq l \leq n_i$ and $\gamma = (9n + 2) + \sum_{t=6}^r [2^{t-5}(3n) - t + 6]$, where $1 \leq i \leq 5$ and for $6 \leq i \leq r$:

$$\lambda(w) = \begin{cases} (\gamma + 1) \frac{l_1 - 2}{2}, & \text{if } w = x_1^{l_1} \\ (\gamma + 3n - 1) - \frac{l_2 - 2}{2}, & \text{if } w = x_2^{l_2} \\ (\gamma + 3n + 1) + \frac{l_3 - 2}{2}, & \text{if } w = x_3^{l_3} \\ (\gamma + 6n - 1) - \frac{l_4 - 2}{2}, & \text{if } w = x_4^{l_4} \\ (\gamma + 9n - 1) - \frac{l_5 - 2}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$$\lambda(x_i^{l_i}) = (\gamma + 9n - 1) + \sum_{t=6}^i [2^{t-5}(3n) - 2t + 11] - \frac{l_i - 2}{2} \text{ Accordingly.}$$

By using the above scheme of labeling, we get the set of edge-sums consecutive integer sequence $S = [\gamma + 2, \gamma + 1 + e]$. Therefore, by Lemma 1.1 can be extended to a reverse edge magic total labeling with magic constant $k = (9n - 2) + \sum_{t=6}^i [2^{t-6}(3n) - t + 5]$.

Theorem 2.3 The graph $G \cong Sb(mn, mn, mn, 2mn, n_6, \dots, n_r)$ admits the reverse edge magic total labeling with $k = 2v - s - 1$ for any odd $n \geq 3$, $r \geq 6$, $n_p = 2^{p-4}kn - 2p + 11$ and $6 \leq p \leq r$

Proof

Let $v = |V(G)|$ and $e = |E(G)|$ then

$$v = (6kn + 1) + \sum_{t=6}^r [2^{m-4}kn - 2m + 11] \text{ and } e = v - 1$$

Let us define $\lambda: V(G) \rightarrow [1, v]$ as follows,

$$\lambda(c) = (4kn + 2) + \sum_{t=6}^r [2^{m-5}kn - m + 6]$$

For odd $1 \leq l \leq n_i$, where $1 \leq i \leq 5$ and for $1 \leq i \leq r$:

$$\lambda(w) = \begin{cases} \frac{l_1 + 1}{2}, & \text{if } w = x_1^{l_1} \\ mn + 1 - \frac{l_2 - 1}{2}, & \text{if } w = x_2^{l_2} \\ (mn + 2) + \frac{l_3 - 1}{2}, & \text{if } w = x_3^{l_3} \\ 2(mn + 2) - \frac{l_4 - 1}{2}, & \text{if } w = x_4^{l_4} \\ (3mn + 2) - \frac{l_5 - 1}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$$\lambda(x_i^{l_i}) = (3mn + 2) + \sum_{t=6}^i [2^{t-5}mn - m + 6] - \frac{l_i - 1}{2} \text{ Accordingly.}$$

For even $1 \leq l \leq n_i$ and $\gamma = (3kn + 2) + \sum_{t=6}^r [2^{t-6} 2kn - (m - 6)]$, where $1 \leq i \leq 5$ and for $6 \leq i \leq r$:

$$\lambda(w) = \begin{cases} (\gamma + 1) \frac{l_1 - 2}{2}, & \text{if } w = x_1^{l_1} \\ (\gamma + mn - 1) - \frac{l_2 - 2}{2}, & \text{if } w = x_2^{l_2} \\ (\gamma + mn + 1) + \frac{l_3 - 2}{2}, & \text{if } w = x_3^{l_3} \\ (\gamma + 2mn - 1) - \frac{l_4 - 2}{2}, & \text{if } w = x_4^{l_4} \\ (\gamma + 3mn - 1) - \frac{l_5 - 2}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$$\lambda(x_i^{l_i}) = (\gamma + 3mn - 1) + \sum_{t=6}^i [2^{t-5}(mn) - 2m + 11] - \frac{l_i - 2}{2} \text{ Accordingly.}$$

By using the above scheme of labeling, we get the set of edge-sums consecutive integer sequence $S = [\gamma + 2, \gamma + 1 + e]$. Therefore, by Lemma 1.1 can be extended to a reverse edge magic total labeling with magic constant $k = (3mn - 2) + \sum_{t=6}^i [2^{t-6}(mn) - t + 10]$.

Theorem 2.4 The graph $G \cong Sb(mn, mn, 2n, 2n + 2, 4n + 3, n_6, \dots, n_r)$ admits the reverse edge magic total labeling with $k = 2v - s - 1$, for any odd $n \geq 3$, $r \geq 6$, $n_p = 2^{p-5}(4n + 2) + 1$ and $6 \leq p \leq r$

Proof

Let $v = |V(G)|$ and $e = |E(G)|$ then

$$v = [(2k + 8)n + 6] + \sum_{t=6}^r [2^{m-5}(4n + 2) + 1] \text{ and } e = v - 1$$

Let us define $\lambda: V(G) \rightarrow [1, v]$ as follows,

$$\lambda(c) = [(2k + 4)n + 4] + \sum_{t=6}^r [2^{m-6}(4n + 2) + 1]$$

For odd $1 \leq l \leq n_i$, where $1 \leq i \leq 5$ and for $6 \leq i \leq r$:

$$\lambda(w) = \begin{cases} \frac{l_1 + 1}{2}, & \text{if } w = x_1^{l_1} \\ mn + 1 - \frac{l_2 - 1}{2}, & \text{if } w = x_2^{l_2} \\ (mn + 2) + \frac{l_3 - 1}{2}, & \text{if } w = x_3^{l_3} \\ (m + 2)n + 2 - \frac{l_4 - 1}{2}, & \text{if } w = x_4^{l_4} \\ (m + 2)n + 4 - \frac{l_5 - 1}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$\lambda(x_i^{l_i}) = [(2k + 4)n + 4] + \sum_{t=6}^r [2^{t-6}(4n + 2) + 1] \frac{l_i-1}{2}$ Accordingly.

For even $1 \leq l \leq n_i$ and $\gamma = (3kn + 2) + \sum_{t=6}^r [2^{t-6}2kn - (m - 6)]$, where $1 \leq i \leq 5$ and for $6 \leq i \leq r$:

$$\lambda(w) = \begin{cases} (\gamma + 1) \frac{l_1 - 2}{2}, & \text{if } w = x_1^{l_1} \\ (\gamma + mn - 1) - \frac{l_2 - 2}{2}, & \text{if } w = x_2^{l_2} \\ (\gamma + mn + 1) + \frac{l_3 - 2}{2}, & \text{if } w = x_3^{l_3} \\ (\gamma + (m + 2)n - 1) - \frac{l_4 - 2}{2}, & \text{if } w = x_4^{l_4} \\ (\gamma + (m + 2)n + 2) - \frac{l_5 - 2}{2}, & \text{if } w = x_5^{l_5} \end{cases}$$

$\lambda(x_i^{l_i}) = [\gamma + (m + 4)n + 2] + \sum_{t=6}^i [2^{t-6}(4n + 2) + 1] - \frac{l_i-2}{2}$ Accordingly.

By using the above scheme of labeling, we get the set of edge-sums consecutive integer sequence $S = [\gamma + 2, \gamma + 1 + e]$. Therefore, by Lemma 1.1 can be extended to a reverse edge magic total labeling with magic constant $k = (mn + 4) + \sum_{t=6}^i [2^{t-6}(4n + 2)]$.

3. Conclusion

In this paper, we have proved that the following subclasses of subdivided stars admit reverse super edge magic total labeling,

- For $n \geq 3$ and $k \geq 1$ are odd, $r \geq 6$, $Sb(mn, mn, mn, 2mn, n_6, \dots, n_r)$ with $n_p = 2^{p-4} - 2p + 11$ for $6 \leq p \leq r$.
- For $n \geq 3$ and $k \geq 1$ are odd, $r \geq 6$, $Sb(mn, mn, 2n, 2n + 2, 4n + 3, n_6, \dots, n_r)$ with $n_p = 2^{p-5}(4n + 2) + 1$ for $6 \leq p \leq r$.

The problem is still open for the remaining subclasses of subdivided stars with different combinations of m and n .

References

- [01] West DB. *An introduction to graph theory*. New Jersey: Prentice-Hall, 1996.
- [02] Gallian JA. A dynamic survey of graph labeling. *Electron J Combin*. 2017;17:42–45.
- [03] Enomoto H, Lladó, AS, Nakamigawa T, Ringel G. Super edge-magic graphs, *SUT J Math*. 1998;34:105–109.
- [04] Baskoro ET, Sudarsana, IW, Cholily YM, How to construct new super edge-magic graphs from some old ones. *J Indones Math Soc (MIHIM)*. 2005;11(2):155–162.
- [05] Figueroa-Centeno RM, Ichishima R, Muntaner-Batle FA. On super edge-magic graph. *Ars Combinatori*. 2002;64: 81–95.
- [06] Javaid M, Hussain M, Ali K, Dar KH. Super edge-magic total labeling on w -trees. *Utilitas Math*. 2011;86:183–191.

- [07] Ngurah AAG, Simanjuntak R, Baskoro ET. On (super) edge-magic total labeling of subdivision of $K_{1,3}$. *SUT J Math*. 2007;43:127–136.
- [08] Salman ANM, Ngurah, AAG, Izzati N. On super edge-magic total labeling of a subdivision of a Star S_n . *Utilitas Mathematica*. 2010;81:275–284.
- [09] Sharief BS. Reverse super edge magic labeling on w-trees. *Int J Comput Eng Res Trends*. 2015;2(11):719–721.
- [10] Sharief BS. Madhusudhan RK. Reverse magic strength of festoon trees. *Ital J Pure Appl Math*. 2014;33:191–200.
- [11] Raheem AM, Javaid M, Baig AQ. On antimagicness of subdivided stars. *Discuss Math Graph Theory*. 2015;35:663–673.
- [12] Raheem A, Baig AQ. Antimagic labeling of the union of subdivided stars, *TWMS J Appl Eng Math*. 2016;6:244–250.
- [13] Raheem A, Baig AQ, Javaid M. On (a,d)-EAT labeling of subdivision of $K_{1,r}$. *J Inf Optim Sci*. 2018;39(3):643–650.
- [14] Venkata RS, Sharief BS. On reverse super edge-magic labeling of graphs, *Int Rev Pure Appl Math*. 2010;6(1):181–188.