Overview on chromatographic and potentiometric based approaches for pK_a determination of sparingly soluble substances

Natalija Nakov^{1*}, Jelena Acevska¹, Katerina Brezovska¹, Aneta Dimitrovska¹

¹*Institute of Applied Chemistry and Pharmaceutical Analysis, Faculty of Pharmacy,
Ss. Cyril and Methodius University, 1000 Skopje, Republic of Macedonia*

Received: February 2018; Accepted: April 2018

Abstract

The dissociation constant (pK_a) is one of the most important physicochemical property of the compounds and it is of great importance for the pharmaceutical industry. Many of the active pharmaceutical ingredients are sparingly soluble in water or have no chromophore, thus the conventional *UV/Vis* spectrometry and potentiometry methods are not applicable. This article presents an overview of several approaches based on RP-HPLC and cosolvent/surfactant potentiometry that have been recently published for the pK_a determination of sparingly soluble active pharmaceutical ingredients. Additionally, different calculation methods that have been used for the estimation of the aqueous dissociation constant are summarized.

Keywords: dissociation constant, pK_a , sparingly soluble substances, determination, reverse-phase liquid chromatography, potentiometry, surfactants

Introduction

The dissociation constant (pK_a) is one of the primary physicochemical property of active pharmaceutical ingredients. The accurate measurement of the pK_a value is of great importance to a wide range of research fields. The pharmacokinetics (ADMET: adsorption, distribution, metabolism, excretion and toxicity) of active pharmaceutical ingredients depend on their pK_a value. The determination of the pK_a value during the early stage studies could provide proper selection of a candidate for active pharmaceutical ingredient among the large number of compounds produced by combinatorial chemistry. The knowledge of the pK_a value for the analyte and its related impurities is essential for analytical method development; providing ruggedness and faster method optimization

(Avdeef, 2001; Kerns & Di, 2004; Manallack, 2007).

Several techniques are used for determination of the dissociation constant. Computational methods are valuable tool for pK_a calculation, especially for large set of compounds present in the screening stage of the drug candidate. There are several computer programs (such as pK_a Pallas program, ACD/ pK_a , CSpKa Predictor, ADME/Tox WEB, etc.) which are based on different theoretical approaches for predicting the pK_a value. It should be noted that the accuracy of the computational methods depends only on the model used to perform the calculations. However, computational methods could give unreliable results for sparingly soluble compounds (Babic et al., 2007; Reijenga et al., 2013; Špirtović-Halilović & Završnik, 2010).

Traditionally, *UV/Vis* spectrometry and potentiometry have been standard methods for the pK_a determination (Erdemgli et al., 2007). The *UV/Vis*

* natalijan@ff.ukim.edu.mk

spectrometric method offers great precision, but it hinges on a pH dependent spectrum of the compound in question. It means that a presence of a chromophore close to the ionization site in the molecule is needed. Additional requirement of this technique is that the tested compound should be pure or its impurities should not absorb in the UV/Vis range, since overlapping spectra could be obtained (Babic et al., 2007; Demiray et al., 2012; Pandey, 2013; Reijenga et al., 2013). Potentiometry is the method of choice for the pK_a determination of water soluble compounds. This technique offers great precision, accuracy and fast determination. The major drawback is the requirement of larger quantity of pure compound having a water solubility of at least 10^{-4} M (Babic et al., 2007; Narasimham et al., 2011; Reijenga et al., 2013).

In the last twenty years, there is a growing interest for the reverse-phase high performance liquid chromatography (RP-HPLC) as a method for the pK_a determination, based on the relationship between the capacity factor and the pH value of the mobile phase (Babic et al., 2007; Manderscheid & Eichinger, 2003; Uhrova et al., 1997; Wiczling et al., 2004). The increased application of this technique for the pK_a determination is due to the several advantages, including very small quantity of sample is required, the solubility is not a concern and the purity of the compound is not a question.

Many active pharmaceutical ingredients are sparingly soluble in water, thus the pK_a determination could not be performed in pure water, and thus water-organic mixtures are needed. It is well established that the increase of the organic content in water-organic mixtures leads to suppression of the basic analyte pK_a and to an increase in the acidic analyte pK_a (Kazakevich & Lobrutto, 2007). In such case, the obtained pK_a values are referred as apparent pK_a ($^s pK_a$) values and consequently an estimation of the aqueous pK_a ($^w pK_a$) value is needed (Babic et al., 2007; Manderscheid & Eichinger, 2003; Uhrova et al., 1997).

Generally, the RP-HPLC technique and modifications of the potentiometric titration are the methodological approaches of choice for the pK_a determination of sparingly soluble compounds. This review article, describes the principles, advantages and drawbacks on several approaches based on RP-HPLC and potentiometry which are recently used for this purpose. Additionally, an overview of different approaches for estimation of the aqueous pK_a values is given.

RP-HPLC approaches for pK_a determination

Isocratic RP-HPLC approach

The HPLC methodological approach of the pK_a determination is based on the assumption that the neutral and ionic forms of the compound have different chromatographic behavior. The determination of pK_a

value by means of HPLC was first reported by Horvath et al. (1977) and by Van de Venne et al. (1978). The most widely used HPLC procedure for the pK_a determination is the isocratic HPLC approach (Babic et al., 2007; Manderscheid & Eichinger, 2003; Uhrova et al., 1997). This approach involves a series of isocratic experiments at different pH value of the mobile phase, using at least three different concentration of the organic modifier (acetonitrile or methanol) in the mobile phase (Demiray et al., 2012; Erdemgli et al., 2007; Gumustas et al., 2010; Huo et al., 2013; Kazakevich & Lobrutto, 2007). More accurate results could be obtained if lower organic modifier concentration is used, thus the organic modifier concentration in the mobile phase should not be more than 40%. The selected organic modifier concentration in the mobile phase should be able to elute the neutral form in a reasonable time, while providing a value for the retention factor (k') of the ionized form of at least one. The $^s pK_a$ value could be determined using nonlinear regression analysis software, such as NLREG program (Demiralay et al., 2009; Gumustas et al., 2010; Şanlı et al., 2010) where the plot k' versus pH value at a particular organic content concentration is fitted to a nonlinear symmetrical sigmoidal model. The inflection point of the curve corresponds to the $^s pK_a$ value of the compound.

The drawback of the isocratic concept is that long retention times could be observed using low organic content mobile phases. Additionally, 8 to 10 experiments should be performed at each organic content composition, which makes this concept a time-consuming (Babic et al., 2007; Wiczling et al., 2004). Beforehand the column stability was the limiting factor for the range of the pK_a values that could be determined. However, nowadays there is variety of stationary phases with wide pH stability range that could fit this purpose, so the chromatographic column stability is not an issue anymore. The isocratic RP-HPLC approach is applicable for simultaneous pK_a determination of small number of compounds, thus this approach is not a method of choice for early stage screening of drug candidates.

Gradient RP-HPLC approach

The methodology for the gradient RP-HPLC determination of pK_a is based on two organic solvent gradient runs and one pH gradient RP-HPLC run (Kaliszan & Wiczling, 2010; Wiczling et al., 2004; Wiczling et al., 2006; Wiczling et al., 2008). The initial organic solvent gradient experiments at two different gradient times should be carried out at fixed pH value, where the ionization of the compound is suppressed (pH 3.0 for acidic and pH 10.5 for basic compounds). These organic solvent gradient runs are intended to define the organic solvent concentration, needed for providing appropriate retention time for completing the pH gradient experiment. Afterwards, a programmed pH gradient experiment is carried out at the chosen organic solvent

concentration in the mobile phase. The pH gradient should be started at pH value that ensures complete suppression of the ionization of the compounds and it ends at a pH value that provides their full ionization. This means that the pH gradient experiment for acidic compound should be started at pH 3.0 and finished at pH 10.5. At the end, one isocratic experiment at pH value that ensures a complete ionization of the compound is needed in order to obtain the retention of the ionized form of the compound.

The described procedure has the advantage of fast pK_a determination of large set of substances in a mixture. Additional increase of the number of analyzed compounds in individual run could be gained with the replacement of the UV detector with time-of-flight mass spectrometry (TOF-MS) detector. Recently, a gradient RP-HPLC-TOF-MS method has been published for determination of the hydrophobicity and dissociation constant of mixture of 40 compounds (Wiczling et al., 2014). The same research team expands the applicability of the proposed method on a diverse group of 161 chemical compounds (Wiczling et al., 2015). Some of the pK_a values derived with this methodology differ from the literature data; therefore, the gradient RP-HPLC approach is useful for a rough, preliminary estimation of the pK_a values of drug candidates.

Solely aqueous RP-HPLC approach

Recently, a new RP-HPLC approach based on solely aqueous mobile phase was published (Volna et al., 2017). The mobile phase is composed of 100% modified Britton-Robinson buffer, where the pH of the eluent is set by mixing the Britton-Robinson buffer with diethylamine (0.1% v/v) in the HPLC pump. In case of very lipophilic compounds, 5% methanol could be added to the mobile phase. This methodology is applicable for compounds having wide range of lipophilicity (log P from -2.26 to 2.26). The use of the solely aqueous mobile phase avoids the need of pK_a determination at several sets of measurements in eluent at different organic solvent content, which speeds up the procedure. Another advantage of this methodology is that there is no need to use more complex calculation, such as Yasuda-Shedlovsky extrapolation (Shedlovsky, 1962), for the estimation of the ${}^w pK_a$ value. Also, it should be emphasized that good agreement between the pK_a values derived with this procedure and literature values was encountered for acidic compounds. Hence, the solely aqueous RP-HPLC approach presents a promising approach for fast and relatively simple approach for pK_a determination of large number of compounds.

Potentiometric approaches for pK_a determination

The potentiometry is a well-established method for

pK_a determination. The simplicity and the low cost of the potentiometric titrations have made it one of the most commonly used methods for pK_a determination. However, the poor solubility of the compounds impedes the traditional potentiometric determination, thus alternative approaches are needed.

Cosolvent (mixed-solvent) approach

The cosolvent approach is based on the measurements of the apparent dissociation constant in different ratios of organic solvent/water mixtures where the ${}^w pK_a$ value is obtained by extrapolation (Avdeef et al., 1999; Fallaven & Schapoval, 1997; Takacs-Novak et al., 1997). According to the reported literature data, the potentiometric titrations could be performed by use of different organic solvents, such as: methanol (Erdeemly et al., 2007; Fallavena & Schapoval, 1997; Herrero-Martinez et al., 2008; Takacs-Novak et al., 1997), dioxane (Fallavena & Schapoval, 1997), acetone (Fallavena & Schapoval, 1997; Takacs-Novak et al., 1997), isopropyl alcohol (Rafols et al., 1997). However, methanol has the widest application for the purpose because the solvation effect of this solvent is closest to water (Avdeef, 1993a; Avdeef, 1993b; Takacs-Novak et al., 1997; Völgyi et al., 2007).

However, some compounds could not be dissolved in single organic solvent/water mixture. To overcome this problem, Völgyi and coworkers (2007) proposed a multicomponent cosolvent (MDM-water mixture) approach. The MDM-water mixture consists of 60% v/v MDM stock solution (equal volumes of methanol, dioxin and acetonitrile) and 40% v/v water (and 0.15 M KCl, for ionic strength adjustment), and this mixture improves the solubility of the compounds. Consequently, the pK_a measurements could be performed using lower proportions of organic solvent which allows sidestep of the long-distance extrapolations to zero organic solvent. Hence, more reliable results could be obtained. The applicability of the MDM-water mixture approach was confirmed through the pK_a determination of compounds having diverse structure (Völgyi et al., 2007; Zapala, 2011).

The apparent ionization constant obtained with the cosolvent approach could be deduced graphically from the Bjerrum plots, n_h^- vs pH (Aldeef, 2003; Kraft, 2003), where the Bjerrum function, n_h^- , is the average number of bound protons at particular pH and the pK_a value is read from the Bjerrum plot at $n_h^-=0.5$.

Surfactant based approach

Generally, the cosolvent concept is more time-consuming. Recently, an alternative approach using nonionic surfactants is proposed (Ravichandiran et al., 2011). In this study, the desired solubility of the poorly

soluble compounds was achieved using low concentrations (2-5%) of Tween 80, Chemophol and Labrasol. The authors of this concept demonstrated that the ionization behavior of several compounds remains unaffected by the use of the nonionic surfactants, with the same level of accuracy without a need of extrapolations to pure water.

However, several literature reports revealed that cationic (cetyltrimethylammonium bromide), anionic (sodium dodecyl sulfate) and non-ionic (Triton X100) micellar solutions shift the dissociation equilibrium as compared to those in pure water (Jaiswal et al., 2005; Nemcova et al., 2009; Yuangin et al., 2002). In case of cationic and anionic surfactant solution, electrostatic effect and polarity effect have an impact on dissociation equilibrium, as for non-ionic surfactants the polarity is the dominant factor. Generally, the pK_a shifts are smaller when nonionic surfactants are used. This suggests that measurements using different concentration of nonionic surfactant should be performed in order to perceive their effect on the dissociation constant.

Estimation of aqueous pK_a value

The literature data suggest that several approaches could be used for the estimation of the ${}^w pK_a$ value. One of the most commonly used approach for the ${}^w pK_a$ value estimation is based on the linear relationship between the ${}^s pK_a$ values and the volume fraction of the organic solvent used for the determination. The intercept of the linear equation (extrapolation to pure water mobile phase) presents the ${}^w pK_a$ value of the compound.

The second method known as Yasuda-Shedlovsky extrapolation establishes linear relationship between the apparent pK_a and the solvent dielectric constant (Sheldovsky, 1962), following the equation:

$${}^s pK_a + \log[H_2O] = \frac{\alpha}{\epsilon} + \beta \quad \text{Eq.1}$$

where ϵ denotes the solvent dielectric constant, $[H_2O]$ is the molar concentration of water in the mobile phase, and α and β are constants.

The plot of ${}^s pK_a$ value + $\log [H_2O]$ versus $1/\epsilon$ is extrapolated to a value corresponding to pure water ($\epsilon=78.3$, $[H_2O] = 55.5$). The ${}^w pK_a$ value could be calculated from the equation 1, using the obtained value for slope and intercept as α and β constants, respectively. This method is most widely used procedure for cosolvent techniques, such as isocratic RP-HPLC and cosolvent potentiometry procedure (Demiray et al., 2012; Herrero-Martinez et al., 2008; Völgyi et al., 2007; Wiczling et al., 2004; Zapala, 2011).

The ${}^w pK_a$ value could be calculated using the empirical equation (Eq.2 - Eq. 4) presented by Algelov et al. (2008).

$$k_{\text{obs}} = \frac{k_{HA} + k_A \cdot 10^{pH - pK_a}}{1 + 10^{pH - pK_a}} \quad (\text{for acidic compounds}) \quad \text{Eq. 2}$$

$$k_{\text{obs}} = \frac{k_B + k_{BH} \cdot 10^{pK_a - pH}}{1 + 10^{pK_a - pH}} \quad (\text{for basic compounds}) \quad \text{Eq. 3}$$

where k_{HA} , k_B and k_A^- , k_{BH}^+ are the neutral and ionic forms of the compound, respectively.

The conversion formula is as follows:

$$pK_a = pH - \frac{\log(k_{HA} - k)}{k - k_A} \quad \text{Eq. 4}$$

where k_{HA} and k_A^- are k' of the neutral and ionic form of the compound, respectively; the k_{HA} is the minimum value of k' and k_A^- is the maximum one. The pH and k values are defined as points at maximum slope of the pH- k' plot. Huo et al. (2013) applied this method to determine the pK_a value of oxysophocarpine. This procedure was also used to calculate the pK_a value of compound with wide range of lipophilicity using the solely aqueous RP-HPLC approach (Volna et al., 2017).

Another empirically based approach takes into account the type (acetonitrile or methanol) and concentration of the organic modifier. For acetonitrile/water mobile phase the aqueous pK_a could be calculated by the following equations:

$${}^w pK_a = {}^s pK_a + (\% \text{ organic}) * A \quad \text{Eq. 5}$$

$${}^w pK_a = {}^s pK_a + (\% \text{ organic}) * B \quad \text{Eq. 6}$$

The value of A is 0.03 and the value of B is 0.02 and they correspond to acidic and basic pK_a shift per 10% ACN, respectively (Kazakevich & Lohrutto, 2007). This calculation is applicable only for the isocratic RP-HPLC approach.

Conclusion

Several RP-HPLC and potentiometric based approaches are available for the determination of the pK_a value of sparingly soluble active pharmaceutical ingredients.

The isocratic RP-HPLC is simple and easy approach for the pK_a determination of compounds in simple mixtures. Although it is a time-consuming, this approach gives very accurate results. Recently solely aqueous RP-HPLC has emerged as a simple alternative approach. However, more literature data are required to evaluate the reliability of the pK_a data obtained using this approach.

The gradient RP-HPLC concept is more applicable for rough estimation of the *pKa* value. The gradient RP-HPLC/MS methodological approach is useful for a preliminary evaluation of the drug candidates, considering the short analysis time.

The cosolvent potentiometric approach is a well-established procedure for the *pKa* determination of pure, sparingly soluble compounds. The major disadvantage of this concept could be seen in the large number of experiments needed for the determination and in the complexity of calculations for estimation of the aqueous *pKa* value. The surfactant concept appears as a simpler approach and it has a great potential, but additional investigation that include a larger number of compounds is needed.

This overview contributes to the enrichment of the scientific data related to the alternative chromatographic and potentiometric based approaches for determination of sparingly soluble active pharmaceutical ingredients.

References

- Angelov, T., Vlasenko, A., Tashkov, W., 2008. HPLC determination of *pKa* of parabens and investigation on their lipophilicity parameters. *J. Lid. Chromatogr. Relat. Technol.* 31, 188-197. Available at: <https://doi.org/10.1080/10826070701738787>.
- Avdeef, A., 1993a. pH-metric log P. Refinement of partition coefficients and ionization constants of multiprotic substances. *J. Pharm. Sci.*, 82, 183-190. Available at: <https://doi.org/10.1002/jps.2600820214>.
- Avdeef, A., Corner, J.E.A., Thomson, J.S., 1993b. pH-metric log P 3. Glass electrode calibration in methanol-water, applied to *pKa* determination of water-insoluble substances. *Anal. Chem.* 65, 42-49. Available at: <https://doi.org/10.1021/ac00049a010>.
- Avdeef, A., Box, K.J., Comer, J.E.A., Hadley, M., Hibbert, C., 1999. pH-metric log P 11. *pKa* determination of water-insoluble active pharmaceutical ingredients in organic solvent-water mixtures. *J. Pharm. Biomed. Anal.* 20, 631-641. Available at: [https://doi.org/10.1016/s0731-7085\(98\)00235-0](https://doi.org/10.1016/s0731-7085(98)00235-0).
- Avdeef, A., 2001. Physicochemical profiling (solubility, permeability and charge state). *Curr. Top. Med. Chem.* 1, 277-351. Available at: <https://doi.org/10.2174/1568026013395100>.
- Aldeef, A., 2003. *Absorption and Drug Development*. Wiley-Interscience: Hoboken, New Jersey.
- Babic, A., Horvat, A., Pavlovic, M.D., Kastelan-Macan, M., 2007. Determination of *pKa* values of active ingredients. *Trends Anal. Chem.* 26, 1043-1061. Available at: <https://doi.org/10.1016/j.trac.2007.09.004>.
- Demiray, E.Ç., Alsancak, G., Ozkan, A.S., 2009. Determination of *pKa* values of nonsteroidal antiinflammatory drug-oxicams by RP-HPLC and their analyses in pharmaceutical dosage forms. *J. Sep. Sci.* 32, 2928-2936. Available at: <https://doi.org/10.1002/jssc.200900234>.
- Demiray, E.Ç., Koç, D., Daldal, Y.D., Çakir, C., 2012. Determination of chromatographic and spectrophotometric dissociation constant of some beta lactam antibiotics. *J. Pharm. Biomed. Anal.* 71, 139-143. Available at: <https://doi.org/10.1016/j.jpba.2012.06.023>.
- Erdemgil, F.Z., Şanlı, S., Şanlı, N., Özkan, G., Barbosa, J., Guiteras, J., Beltrán, J.L., 2007. Determination of *pKa* values of some hydroxylated benzoic acids in methanol-water binary mixtures by LC methodology and potentiometry. *Talanta* 72, 489-496. Available at: <https://doi.org/10.1016/j.talanta.2006.11.007>.
- Fallavena, P.R., Schapoval, E.E., 1997. *pKa* determination of nimesulide in methanol-water mixtures by potentiometric titrations. *Int. J. Pharm.* 158, 109-112. Available at: [https://doi.org/10.1016/S0378-5173\(97\)00221-4](https://doi.org/10.1016/S0378-5173(97)00221-4).
- Gumustas, M., Şanlı, S., Şanlı, N., Ozkan, S., 2010. Determination of *pKa* values of some antihypertensive active pharmaceutical ingredients by liquid chromatography and simultaneous assay of lercanidipine and enalapril in their binary mixtures. *Talanta* 82, 1528-1537. Available at: <https://doi.org/10.1016/j.talanta.2010.07.037>.
- Herrero-Martinez, J.M., Repolles, C., Bosch, E., Roses, M., Rafols, C., 2008. Potentiometric determination of aqueous dissociation constant of flavonols sparingly soluble in water. *Talanta* 74, 1008-1013. Available at: <https://doi.org/10.1016/j.talanta.2007.08.007>.
- Horvat, C., Melander, W., Molnar, I., 1977. Liquid chromatography of ionogenic substances with nonpolar stationary phases. *Anal. Chem.* 49, 142-154. Available at: <https://doi.org/10.1021/ac50009a044>.
- Huo, H., Li, T., Zhang, L., 2013. *pKa* determination of oxysophocarpine by reversed-phase high performance liquid chromatography. *SpringerPlus* 2, 270. Available at: <https://doi.org/10.1186/2193-1801-2-270>.
- Jaiswal, P.V., Ijeri, V.S., Srivastava, A.K., 2005. Effect of surfactants on the dissociation constant of ascorbic and maleic acids. *Colloids Surf B Biointerfaces* 46, 45-51. Available at: <https://doi.org/10.1016/j.colsurfb.2005.09.001>.
- Kaliszan, R., Wiczling, P., 2010. Gradient reversed-phased high-performance chromatography of ionogenic analytes. *Trends Anal. Chem.* 30, 1372-1381. Available at: <https://doi.org/10.1021/ac034999v>.
- Kazakevich, Y., Lobrutto, R., 2007. *HPLC for pharmaceutical scientists*, fourth Ed. John Wiley & Sons, Inc.: Hoboken, New Jersey.
- Kerns, E.H., Di, L., 2004. Physicochemical profiling: overview of the screens. *Drug Discov. Today Technol.* 1, 343-348. Available at: <https://doi.org/10.1016/j.ddtec.2004.08.011>.
- Kraft, A., 2003. The determination of the *pKa* of multiprotic, weak acids by analyzing potentiometric acid-base titration data with difference plots. *J. Chem. Ed.* 80, 554-559. Available at: <https://doi.org/10.1021/ed080p554>.
- Kubik, L., Struck-Lewicka, W., Kaliszan, R., Wiczling, P., 2015. Simultaneous determination of hydrophobicity and dissociation constant for a large set of compounds by gradient reverse phase high performance liquid chromatography-mass spectrometry technique. *J. Chromatogr. A* 1416, 31-37. Available at: <https://doi.org/10.1016/j.chroma.2015.09.003>.
- Manallack, D.T., 2007. The *pKa* Distribution of Active pharmaceutical ingredients: Application to Drug Discovery. *Perspec. Medicin. Chem.* 1, 25-38.
- Manderscheid, M., Eichinger, T., 2003. Determination of *pKa* values by Liquid Chromatography. *J. Chrom. Sci.* 41, 323-

326. Available at:
<https://doi.org/10.1093/chromsci/41.6.323>.
- Narasimham, L., Barhate, V.D., 2011. Physico-chemical characterization of some beta blockers and anti-diabetic active pharmaceutical ingredients - potentiometric and spectrophotometric pKa determination in different co-solvents. *Eur. J. Chem.* 2, 36-46. Available at: <https://doi.org/10.5155/eurjchem.2.1.36-46.371>.
- Némecová, I., Rychlovský P., Pudilová, D., 2009. Effect of surfactants and ionic strength on dissociation constant of hydrochlorides of phenothiazine derivatives. *Anal. Lett.* 42, 646-658. Available at:
<https://doi.org/10.1080/00032710902722046>.
- Pandey, M.M., Jaipal, A., Kumar, A., Malik, R., Charde, S.Y., 2013. Determination of pKa of felodipine using UV-Vis spectroscopy. *Spectrochim. Acta A Mol. Biomol. Spectrosc.* 115, 887- 890. Available at:
<https://doi.org/10.1016/j.saa.2013.07.001>.
- Rafols, C., Roses, M., Bosch, E., 1997. Dissociation constant of several non-steroidal anti-inflammatory active pharmaceutical ingredients in isopropyl alcohol/water mixtures. *Anal. Chim. Acta* 350, 249-255. Available at: [https://doi.org/10.1016/S0003-2670\(97\)00307-3](https://doi.org/10.1016/S0003-2670(97)00307-3).
- Ravichandiran, V., Devarajan, V., Masilamani, K., 2011. Determination of ionization constant (pKa) for poorly soluble active pharmaceutical ingredients by using surfactants: a novel approach. *Pharm. Lett.* 3, 183-192.
- Reijenga, A., Van Hoof, A., Van Loon, A., Teunissen, B., 2013. Development of methods for determination of pKa values. *Anal. Chem. Insights* 8, 53-71. Available at: <https://doi.org/10.4137/ACIS12304>.
- Şanlı, N., Şanlı, S., Ozkan, G., Denizli, A., 2010. Determination of pKa of some sulfonamides by LC and LC-PDA methods in acetonitrile-water binary mixtures. *J. Braz. Chem. Soc.* 21, 1952-1960. Available at: <http://dx.doi.org/10.1590/S0103-50532010001000022>.
- Sheldovsky, T., 1962. in Peasce, B. (Eds), *Electrolytes*. Pergamon Press, New York.
- Špirtović-Halilović, S., Završnik, D., 2010. Computer programs for calculating pKa: a comparative study for 3-(3-(2-nitrophenyl) prop-2-enoyl)-2H-1-benzopyran-2-one. *J. Serb. Chem. Soc.* 75, 243-248. Available at: <https://doi.org/10.2298/JSC1002243S>.
- Takacs-Novak, K., Box, K.J., Avdeef, A., 1997. Potentiometric pKa determination of water-insoluble compounds: validation study in methanol/water mixtures. *Int. J. Pharm.* 151, 235-248. Available at: [https://doi.org/10.1016/S0378-5173\(97\)04907-7](https://doi.org/10.1016/S0378-5173(97)04907-7).
- Van de Venne, J.L.M., Hendrix J.L.H.M., Deeler, R.S., 1978. Retention behavior of carboxylic acids in reverse-phase column liquid chromatography. *J. Chromatogr.* 167, 1-16. Available at:
[https://doi.org/10.1016/S0021-9673\(00\)91142-7](https://doi.org/10.1016/S0021-9673(00)91142-7).
- Völgyi, G., Ruiz, R., Box, K., Comer, J., Bosch, E., Takacs-Novák, K., 2007. Potentiometric and spectrophotometric pKa determination of water-insoluble compounds: Validation study in a new cosolvent system. *Anal. Chim. Acta*, 583, 418-428. Available at:
<https://doi.org/10.1016/j.aca.2006.10.015>.
- Volna, T., Motyka, K., Hlavác, J., 2017. RP-HPLC determination of dissociation constant using solely aqueous mobile phase. *J. Pharm. Biomed. Anal.* 134, 143-148. Available at:
<https://doi.org/10.1016/j.jpba.2016.11.038>.
- Uhrova, M., Miksik, I., Deyl, Z., Bellini, S., 1997. Determination of dissociation constants by separation methods (HPLC and CE). Theoretical background and guidelines for application. *Process Contr. Qual.* 10, 151-167.
- Yuanqin, Z., Fan, L., Xiaoyan, L., Jing, L., 2005. The effect of surfactant micelles on the dissociation constant and transition points and transition intervals of acid-base indicators. *Talanta* 56, 705-710. Available at: [https://doi.org/10.1016/s0039-9140\(01\)00618-x](https://doi.org/10.1016/s0039-9140(01)00618-x).
- Wiczling, P., Markuszewski, M.J., Kaliszan, R., 2004. Determination of pKa by pH gradient reversed-phase HPLC. *Anal. Chem.* 76, 3069-3077. Available at: <https://doi.org/10.1021/ac049807q>.
- Wiczling, P., Kawczak, P., Nasal, A., Kaliszan, R., 2006. Simultaneous determination of pKa and lipophilicity by gradient RP HPLC. *Anal. Chem.* 78, 239-249. Available at: <https://doi.org/10.1021/ac0512103>.
- Wiczling, P., Markuszewski, M.W., Markuszewski, M.J., Kaliszan, R., 2008. The application of gradient reverse-phase liquid chromatography to the pKa and log kw determination of polyprotic analytes. *J. Chromatogr. A* 1214, 109-114.
- Wiczling, P., Struck-Lewicka, W., Kubik, L., Siluk, D., Markuszewski, M.J., Kaliszan, R., 2014. The simultaneous determination of hydrophobicity and dissociation constant by liquid chromatography-mass spectrometry. *J. Pharm. Biomed. Anal.* 94, 180-187. Available at: <https://doi.org/10.1016/j.jpba.2014.01.038>.
- Zapala, L., 2011. Potentiometric studies on the equilibria of flufenamic acid in aqueous solutions and in two-phase organic solvent+water systems. *J. Solution Chem.* 40, 198-210. Available at: <https://doi.org/10.1007/s10953-010-9642-6>.

Резиме

Преглед на хроматографски и потенциометриски методи за определување на pK_a вредност на слабо растворливи супстанции

Наталија Наков^{1*}, Јелена Ацевска¹, Катерина Брезовска¹, Анета Димитровска¹

¹Институт за применета хемија и фармацевтски анализи, Фармацевтски Факултет,
Универзитет „Св. Кирил и Методиј“, 1000 Скопје, Македонија

Клучни зборови: константа на дисоцијација, pK_a , слабо растворливи супстанции, определување, реверзно-фазна течна хроматографија, потенциометрија, сурфактанти

Константата на дисоцијација (pK_a), како една од најважните физичко-хемиски својства на активните супстанции, има големо значење за фармацевтска индустрија од повеќе аспекти. Повеќето активни супстанции се нерастворливи во вода или немаат хромофор, поради што не се применливи конвенцијалните техники (UV/Vis спектрометрија и потенциометрија) за определување на константата на дисоцијација. Во овој ревијален труд даден е преглед на новите пристапи што се користат за определување на pK_a вредноста на слабо растворливите супстанции, со посебен осврт на реверзно-фазната течна хроматографија и на потенциометриски титрации во присуство на корастворувач/сурфактант. Дополнително, даден е осврт на различни пристапи што се применуваат за пресметување на константата на дисоцијација.