Use of Different Agro-wastes in the Cultivation of *Pleurotus ostreatus* (Jacq.) Kummer

Mary O. Iwuagwu

Abia State University, Uturu, Nigeria, mary.iwuagwu@abiastateuniversity.edu.ng

Deborah S. Nwaukwa

Abia State University, Uturu, Nigeria, debby4dance@gmail.com

Chidi E. Nwaru

Abia State University, Uturu, Nigeria, nwaru.chidi@abiastateuniversity.edu.ng

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>

Part of the [Agriculture Commons](#), and the [Botany Commons](#)

Recommended Citation

Iwuagwu, M. O., Nwaukwa, D. S., & Nwaru, C. E. (2020). Use of Different Agro-wastes in the Cultivation of *Pleurotus ostreatus* (Jacq.) Kummer, *Journal of Bioresource Management*, 7 (2).

DOI: <https://doi.org/10.35691/JBM.0202.0129>

ISSN: 2309-3854 online

(Received: Apr 17, 2020; Accepted: May 15, 2020; Published: Jun 1, 2020)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in *Journal of Bioresource Management* by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

Use of Different Agro-wastes in the Cultivation of *Pleurotus ostreatus* (Jacq.) Kummer

Cover Page Footnote

The authors are grateful to Prof. I.C. Okwulehie for his invaluable assistance

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

USE OF DIFFERENT AGRO-WASTES FOR CULTIVATION OF *PLEUROTUS OSTREATUS* (Jacq.) Kummer

MARY O. IWUAGWU, DEBORAH S. NWAUKWA AND CHIDI E. NWARU

Abia State University, Uturu, Nigeria

*corresponding author: mary.iwuagwu@abiastateuniversity.edu.ng

ABSTRACT

Agro-wastes are abundant in both rural and urban areas and when carelessly disposed off to the environment by dumping or burning, they lead to environmental pollution, and consequently to health hazards. Growing of edible mushroom on these wastes can contribute to decrease in environment pollution. This work investigated the growth and yield of *Pleurotus ostreatus* on different agro-wastes: plantain midrib (PM), corn straw (CS), sugarcane bagasse (SB) and their combinations (PM+CS, PM+SB, CS+SB, CS+PM+SB) with a view to finding the best substrate for the growth of *P. ostreatus*. The experiment was carried out using completely randomized design with seven treatments and three replications. Results revealed that mycelia growth period and days to pinhead formation were faster in corn straw (7 and 11 days, respectively from the day of inoculation), this was followed by CS+SB (8 and 12 days, respectively) while CS+PM+SB recorded the longest days (13 and 17, respectively). Stipe length ranged from 2.6 cm in PM to 3.3 cm in CS+PM while the pileus diameter was between 3.3 cm in CS and PM and 4.3 cm in PM+SB. Number of fruiting bodies, fresh and dry weights of the mushroom were significantly higher in CS than in the other substrates. The least number of fruiting bodies, fresh and dry weights were recorded in CS+PM+SB.

Keywords: Agro-wastes, mycelium, growth period, *Pleurotus ostreatus*, pollution.

INTRODUCTION

The genus, *Pleurotus* is commonly known as Oyster mushroom. It belongs to the class, Basidiomycetes and a group of fungi known as white-rot fungi as they produce a white mycelium (Tsujiyama and Ueno, 2013). These are edible mushrooms, highly nutritious with high contents of protein, fiber, vitamins and minerals (Mattila et al., 2001; Shin et al., 2007). They can be effectively used as appetizers and also as an ingredient in soups, sauces, salads, and meat dishes (Mattila et al., 2001). *Pleurotus* sp. is reported to be suitable for inclusion into calorie-controlled diet because of its characteristic high water content and low calorific value (1510 kJ-

edible parts) (Jaworska and Bernas, 2009). They have several beneficial properties (anti-fungal, anti-oxidant, etc.) as they contain important bioactive molecules such as phenolic compound, terpenes, steroids and polysaccharides (Yang et al., 2013; Zhang et al., 2016).

There are over 2000 species of mushrooms that exist in nature (Manzi et al., 2001) but only about 35 species have been commercially cultivated (Aida et al., 2009). *Pleurotus* spp. are cultivated worldwide and rank second in the world mushroom market after *Lentinus edodes* (Royse et al., 2017; Bellettini et al., 2019). *Pleurotus* spp. are one of the most extensively studied white-rot fungi because of their ability to degrade cellulose, hemi-cellulose and lignin (Machado et al., 2016). Hence, they can be

cultivated on various lignocellulosic substrates (Savoie et al., 2007; Li and Shah, 2016). These substrates are majorly agro-industrial wastes, which are produced in enormous quantities causing disposal problems and consequently environmental pollution/health risks (Garg and Gupta, 2009). Utilization of these wastes in mushroom cultivation can help reduce these problems.

Many different lignocellulosic materials have been tested for their suitability in the cultivation of *Pleurotus* spp. including grass straws (Okwulehie and Okwujiaku, 2008), sawdust, cassava peels, corn cobs, melon chaff, paper waste and water hyacinth (Ambi et al., 2011), soybean straw, wheat straw, paddy straw, sugarcane bagasse, sunflower stalks, maize stalks, domestic waste, used tea leaves, fruit waste, newspaper, bamboo leaves, saw dust (Dehariya and Vgas, 2013), corn cobs (Itelima, 2012), cotton waste (Shah et al., 2004), palm leaves (Alananbeh et al., 2014), maize stover and thatch grass (Fanadzo et al., 2010) among others.

There is an ever-increasing demand for mushrooms with the total demand for all the species estimated at 60.0 million tons per annum and the world output at about 34 million tons (Panjikkaran and Mathew, 2012). China contributes about half of the world's mushroom production. Whereas, South Africa (the only African country estimated) contributes less than 0.3 % (FAO, 2009). There is therefore no gainsaying that the production level in Nigeria and the whole of Africa is insignificant. Research efforts directed in the area of exploring more substrates and determining the best for the cultivation of mushroom more especially *Pleurotus* spp., that is widely adapted, will no doubt enhance its production.

The aim of this study was to compare the growth and yield responses of *P. ostreatus* cultivated on different agro-

wastes, namely corn straw, plantain midrib, sugarcane bagasse and their combinations with a view to determine the best substrate for the cultivation of *P. ostreatus*.

MATERIALS AND METHODS

Study Location

The study was conducted at the mushroom unit of Michael Okpara University of Agriculture, Umudike, in Southeastern Nigeria.

Sources of Materials

One bottle of *Pleurotus ostreatus* mother spawn was obtained from the mushroom house of Plant Science and Biotechnology Department, Michael Okpara University of Agriculture Umudike, Abia State. Red sorghum grain (*Sorghum bicolor*) was purchased from Umuahia main market, Abia State, Nigeria and was utilized as substrate for spawn multiplication. Plantain midrib, sugarcane bagasse and corn stalk were obtained from the agricultural farm of Michael Okpara University of Agriculture, Umudike, Abia State.

Spawn Preparation

The method described by Stamets (2000) was adopted in the multiplication of spawn. Substrate preparation was carried out according to the method described by Sharma (2003).

Inoculation of the Substrates

After cooling, equal amount (30 g) of *Pleurotus ostreatus* grain spawn was inoculated into each of the buckets containing the substrate, by gently placing it on layers of the substrate. The inoculated buckets were covered and transferred onto wooden racks in the cropping room. The buckets were covered with a black polythene

bag to serve as dark incubation for ramification to take place. The cropping room was well-lit, ventilated and constantly flooded with clean water to maintain the environmental conditions (good aeration, temperature between 18-32 °C, high humidity) needed for the vegetative growth of the mushroom.

Induction of Fruiting Bodies

In order to induce fruiting body formation in the substrates, the polythene bag was removed on the 10th day, when it was assumed that the mycelia have fully colonized the substrate (Plate 1). The bucket covers were removed to increase aeration and flooding of the floors was reduced so as to reduce the humidity of the room. The buckets were routinely observed until immature mushrooms started to emerge as primordia.

Experimental Design and Treatments

The experiment was laid in a completely randomized design. Each treatment was replicated thrice. The treatments were: plantain midrib (PM), corn straw (CS), sugarcane bagasse (SB), 500 g plantain midrib + 500 g corn straw (PM + CS), 500 g plantain midrib + 500 g sugarcane bagasse (PM + SB), 500 g corn straw + 500 g sugarcane bagasse (CS + SB) and 400 g corn straw + 300 g plantain midrib + 300 g sugarcane bagasse (CS + PM + SB).

Plate 1: Substrates fully colonized by *P. ostreatus* mycelium

Assessment Parameters

i. Vegetative Growth

The mycelium growth period was recorded as the time (in days) taken from inoculation of the spawn to the time the spawn fully ramified and colonized the substrate. While days to pinhead formation were recorded as the number of days taken from the time (days) the spawn fully colonized the substrate to the time fruiting bodies appeared on the substrates.

ii. Yield and Yield Attributes

At each flush (three flushes), the fruiting bodies produced (Plate 2) from each substrate were harvested by hand-picking. The number of fruiting bodies in different substrates was counted and total of the three flushes was recorded. The diameter of the cap was measured in centimeter (cm) with a transparent plastic ruler from one edge to the other edge when it had fully expanded. The length of the stripe was also measured (cm) using a transparent plastic ruler. The fresh weight of the fruiting-bodies was determined by weighing them immediately after harvesting, using an electronic

weighing balance. The dry weight was taken after drying them in an oven at 60 °C.

Statistical Analysis

Data collected was analyzed statistically using ANOVA. Means that were found significant were separated using Fisher's Least Significant Difference (LSD) at a significance level of $P < 0.05$.

RESULTS

The results for period of mycelium growth, days to pinhead formation, stipe length and pileus diameter are presented in Table 1. There was a significant ($P < 0.05$) difference in the period of mycelium growth, which is the number of days taken by the fungus to fully colonize the substrate.

Table 1: Effect of substrates on period of mycelium growth (PMG), days to pinhead formation (DPH), stipe length (SL) and pileus diameter (PD)

Treatment	PMG (days)	DPH (days)	SL (cm)	PD (cm)
PM	16	10	2.6	3.3
CS	11	7	3.1	3.3
SB	-	-	-	-
PM+ CS	16	12	3.3	3.5
PM+SB	16	12	2.8	4.5
CS+SB	12	8	2.8	3.4
CS+PM+SB	17	13	2.7	3.4
LSD	4.0	2.2	0.5	1.0

PM = plantain midrib; *CS* = corn straw; *SB* = sugarcane bagasse; *PM+CS* = plantain midrib + corn straw; *PM+SB* = plantain midrib + sugarcane bagasse; *CS+SB* = corn straw + sugarcane bagasse; *CS+PM+SB* = corn straw + plantain midrib + sugarcane bagasse

Figure 1: Effect of substrates on the number of fruiting bodies on *P. ostreatus*.

PM = plantain midrib; *CS* = corn straw; *SB* = sugarcane bagasse; *PM+CS* = plantain midrib + corn straw; *PM+SB* = plantain midrib + sugarcane bagasse; *CS+SB* = corn straw + sugarcane bagasse; *CS+PM+SB* = corn straw + plantain midrib + sugarcane bagasse

Statistical analysis revealed significant effect of substrate formulation on time (days) taken for pinhead formation. This ranged from 7 days in corn straw to 13 days in the three combinations (*CS* + *PM* + *SB*). It took significantly shorter time for the pinheads to appear on corn straw alone in contrast to the other substrates except in the combination of corn straw and sugarcane bagasse. The number of days taken for the pinheads to appear on the other substrates did not differ significantly ($P < 0.05$).

The stipe length of the fruiting bodies of *P. ostreatus* was significantly affected by the substrate formulation. Fruiting bodies harvested from plantain midrib plus corn straw had the highest stipe length that was significantly ($P < 0.05$) different from the stipe lengths of those harvested from the three combinations *CS+PM+SB* and sole plantain midrib, which recorded the least. Highest pileus diameter was recorded in fruiting bodies harvested from combination of plantain midrib plus sugarcane bagasse.

Plate 2: *P. ostreatus* fruiting bodies growing in different substrates contained in a bucket.

The number of fruiting bodies produced was significantly highest in corn straw substrate (Figure 1). The least number of fruiting bodies was produced by the combination of all three substrates.

The highest fresh weight was obtained in sole corn straw substrate with a mean value of 193.5 g/kg of substrate and this was significantly ($P < 0.05$) different from the others. The least fresh weight (49.5 g/kg) was recorded in the all-three substrates combination (*CS+PM+SB*), though it was not significantly different from that of

plantain midrib mixed with sugarcane bagasse (67.5 g/kg) (Figure 2).

Similarly, the highest mushroom dry weight of 14.7 g/kg, was obtained in sole corn straw substrate. This value was

significantly different from those obtained from the other treatments (Figure 3). The least dry weight of 4.2 g/kg was obtained in plantain midrib mixed with sugarcane bagasse.

Figure 2: Effect of substrates on fresh weight of *P. ostreatus*.

PM = plantain midrib; CS = corn straw; SB = sugarcane bagasse; PM+CS = plantain midrib + corn straw; PM+SB = plantain midrib + sugarcane bagasse; CS+SB = corn straw + sugarcane bagasse; CS+PM+SB = corn straw + plantain midrib + sugarcane bagasse

Figure 3: Effect of substrates on dry weight of *P. ostreatus*.

PM = plantain midrib; CS = corn straw; SB = sugarcane bagasse; PM+CS = plantain midrib + corn straw; PM+SB = plantain midrib + sugarcane bagasse; CS+SB = corn straw + sugarcane bagasse; CS+PM+SB = corn straw + plantain midrib + sugarcane bagasse

DISCUSSION

The most important phases in the cultivation of mushrooms are spawn running and pinhead formation. Spawn running or period of mycelium growth refers to the period of mycelia growing on the substrate from inoculation to full ramification. The period of mycelia growth in this study ranged from 11 days in CS to 17 days in CS+PM+SB. This range was shorter than the ranges of 14-21 days and 17-23 days reported by Tan (1981) and Shah et al. (2004), respectively. The observable difference among the treatments in period of mycelia growth may be due to the composition of the substrate. Chang and Miles (2004) reported that the variation in the number of days taken for the spawn to complete colonization of a given substrate is a function of the fungus strain, growth condition and substrate type. According to Stamets (2000), longer spawn run can be attributed to the compactness of the substrate owing to small particle size of the substrate, which interferes with mycelial growth of the mushroom.

Pin head formation (primordium initiation) was observed following the invasion of the substrates by *P. ostreatus* mycelia. The result of the present study indicated that corn straw (CS) substrate positively influenced the period of appearance of the pin head as the fruiting bodies appeared earlier on corn straw (7 days after full colonization and 18 days from spawning) than the other substrates. This is in agreement with the report of Stanley et al. (2010). Naraian et al. (2009) attributed earlier appearance of pin head in corn straw to its high content of lignin, which upon enzymatic breakdown by the fungus, releases enough nutrients for the fungal growth.

The time required for the formation of pinhead, which was between 18-30 days from spawning was comparable with other

similar studies where pinhead appeared around 20-23 days and 28-34 days on different substrates (Fan et al., 2000; Islam et al., 2009). Shah et al. (2004) found that pinheads appeared in about 24 – 30 days from spawning. Variation in mycelia growth rate, colonization and primordial initiation has also been observed when other *Pleurotus* mushroom species were grown on a wide range of substrates (Islam et al., 2009; Dehariya and Vyas, 2013; Hwang et al., 2015).

The results of the study showed that fructification of *P. ostreatus* occurred in all the substrates (Plates 3 and 4) except on undecomposed sugarcane bagasse. This indicates that these substrates contained nutrients that supported the growth of the mushroom (Okwulehie and Okwujiako, 2008; Wabali and Wocha, 2013). In addition, *Pleurotus* spp. have a high saprophytic ability and can successfully grow on a variety of lignocellulosic substrates (Machado et al., 2016; Iwuagwu et al., 2017).

The stipe height ranged from 2.6 – 3.3 cm. This is in conformity with the range of 2.6 – 3.6 cm earlier reported by Iwuagwu et al. (2017) in saw dust, corn husk and their combination. Dehariya and Vyas (2013) reported that stipe height of *P. sajor-caju* ranged from 2.5 – 2.9 cm in conventional substrates (straws of soybean, wheat, paddy, sunflower and maize) and 2.3 – 3.5 cm in non-convectional substrates (domestic waste, fruit waste, used tea leaves, bamboo leave and sawdust). Islam et al. (2009) observed a stipe height of *P. flabellatus*, an oyster mushroom cultivated on various substrates as between 2.20 and 3.59 cm.

Pileus diameter differed significantly among the substrates and ranged from 3.3 to 4.5 cm. This range was within the range (1.92-4.78 cm) reported by Okwulehie and Okwujiako (2008). Bonatti et al. (2004) had earlier observed that *P. ostreatus* cap ranged

from 2–15 cm in diameter while the stipe was very short. However, the significant difference in the diameter of the pileus could be due to texture and substrate formulation as well as nutrients in the substrate, which possibly affected the composition of the final mushroom growth substrate qualities such as water holding capacity and degree of aeration (Reyes et al., 2009).

CONCLUSION

This work investigated the use of different agro-wastes namely corn straw, plantain midrib, sugarcane bagasse, and their combinations on the cultivation of *P. ostreatus*. Results of this study have shown that *P. ostreatus* can be cultivated successfully on sole corn husk and plantain mid rib. Hence, corn straw may be considered as a good substrate for the cultivation of *P. ostreatus*.

CONFLICT OF INTEREST

Authors declare no conflict of interest.

REFERENCES

- Aida FMNA, Shuhaimi M, Yazid M and Maaruf AG (2009). Mushroom as a potential source of prebiotics: a review. *Trends Food Sci Technol.*, 20: 567–575.
- Ambi AA, Pateh UU, Sule MI and Nuru GF (2011). Studies on cultivation of some wild edible mushroom on some lignocellulosic substrates in main Campus, Ahmadu Bello University, Zaria. *Int J Pharm Res Inno.*, 3: 14-18.
- Alananbeh KM, Bouqellah NA and Al Kaff NS (2014). Cultivation of oyster mushroom *Pleurotus ostreatus* on date-palm leaves mixed with other agro-wastes in Saudi Arabia. *Saudi J Biol Sci.*, 21: 616–625.
- Bellettini MB, Fiorda FA, Maieves HA, Teixeira GL, Avila S, Hornung PS, Junior AM and Ribani RH (2019). Factors affecting mushroom *Pleurotus* spp. *Saudi J Biol Sci.*, 26: 633–646.
- Bonatti M, Karnopp P, Soares HM and Furlan SA (2004). Evaluation of *Pleurotus* species nutritional characteristics when cultivated in different lignocellulose wastes. *Food Chem.*, 88: 425-428.
- Chang ST and Miles PG (2004). In: *Mushroom Cultivation, Nutritional Value, Medicinal Effect and Environmental Impact*. 2nd eds. CRC Press, Boca Raton.
- Dehariya P and Vyas D (2013). Effect of different agro-waste substrates and their combinations on the yield and biological efficiency of *Pleurotus Sajo-caju*. *IOSR J Pharm Biol Sci.*, 8 (3): 60-64.
- Fan L, Pandey A, Mohan R and Socco CR (2000). Use of various coffee industry residues for the cultivation of *Pleurotus ostreatus* in solid state fermentation. *Acta Biotechnol.*, 20 (1): 41-52
- Fanadzo M, Zireva DT, Dube E and Mashingaidze AB (2010). Evaluation of various substrates and supplements for biological efficiency of *Pleurotus sajor-caju* and *Pleurotus ostreatus*. *Afr J Biotechnol.*, 9 (19): 2756–2761.
- FAO (2009). *World mushroom and truffles: Production, 1961-2007*; United Nations, FAO, FAOStat (8/28/2009)
- Garg VK and Gupta R (2009). Vermicomposting of agro-industrial processing waste. In: *Biotechnology for Agro-Industrial Residues Utilization*. Springer, Netherlands: pp. 431–456.

- Hwang S, Li Y and Lin H (2015). The use of sawdust mixed with ground branches pruned from wax apple or indian jujube as substrate for cultivation of king oyster mushroom (*Pleurotus eryngii*). *Hortsci.*, 50 (8): 1230–1233.
- Islam MZ, Rahman H and Hafiz F (2009). Cultivation of oyster Mushroom (*Pleurotus flabellatus*) on different substrates. *Int J Sustain Crop Prod.*, 4: 45-48.
- Itelima JU (2012). Cultivation of mushroom (*Pleurotus ostreatus*) using corn cobs and saw dust as the major substrates. *Global J Agric Sci.*, 11 (1): 51-56.
- Iwuagwu MO, Levi O and Ndukaji-Clement C (2017). Evaluation of different agro-wastes for the cultivation of oyster mushroom (*Pleurotus ostreatus* (Jacq) P. Kumn). Pp 138 – 142. In: Okoronkwo, N.E., Ogbonnaya, C. I. (eds.). *Fungi: A Key to Food Security*. Proceedings of the 9th Annual Conference of Mycological Society of Nigeria, held at ABSU, Nigeria; December 10 – 13, 2017.
- Jaworska G and Bernas E (2009). Qualitative changes in *Pleurotus ostreatus* (Jacq.: Fr.) Kumm mushrooms resulting from different methods of preliminary processing and periods of frozen storage. *J Sci Food Agric.*, 89: 1066–1075.
- Li S and Shah NP (2016). Characterization, antioxidative and bifidogenic effects of polysaccharides from *Pleurotus eryngii* after heat treatments. *Food Chem.*, 197: 240–249.
- Machado ARG, Teixeira MFS, Kirsch LS, Campelo MCL and Oliveira IMA (2016). Nutritional value and proteases of *Lentinus citrinus* produced by solid state fermentation of lignocellulosic waste from tropical region. *Saudi J Biol Sci.*, 23 (5): 621–627.
- Manzi P, Aguzzi A and Pizzoferrato L (2001). Nutritional value of mushroom widely consumed in Italy. *Food Chem.*, 73: 321-325.
- Mattila P, Konko K, Eurola M, Pihlawa JM, Astola J, Lietaniemi V, Kumpulainen J, Valtonen M and Piironen V (2001). Content of vitamins, minerals elements and some phenolic compounds in cultivated mushrooms. *J Agric Food Chem.*, 49: 2343-2348.
- Naraian R, Sahu RK, Kumar S, Garg SK, Singh CS and Kanaujia RS (2009). Influence of different nitrogen rich supplements during cultivation of *Pleurotus florida* on corn cob substrate. *Environmentalist*, 29: 1-7.
- Okwulehie IC and Okwujiako IA (2008). The use of local Nigerian substrate for the production of *Pleurotus ostreatus* var. *florida* (Eger) sporophores. *Dynamic Biochem Process Biotechnol Mol Biol.*, 2: 38-40.
- Panjikkaran ST and Mathew D (2013). An environmentally friendly and cost-effective technique for the commercial cultivation of oyster mushroom [*Pleurotus florida* (Mont.) Singer]. *J Sci Food Agric.*, 93: 973-976.
- Reyes RG, Lopez LL, Kumakura K, Kalaw SP, Kumakura TE and Eguchi F (2009). *Corpinus comatus*, a newly domesticated wild nutraceutical mushroom in the Philippines. *J Agric Technol.*, 5 (2): 299-316.
- Royse DJ, Baars JJP and Tan Q (2017). Current overview of mushroom production in the world. In: Diego CZ, Pardo-Giménez A (eds.). *Edible and Medicinal Mushrooms: Technology and Applications*. Wiley-Blackwell, USA: pp 592.

- Savoie JM, Salmones D and Mata G (2007). Hydrogen peroxide concentration measured in cultivation substrates during growth and fruiting of the mushrooms. *Agaricus bisporus* and *Pleurotus* spp. *J Sci Food Agric.*, 87: 1337–1344.
- Shah ZA, Ashraf M and Ishitiaq M (2004). Comparative study on cultivation and yield performance of oyster mushroom (*Pleurotus ostreatus*) on different substrates (wheat straw, leaves, sawdust). *J Nutr.*, 3: 158-160.
- Sharma BB (2003). Effects of different substrates on spawn growth and yield of pink oyster mushroom *Pleurotus djamor* (Fri.) Boedijn. *J Mycol Plant Pathol.*, 3 (2): 265-268.
- Shin CK, Yee CF, Shya LJ and Atong M (2007). Nutritional properties of some edible mushrooms in Sabah. *J Appl Sci.*, 7 (15): 2216-2221.
- Stamets P (2000). *Growing Gourmet and Medicinal Mushrooms*. 3rd ed. Ten Speed Press, Berkley (CA): pp. 574.
- Stanley HO (2010). Effect of substrate on spawn production and mycelial growth of oyster mushroom species. *Agric Biol J North Ame.*, 1 (5): 817-820.
- Tan KK (1981). Cotton waste is a good substrate for cultivation of *Pleurotus ostreatus* the oyster mushroom. *Mushroom Sci.*, 11: 705-710.
- Tsujiyama S and Ueno H (2013). Performance of wood-rotting fungi-based enzymes on enzymic saccharification of rice straw. *J Sci Food Agric.*, 93: 2841–2848.
- Wabali CV and Wocha I (2013). Effect of Nutrient concentration on the yield of mushroom (*Pleurotus ostreatus*). *Greener J Agric Sci.*, 3 (6): 437-444.
- Yang W, Guo F and Wan Z (2013). Yield and size of oyster mushroom grown on rice/wheat straw basal substrate supplemented with cotton seed hull. *Saudi J Biol Sci.*, 20: 333–338.
- Zhang JJ, Li Y, Zhou T, Xu DP, Zhang P, Li S and Li HB (2016). Bioactivities and health benefits of mushrooms mainly from China. *Molecules*, 21 (7): 938.