Feeding Ecology, Behaviour and Habitat Utilization of Black Drongo (*Dicrurus Macrocerus*) in Pothwar Plateau, Pakistan

Muhammad Bilal

Department of Biology, Virtual University of Pakistan, Lahore, Pakistan, shbilal7707@gmail.com

Zeeshan Khalid

WWF-Pakistan, Ferozepur Road, Lahore, Pakistan, chz39@hotmail.com

Ali Hasnain Mosvi

Department of Wildlife Management, PMAS-Arid Agriculture University, Rawalpindi, Pakistan

Amir Naseer

Department of Wildlife Management, PMAS-Arid Agriculture University, Rawalpindi, Pakistan, aamirnaseer70@yahoo.com

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>

Part of the [Biodiversity Commons](#), [Ornithology Commons](#), and the [Zoology Commons](#)

Recommended Citation

Bilal, M., Khalid, Z., Mosvi, A. H., & Naseer, A. (2020). Feeding Ecology, Behaviour and Habitat Utilization of Black Drongo (*Dicrurus Macrocerus*) in Pothwar Plateau, Pakistan, *Journal of Bioresource Management*, 7 (2).

DOI: <https://doi.org/10.35691/JBM.0202.0131>

ISSN: 2309-3854 online

(Received: May 5, 2020; Accepted: May 8, 2020; Published: Jun 1, 2020)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in Journal of Bioresource Management by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

Feeding Ecology, Behaviour and Habitat Utilization of Black Drongo (*Dicrurus Macrocerus*) in Pothwar Plateau, Pakistan

Cover Page Footnote

ACKNOWLEDGEMENT Authors are highly thankful to all those who assisted in the present study, and provided revision of the manuscripts. Special thanks to Haris Mughal, Noman Ali, Shahzaib Ahmed and Tayyab Shehzad, who assisted in field activities and data collections.

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

FEEDING ECOLOGY, BEHAVIOUR AND HABITAT UTILIZATION OF BLACK DRONGO (*DICRURUS MACROCERCUS*) IN POTHWAR PLATEAU, PAKISTAN

MUHAMMAD BILAL^{1*}, ZEESHAN KHALID², ALI HASNAIN MOSVI³ AND AMIR NASEER³

¹Department of Biology, Virtual University of Pakistan, Lahore, Pakistan

²WWF-Pakistan, Ferozepur Road, Lahore, Pakistan

³Department of Wildlife Management, PMAS-Arid Agriculture University, Rawalpindi, Pakistan

*Corresponding author: shbilal7707@gmail.com

ABSTRACT

Black Drongo is an insectivorous bird native to Pakistan. The present study focused on its feeding preferences, behaviour, ecology and habitat relationships. The study was carried out from first week of December, 2019 to the second week of April, 2020 in the two villages of Rawalpindi named as Daultala (Location-I) and Nata Gujarmall (Location-II). Observations were made by point count method on foot using line transect method. Air (52.27%) was the most commonly used feeding substrate. Aerial feeding mode (52.27%) is preferred over plant or tree gleaning (n=58) and ground feeding (n=47). Most common perching site was man made obstacle, wire (37.72%). Thirteen indigenous tree species and two exotic trees including Eucalyptus (*Eucalyptus camaldulensis*) and Bottle brush (*Callistemon citrinus*) were used as perching site by Black Drongo throughout the study. Black Drongo remained solitary (36.36%) during feeding rituals and shared (63.63%) its intimate habitat with other birds. At Location-I and Location-II, habitat was shared with 12 and 16 bird species, respectively. The mean time of prey capturing attempt was lower at Location-I (1.93 ± 0.17 min) than Location-II (2.16 ± 0.16 min). The time range of prey capturing attempt was (0.1 ± 0.16 min) to (11.5 ± 0.17 min). Black Drongo captured prey successfully by first attempt (59.54%), maximum attempts noted were 6. After preying attempts Black Drongo came back to same perch site (67.27%); which is a representative of resource abundance.

Keywords: Black Drongo, feeding, behaviour, ecology, habitat

INTRODUCTION

Black Drongo (*Dicrurus macrocercus*: family Dicruridae) is an insectivorous bird, found in South Asia and the Indian subcontinent (Grimmett et al., 2009). It is native to Pakistan, present in farms and open grasslands in Sindh and Punjab (Grimmett et al., 2009; BirdLife International, 2020). It is major constituent of agricultural ecosystem and check the build-up of harmful insects of agriculture crops (Kaur and Kler, 2018). Finding a species' resource preference is essential to basic and applied ecology (Charmantier et al., 2008).

Present study sought out to find foraging behaviour, perch site, perch height, inter and intraspecific behaviour, time of recurrent attack to capture prey,

number of attempts for successful capture and second perch site (after attack) of Black Drongo in agriculture landscapes of Pothohar plateau (Punjab, Pakistan).

MATERIALS AND METHODS

Study Site

Present study was carried out at two villages of Pothohar plateau, viz. Daultala ($33^{\circ}11'33''\text{N}$, $73^{\circ}08'25''\text{E}$: scarce vegetation and trees) and Nata Gujarmall ($33^{\circ}10'21''\text{N}$, $73^{\circ}09'34''\text{E}$) (District Rawalpindi: Punjab, Pakistan). Eucalyptus (*Eucalyptus camaldulensis*), pomegranate (*Punica granatum*), sukh chain (*Pongamia pinnata*), sacred fig (*Ficus religiosa*), bottle brush (*Callistemon rigidus*), guava (*Psidium guajava*), dhrek (*Melia azedarach*), beri

(*Ziziphus mauritiana*), keekar (*Acacia nilotica*), phulai (*A. modesta*), Cypress spp., Java plum (*Syzygium cumini*), neem (*Azadirachta indica*), sheesham (*Dalbergia sissoo*), giant milkweed (*Calotropis gigantea*) are important tree species.

Data Collection

Two workers walked (Jhonson, n.d.) on foot from 06:30 to 10:30 a.m. and from 14:30 to 18:30 p.m. (Verner, 1985) twice a week between November 2019 and April 2020 actively searching Black Drongo until observations on foraging behaviour (aerial, plants/ trees or ground), perch site, perch height, time of recurrent attack to capture prey, second perch site and number of attempts for a successful capture were recorded in the field on data sheet. Black Drongo was observed with naked eyes (Ali, 2002) and by using Russian Tecno Sehfeld Military Binoculars (20x56) without disturbing the animal. Perching height of Black Drongo was found using stick method (Hairiah et al., 2001) and measuring tape to find out distance between observer and perch. Photographs were also taken with the help of camera.

Whenever and wherever Black Drongo was spotted, the behaviour of Black Drongo with conspecifics and with other species present in the vicinity (10 m) and aforementioned parameters were recorded (Verner, 1985; Kaur and Kler, 2018). When a Black Drongo was found it was observed for a maximum of 10 minutes. Observations with respect to other birds (either sharing same perch site or feeding substrate) were recorded within 5 m of radius. The data was collected only once per bird to exclude prejudices, one of them was noted. Within a radius of 10 m no other observations were made to make observations independent of each other. Observations were recorded on each sighted bird and feeding habits were identified by swallowing behaviour (Dinsmore, 1973; Grubb, 1976; Kour and

Sahi, 2012). Only dominant vegetation and vegetation used by Black Drongo either directly for perching or indirectly for gleaning was recorded and all others were excluded. To minimize vegetation sampling bias, only the first used vegetation was recorded.

Statistical Analysis

For statistical analysis Past 4.01 was used. Student's t-test was carried out to on the average perching site and average perching height of Black Drongo to find whether there was any significant difference between the perching heights of both locations (Kaur and Kler, 2018) and for the time of recurrent attack. The values obtained are represented in Mean $\pm$ SE. Results were tabulated (Sidra et al., 2013) and charts were made by using Microsoft Excel 2010.

RESULTS AND DISSCUSIONS

A total of 220 observations were recorded, 110 from each location and following results were obtained:

Foraging Behaviour

Black Drongo used 3 main foraging substrates named as air, plant or tree and ground at both locations collectively shown in Figure 1. At Location-I air (n=54) was used more than other two substrates with a total of 49.09% times. It was the most used substrate when at the site of feeding there were disturbances in crops or grasses, which caused insects to fly. After aerial feeding the most used foraging substrate was ground (n=35) that is 38.81%, it may be due to the reduced cultivation of fields and presence of only grasses, which are roosting and feeding sites for insects. At Location-I ground feeding and insect picking from ground was more than plant gleaning, this result contradicted with the results find by Kaur and Kler (2018). The plant used (n=21) was least used foraging mode of feeding (19.09%).

Figure 1: Foraging behaviour of Black Drongo at both locations, showing the total number of observations in comparison with the mode of feeding.

While at Location-II, air (n=61) was used a total of 55.45% time to forage on insects, plant/tree substrate (n=37) was 33.63% of time to catch a prey and ground (n=12) only 10.90%. At location-II the most prevalent feeding substrate was air. Air (52.27%) was the most used substrate for feeding, followed by plants or trees (26.36%) and ground (21.36%) at both

locations. It was observed from the results that mostly insects were preyed on when they were flying. This result conforms with the findings by Kaur and Kler (2018). During the whole study, Black Drongo was not seen perching on plants of wheat (*Triticum aestivum*) and Oat (*Avena sativa*) (dominant crops) which contradicts findings from the previous study by Kaur and Kler (2018).

The higher number of aerial feeding at both locations means more insect consumption (Okosodo et al., 2016; Kaur and Kler, 2018) as shown in Table 1. Black Drongo only foraged from a perch to ground; where it may not be considered as active ground feeding like Common Myna do (Kaur and Kler, 2018). Ground foraging was the least common method of feeding at Location-II. This was due to the more vegetation and prey abundance at Location-II that provides sufficient food in aerial feeding and presumably they did not feed on ground. Thus, prey abundance plays an important role (Park et al., 2008) in determining the foraging mode (Airola and Barrett, 1985).

Table 1: Feeding behaviour of Black Drongo at both locations with inter-site difference.

Feeding Behaviour	Location-I		Location-II		Inter-site Difference
	Numbers	Percentage	Number	Percentage	
Air/ Aerial feeding	54	49.09%	61	55.45%	7
Plants gleaning	21	19.09%	37	33.63%	16
Ground feeding	35	31.81%	12	10.90%	23

Perch Site

The most used perch was electrical wire (37.72%) at Location-I. The results of this investigation are similar to the previous study done by Kaur and Kler (2018). Black Drongo also used roofs and walls (n=10) of un-cemented houses near agricultural fields. The indigenous tree species of Pomegranate Tree (*Punica granatum*) (n=7), Phulai (*Acacia modesta*) (n=6), Gum Arabic Tree (*Acacia nilotica*)

(n=5), Sukh Chain (*Pongamia pinnata*) (n=4), Chinaberry Tree (*Melia azedarach*) (n=4), Indian Plum Tree (*Ziziphus mauritiana*) (n=3), Common Guava (*Psidium guajava*) (n=3), Cypress spp. (n=2) and Java Plum (*Syzygium cumini*) (n=1) were used by Black Drongo to perch on. Near human dwellings it was found to be perched dominantly on wire and some accidental perch on Dhrek and Java Plum. Eucalyptus (*Eucalyptus camaldulensis*) (n=17) and Bottle brush

(*Callistemon citrinus*) (n=5) were the two exogenous species selected by Black Drongo for perching.

Figure 2: A Black Drongo perching on a wire.

Due to more tree species present at Location-II, Black Drongo used a variety of perching sites, a total of 13 indigenous tree species namely, Gum Arabic Tree (*Acacia nilotica*) (n=12), Indian Plum tree (*Ziziphus mauritiana*) (n=9), Chinaberry

Tree (*Melia azedarach*) (n=7), Indian Lilac (*Azadirachta indica*) (n=4), Indian Rosewood (*Dalbergia sissoo*) (n=3), Pomegranate Tree (*Punica granatum*) (n=2), Sukh Chain (*Pongamia pinnata*) (n=2), Mango Tree (*Mangifera indica*) (n=2), Sacred fig (*Ficus religiosa*) (n=2), Guava (*Psidium guajava*) (n=1), Cypress spp. (n=1), Phulai (*Acacia modesta*) (n=6) were used. While 2 species Eucalyptus (*Eucalyptus camaldulensis*) (n=11), Bottle brush (*Callistemon citrinus*) (n=7), were the exotic species used (Table 1). Most widely used perch was wire (n=40). Roofs (n=6) were also used but less than at Location-I due to the presence of more natural perch sites. It was also seen that Black Drongo utilized a greater number of tree species at Location-II. It may be due to undisturbed habitats and less cultivated fields there. The least used trees for perching were Guava, Cypress spp. and Phulai (n=1 for each). Similar results were found by previous studies on vegetation preferences by insectivorous birds (Gabbe et al., 2002).

Table 2: tree species utilized by Black Drongo at both locations for perching purposes.

Family	Tree Species	Scientific Name	Location-I	Location-II
Indigenous species				
Fabaceae	Phulai	<i>Acacia modesta</i>	5.45%	0.9%
	Indian Rosewood	<i>Dalbergia sissoo</i>	-	2.72%
	Gum Arabic Tree	<i>Acacia nilotica</i>	4.54%	10.9%
	Sukh Chain	<i>Pongamia pinnata</i>	3.63%	1.81%
Myrtaceae	Java Plum	<i>Syzygium cumini</i>	0.9%	-
	Guava	<i>Psidium guajava</i>	2.72%	0.9%
Meliaceae	Chinaberry Tree	<i>Melia azedarach</i>	3.63%	6.36%
Cupressaceae	Cypress spp.	-	1.81%	0.9%
Meliaceae	Indian Lilac	<i>Azadirachta indica</i>	-	3.63%
Anacardiaceae	Mango Tree	<i>Mangifera indica</i>	-	1.81%
Moraceae	Sacred fig	<i>Ficus religiosa</i>	-	1.81%
Rhamnaceae	Indian Plum Tree	<i>Ziziphus mauritiana</i>	2.72%	8.18%
Punicaceae	Pomegranate Tree	<i>Punica granatum</i>	6.36%	1.81%
Exotic species				
Myrtaceae	Eucalyptus	<i>Eucalyptus camaldulensis</i>	15.45%	10.0%
	Bottle Brush	<i>Callistemon citrinus</i>	4.54%	6.36%

Perching Height

A total of 220 observations were carried out for perching height utilization at both locations, 110 from each location. The means of perching heights from the two locations were significant ($p=0.83$). The average perching heights of both locations were not too different (5.88 ± 0.25 m and 5.95 ± 0.23 m respectively). The maximum perching height was observed to be 10 m and 11 m at Location-I and Location-II, respectively.

The minimum height utilized was more at Location-II (1 m) than at Location-I (0.7 m). It may be concluded that Black Drongo is height generalist and can utilize all perches irrespective of their

height which ranges from 0.7-11 m. Table 3 shows the maximum utilized height was between 9-11 m with frequencies of 34.54% and 35.45% at Location-I and Location-II, respectively. It possibly depends upon available food, ease of access and vegetation. Black Drongo used wire as a more common perch with average height of 9.5 m a total of 37% time, this finding is consistent with a previous study by Kaur and Kler (2018). During inference strengthening surveys (other than morning and evening) it was observed that Black Drongo perched near the ground within 0.7-2.5 m to pick insects that are disturbed by gardeners when cutting grasses.

Table 3: Height cohorts and perching height frequencies of Black Drongo at both locations with inter-site difference.

Height Cohorts	Frequency at Location-I		Frequency at Location-II		Inter- site Difference
	Number	Percentage	Number	Percentage	
0.7-2 m	4	3.63%	5	4.54%	1
3-5 m	32	29.09%	30	27.27%	2
6-8 m	36	32.72%	36	32.72%	0
9-11 m	38	34.54%	39	35.45%	1

Inter- and Intra-specific Behaviour

The behaviour of Black Drongo with other birds either sharing same site for feeding or perching within 5 m at both locations was observed. Usually Black Drongo is considered an aggressive and partly territorial bird (eBird, n.d.; BirdLife International, 2020), mostly seen perching on telephone and electrical wires (Grimmett et al., 2009). It was seen that Black Drongo preferred to not share its habitat with other birds and remained solitary ($n=46$) at Location-I. However, during surveys, Black Drongo shared either foraging or perching site with 13 other bird species. It also shared its intimate vicinity with other Drongos along with other bird species. At the Location-I, Black Drongo was found clustered with Jungle Babbler, Crested Lark, White-

Browed Wagtail, Bank Myna and Rufous Treepie, these species were not present in close vicinity of Black Drongo at Location-II.

At Location-II, it was found with a lower frequency for solitary behaviour ($n=34$) compared to clustered ($n=76$) with other species. The frequency to be found with other Black Drongos ($n=26$) was less than Location-I. It might be due to prey abundance at Location-II that restrained them to be clustered at a place. This resulted in a greater number of sharing of habitat with other 17 species. In these 17 species some were not found with Black Drongo at Location-I e.g. Green Bee-eater, Greater Coucal, Cattle Egret, Brown Rock Chat, House Sparrow, Asian koel, Brahminy Starling, Crested Lark and Barn Swallow. The above mentioned seven species were found within 5 m of either

foraging or perching site. It can be inferred from the data that Black Drongo from this area shared their habitat (n=140) with other species more frequently than remaining solitary (n=80) when either

perching or feeding (Table 4). In addition, it was seen that Black Drongo remained relatively less active during the day time with exception of times when special treats of insects given when grass was trimmed.

Table 4: List of bird species found with Black Drongo at either perching site or sharing feeding habitat at both locations.

Order	Common Name	Scientific Name	Location-I	Location-II
Passeriformes	Red vented Bulbul	<i>Pycnonotus cafer</i>	5.12%	8.25%
	House Crow	<i>Corvus splendens</i>	7.17%	8.71%
	Common Myna	<i>Acridotheres tristis</i>	11.28%	11.46%
	Black Drongo	<i>Dicrurus macrocercus</i>	18.97%	16.05%
	Bank Myna	<i>Acridotheres ginginianus</i>	5.12%	-
	Rufous Treepie	<i>Dendrocitta vagabunda</i>	1.25%	-
	Long-tailed Shrike	<i>Lanius schach</i>	5.12%	6.42%
	White Wagtail	<i>Motacilla alba</i>	3.58%	1.37%
	White-browed Wagtail	<i>Motacilla maderaspatensis</i>	1.53%	-
	Brown Rock Chat	<i>Cercomela fusca</i>	-	4.12%
	House Sparrow	<i>Passer domesticus</i>	8.71%	11.0%
	Brahminy Starling	<i>Sturnia pagodarum</i>	-	1.83%
	Jungle Babbler	<i>Turdoides striatus</i>	3.58%	-
	Crested Lark	<i>Galerida cristata</i>	5.64%	-
	Barn Swallow	<i>Hirundo rustica</i>	-	0.92%
Coraciiformes	Green Bee-eater	<i>Merops orientalis</i>	-	5.50%
	Indian Roller	<i>Coracias benghalensis</i>	2.56%	2.29%
Cuculiformes	Asian Koel	<i>Eudynamys scolopaceus</i>	-	0.45%
	Greater Coucal	<i>Centropus sinensis</i>	-	0.45%
Columbiformes	Eurasian Collared-Dove	<i>Streptopelia decaocto</i>	7.17%	8.25%
Charadriiformes	Red-wattled Lapwing	<i>Vanellus indicus</i>	4.10%	3.21%
Ciconiiformes	Cattle Egret	<i>Bubulcus ibis</i>	9.23%	9.63%

Time of Recurrent Attack

The successful capturing was confirmed visually by observing the characteristic swallowing movement with binoculars (Dinsmore, 1973; Grubb, 1976). The mean value 1.93 ± 0.17 minutes suggested that Black Drongo used less time to locate and attack a prey at Location-I compared to Location-II (2.16 ± 0.16 min.). The time of attack was significantly less at both Locations on Wheat (*Triticum aestivum*) and Oat (*Avena sativa*).

Minimum time was 0.4 ± 0.17 min and 0.1 ± 0.16 min at Location-I and

Location-II, respectively. While the maximum time of attack was 11.5 ± 0.17 min at Location-I and 9 ± 0.16 min at Location-II. As discussed earlier, at Location-I Black Drongo stayed solitary and less competition was seen compared to Location-II. Maximum prey was captured between 0.1-3 minutes with frequency of 74.54% and 88.18% at Location-I and Location-II respectively (Table 5). Minimum inter-site difference was for 8-11.5 m. Hence, it may be inferred that time of attack is determined by prey abundance, vegetation density and less competition with conspecifics and other insectivorous species.

Table 5: Time of recurrent attack of Black Drongo to capture a prey.

Time of attack	At Location-I		At Location-II		Inter-site Difference
	Number	Percentage	Number	Percentage	
0.1-3 min	82	74.54%	97	88.18%	15
3-5 min	17	15.45%	8	7.27%	9
5-8 min	9	8.18%	4	3.63%	5
8-11.5 min	2	1.82%	1	0.90%	1

Number of Attempts

Black Drongo successfully captured the prey by its first attempt, with the overall percentage 59.54% (Table 6). At Location-I the mean value was 1.45 ± 0.07 attempts which is less than 1.75 ± 0.09 attempts at Location-II. It was noted that maximum attempts (n=6) were made at Location-I.

Number of attempts show the relative strength of Black Drongo to capture its prey. A greater number of 1st attempts (Table 6) represent its successful

preying. It may be inferred that Black Drongo is adapted to preying on insects. Okosodo et al. (2016) found that diet of Black Drongo constitutes of (86.6%) insects. It was observed that a higher number of attempts showed less target insect concealment and disturbances (that provide stimulus of flight initiation in Black Drongo). Dense vegetation was present at Location-II, which can impact searching of potential prey, Christopher and Whelan (2001) reported similar results where vegetation organization influenced the feeding of insectivorous birds.

Table 6: Number of attempts made to capture prey by Black Drongo at Location-I and Location-II with relation to inter-site difference.

Number of Attempts	At Location-I		At Location-II		Inter- site Difference
	Number	Percentage	Number	Percentage	
1	70	63.64%	61	55.45%	9
2	33	30.0%	31	28.18%	2
3	5	4.54%	8	7.27%	3
4	1	0.90%	7	6.36%	8
5	0	0	3	2.72%	3
6	1	0.90%	0	0	1

Second Perch Site

No previous studies were done on second perch site of Black Drongo. Hence, present study sought to find out the second perch site after effective catching or an unsuccessful attempt. Data showed that out of 220 observations, it returned to the same spot 67.27% times. While failing to retreat on the previous perch site (32.72%) might be due to the unavailability of preferred prey, which compelled them to try for another perch site. At Location-I and Location-II it returned 66.36% and 73.63% of times, respectively.

The success of first attempt may have caused them to perceive the suitability of the site for foraging. Data of Number of Attempts at both locations reinforced the same hypothesis. When Black Drongo attempted more to catch but could not do so, they often retreated on the same spot at Location-II more often than Location-I. The second perch site selection hinges on the availability of potential prey, ease of access, vegetation density (Airola and Barrett, 1985) and competition of feeding (Park et al., 2008). As mentioned earlier there was more potential prey with

ease of access and thus competition was less at Location-II.

Figure 3: Overall percentage of second perch site of Black drongo at both locations.

CONCLUSION

Black Drongo utilized air as a dominant feeding substrate. While most preferred perching on wire, Black Drongo is height generalist when it comes to perching. It utilizes two exotic tree species namely Eucalyptus and Bottle Brush. Black Drongo can share its intimate vicinal habitat with 19 other bird species, which indicate its behavioural plasticity. Perhaps due to location and availability of prey, Black Drongo was observed to be a competent predator of insects. Perch site specificity, especially at the second location, compels them to come at the same perch after attempts to catch prey, depending upon availability of food and other environmental factors. It may be concluded that Black Drongo can best utilize its ecological niche. This study will provide guidance to future studies on Black Drongo.

ACKNOWLEDGEMENT

Author is highly thankful to all those who assisted in the present study, and provided revision of the manuscripts. Especially Haris Mughal, Noman Ali and Tayyab Shehzad, who assisted me in field activities and data collection.

CONFLICT OF INTEREST

Authors declare no conflict of interest in this study.

LITERATURE CITED

- Abbas S, Tabassum R, Khan MZ, Hussnain S, Khan G, and Awan S (2014). Avian Diversity in Central Karakoram National Park, Gilgit-Baltistan. *Int J Agric Biol.* 16 (2): 377-382.
- Airola DA and Barrett RH (1985). Foraging and habitat relationships of insect-gleaning birds in a Sierra Nevada Mixed-Conifer forest. *The Condor*, 87: 205-216.
- Ali S (2002). *The Book of Indian Birds*. Bombay: Bombay Natural History Society, Oxford University Press, Inc. Bombay.
- Arif M and Malik MA (2009). Economic feasibility of proposed cropping patterns under different soil moisture regimes of Pothwar plateau. *Int J Agric Biol.* 11: 27-32.
- Bibi F and Ali Z (2013). Measurement of diversity indices of avian communities at Taunsa Barrage Wildlife Sanctuary, Pakistan. *J Anim Plant Sci.* 23(2): 469-474.
- BirdLife International (2020). Species factsheet: *Dicrurus macrocercus*. Retrived from: <http://www.birdlife.org>
- Blundell VJ (1974). The Wandjina cave paintings of Northwest Australia. *Arct Anthropol.* 11: 213-223.
- Charmantier A, McCleery RH, Cole LR, Perrins C, Kruuk LE and Sheldon BC (2008). Adaptive phenotypic plasticity in response to climate change in a wild bird population. *Science*, 320: 800-803.
- Christopher J and Whelan (2001). Foliage structure influences foraging of insectivorous forest birds: an experimental study. *Ecology*, 82 (1): 219-231.

- Dinsmore JJ (1973). Foraging success of Cattle Egrets, *Bubulcus ibis*. *Am Midl Nat.*, 89(1): 242-246.
- Donald PF, Green MF and Heath MF (2001). Agricultural intensification and the collapse of Europe's bird populations. *P Roy Soc B-Biol Sci.*, 268: 25-29.
- eBird (n.d.). Black Drongo. Retrived on April 19, 2020 from: <http://www.ebird.org/species/bladr01/PK>
- Gabbe AP, Robinson SK and Brawn JD (2002). Tree-species preferences of foraging insectivorous birds: implications for Floodplain forest restoration. *Conserv Biol.*, 16(2): 462-470.
- Ghani A and Maalik S (2020). Assesment of diversity and relative abundace of insect fauna associated with *Triticum aestivum* from district Sialkot, Pakistan. *J. King Saud Univ. Sci.*, 32 (1): 986-995.
- Grimmett R, Roberts T and Inskipp T (2009). *Birds of Pakistan*. Helm Field Guides. Yale University Press.
- Grubb TC (1976). Adaptiveness of foraging in the Cattle Egret. *Wilson Bull.*, 88: 145-148.
- Hairiah K, Sitompul SM, Noordwijk MV and Palm C (2001). Methods for sampling carbon stocks above and below ground. Bogor, Indonesia. Retrieved from: <http://www.icraf.cgiar.org/sea>
- Jhonson MD (n.d.). Evaluation of an arthropod sampling technique for measuring food availability for forest insectivorous birds. *J Field Ornithol.*, 71 (1): 88-109.
- Kaur G and Kler TK (2018). Breeding biology of Black Drongo (*Dicrurus macrocercus*) in relation to nest-site selection and habitat characterization in agricultural landscape. *J Entomol Zool Stud.*, 6 (5): 2142-2147.
- Kaur G and Kler TK (2018). Feeding behaviour and perching preferences of black drongo (*Dicrurus macrocercus*) in Ludhiana district (Punjab). *J Entomol Zool Stud.*, 6 (4): 232-239.
- Kaur S and Kler TK (2019). Food and foraging niches of carnivorous bird species in orchards: implications for their conservation. *J Anim Res.*, 9 (2): 325-334.
- Khan BN and Ali Z (2015). Assessment of birds' fauna, occurrence status, diversity indices and ecological threats at Mangla dam, AJK. *J Anim Plant Sci.*, 25 (3): 397-403.
- Kour DN and Sahi DN (2012). Studies on the community ecology of cattle egrets *Bubulcus ibis coromandus* (Boddaert) in Jammu (Jammu and Kashmir), India. *Int J Biodivers Conserv.*, 4 (13): 439-445.
- Mayr E and Amadon D (1951). A classification of recent birds. *Am Mus Novit.*: 1-42.
- Okosodo EF, Orimaye JO and Odewumi OS (2016). Diet and foraging ecology of Fork Tailed Drongo (*Dicrurus adsimilis*) in Leventis Foundation Nigeria, Agricultural School South West Nigeria. *Int J Environ Agric Biotech.*, 1 (2): 252-256.
- Park C, Hino T and Ito H (2008). Prey distribution, foliage structure, and foraging behavior of insectivorous birds in two oak species (*Quercus serrata* and *Q. variabilis*). *Ecol Res.*, 23: 1015-1023.
- Sarwar M (2019). Density and diversity of birds in the agricultural landscape of Pothwar Plateau, Pakistan. *J Anim Plant Sci.*, 29 (3): 864-870.
- Sekercioglu CH (2002). Forest fragmentation hits insectivorous birds hard. *Sci World J.*: 62-64.
- Sidra S, Ali Z and Chaudry MN (2013). Avian diversity at new campus of Punjab University in relation to

- land use change. Pak J Zool., 45 (4): 1069-1082.
- Verner J (1985). Assesment of counting techniques. Current Ornithology, vol. 2, R.F. Johnston Ed., Plenum Press, New York: pp. 247-302.