Plants and Their Uses; How Communities of Tolipir, Pakistan Get Benefit

Abu ul Hassan Faiz

Women University of Azad Jammu and Kashmir, Bagh, sabulhussan@gmail.com

Mehboob ul Hassan

Department of Education, University of Punjab, Pakistan

Fida Muhammad Khan

Quaid-i-Azam University, Islamabad, Pakistan

Lariab Zahra Faiz

Women University of Azad Jammu & Kashmir, Bagh

Hina Farooq

Women University of Azad Jammu & Kashmir, Bagh

Follow this and additional works at: <https://corescholar.libraries.wright.edu/jbm>

Part of the [Biodiversity Commons](#), [Biology Commons](#), and the [Population Biology Commons](#)

Recommended Citation

Faiz, A. H., Hassan, M. u., Khan, F. M., Faiz, L. Z., & Farooq, H. (2018). Plants and Their Uses; How Communities of Tolipir, Pakistan Get Benefit, *Journal of Bioresource Management*, 5 (3).

DOI: <https://doi.org/10.35691/JBM.8102.0090>

ISSN: 2309-3854 online

(Received: Oct 19, 2019; Accepted: Oct 20, 2019; Published: Jul 1, 2018)

This Article is brought to you for free and open access by CORE Scholar. It has been accepted for inclusion in Journal of Bioresource Management by an authorized editor of CORE Scholar. For more information, please contact library-corescholar@wright.edu.

Plants and Their Uses; How Communities of Tolipir, Pakistan Get Benefit

© Copyrights of all the papers published in Journal of Bioresource Management are with its publisher, Center for Bioresource Research (CBR) Islamabad, Pakistan. This permits anyone to copy, redistribute, remix, transmit and adapt the work for non-commercial purposes provided the original work and source is appropriately cited. Journal of Bioresource Management does not grant you any other rights in relation to this website or the material on this website. In other words, all other rights are reserved. For the avoidance of doubt, you must not adapt, edit, change, transform, publish, republish, distribute, redistribute, broadcast, rebroadcast or show or play in public this website or the material on this website (in any form or media) without appropriately and conspicuously citing the original work and source or Journal of Bioresource Management's prior written permission.

PLANTS AND THEIR USES; HOW COMMUNITIES OF TOLIPIR, PAKISTAN GET BENEFIT

ABU UL HASSAN FAIZ^{*1}, MEHBOOB UL HASSAN², FIDA MUHAMMAD KHAN³, LARIAB ZAHRA FAIZ¹ AND HINA FAROOQ¹

¹Women University of Azad Jammu and Kashmir, Bagh, AJK, Pakistan

²Department of Education, University of Punjab, Pakistan

³Department of Microbiology, Quaid-i-Azam University, Islamabad, Pakistan

*Corresponding author: sabulhussan@gmail.com

ABSTRACT

The present study documented ethnobotanical uses of 156 plant species belonged to 93 genera and 51 families listed from five villages of Tolipir landscape (Ali Sojhal, Kanchi Kot, Khorri Chana, Kahoo Kot and Noor Kot villages). Study revealed that most of the species were used for medical treatment, where leaves found to be the most frequently used part for preparation of indigenous recipes and fodder purpose. The current study empirically contributes a significant amount of ethnobotanical knowledge and depicts its strong connection with indigenous traditions. It is an urgent need to document indigenous uses of plants for future domestication.

Key Words: Ethnobotany, Tolipir, Indigenous knowledge, Conservation

INTRODUCTION

Wild plants have been used for human benefit such as food, medicine, fuel and timber from a very long time. The utilization of wild plants to fulfill their needs is prevalent in many rural communities of the world and often they have considerable knowledge about their uses (Ahmad et al., 2014). Most of the population from the Third world countries rely on their ethnobotanical knowledge for medical treatment (Luitel et al., 2014). The use of plants as medicine is slowly increasing in the developed world because they have minor or no side effects (Jordan et al., 2010).

The traditional knowledge of medical use of flora possessed by locals is an important asset that can be used as a baseline for further research into plant-use for treatment (Etkin et al., 1993). The documentation of traditional knowledge aids in the preservation of indigenous culture identifies threatened species and contributes

to the conservation and management of plant diversity (Watson et al., 2003). No previous study about ethnological aspects of plants was conducted in Tolipir landscape except phytosociology of the area (Faiz et al., 2014, 2015). Therefore, in the present study, we aimed to document indigenous knowledge of the use of medicinal plants in the Tolipir landscape.

MATERIAL AND METHODS

Tolipir landscape is one of the biodiversity hotspots of Azad Jammu and Kashmir. It is located at longitude 73° 53 5.316" east and latitude 33° 32 9.925" north. The altitude ranges from 500-800 meters above sea level. The area falls under dry Sub-tropical type in which *Olea ferruginea*, *Accacia modesta* and *Pinus roxburghii* (Chir Pine) are the most dominant tree species in the area (Kharkwal and Rawat, 2010). The vegetation in the area comprises of a wide variety of trees, herbs, shrubs and climbers. Ground cover comprises of a wide variety of

angiosperms along with ferns and mosses. Field surveys were conducted during August 2013-July 2014. Visual Appraisal Approach (VAA) and Rapid Rural Appraisal (RRA) methods were employed to document ethnobotanical information through oral interviews and semi-structured questionnaires from local herbalists (Hakeems) and the elderly people who were familiar with traditional uses of plants particularly for medicines, fodder, fuel, timber, food, cosmetics and etc. by following (Martin et al., 1995). The utility value of plants was calculated by using the following statistical formula:

$$UVs = \Sigma Uis / nis.$$

Where (Uis) stands for single informant and (nis) stands for the total number of people interviewed about that particular species.

Figure 1: A simplified satellite imaginary of Tolipir landscape with sampling villages Ali Sojhal, Kanchi Kot, Khor Chana, Kahoo Kot and Noor Kot (source: Arc View 3.3 and Google Earth pro 4.2).

RESULTS AND DISCUSSION

The present survey found a list of 158 plants species and most of the plants were used as dual purpose according to respondent of local inhabitants of Tolipir landscape. The

list of the plants and their respective uses is outlined in Table 1.

Ethnobotany is the study of traditional pattern of exploitation of indigenous plants by the local populace of a particular culture or region. Ethnobotanical information helps ecologists, pharmacologist, taxonomists, watersheds managers and wildlife workers in their efforts for improving the plant wealth of an area (Faiz et al., 2014).

Tolipir landscape has a greater number of plants (158 species) than any other study area of Kashmir previously recorded such as 36 plants in Leepa valley (Ishtiaq et al., 2012) and 40 plant species in Neelum valley, (Adeel et al., 2011). It is imperative that this diverse richness of species be conserved for the healthy maintenance of the ecosystem. The plants used for ailments of the skin include *Albizia lebbek*, *Berberis lyceum*, *Fumaria indica*, *Myrsine africana*, *Plantago lanceolata*, *Rumex dentatus* and *Verbascum thapsus*. The plants used for treating different bowel illnesses include *Albizia lebbek*, *Berberis lyceum*, *Mentha royleana*, *Ficus palmata*, *Fumaria indica*, *Myrsine africana*, *Trichodesma indicum*, *Verbascum thapsus*, *Valeriana jatamansi*, *V. pyrolifolia* and *Zanthoxylum armatum*. *A. indica*, *Juglans regia*, *Sarcococca sligna* and *Trichodesma indicum* are used for aching joints. These are just some of the uses of the select few plants. Further studies are required to investigate the other possible uses of these plants, their sustainable cultivation and production of medicines for masses.

Pyrus malus, *P. pashia*, *Prunus armeniaca*, *Punica granatum*, *Juglans regia*, *P. persica*, *Ficus palmata*, *Viburnum grandiflorum*, *Rubus fruticosus*, *Duchesnea indica* and *Fragaria nubicola* are used in diet.

Abies pindrow, *P. roxburgii*, *P. wallichiana*, *Populus ciliata* and *J. regia* are used for furniture and constructional

purposes. *A. indica*, *Clematus grata*, *Salix acmophylla*, *Dicliptera bupleuroides*, *Cederala serrata* and *Q. incana* are utilized for fuel.

A. indica, *A. lebbek*, *Prunus persica*, *Viburnum grandiflorum*, *Myrsine Africana*, *Adiantum capillus-veneris*, *Celtis caucasica*, *Hedera nepalensis*, *Oenothera rosea*, *Rumex hastatus*, *Trifolium repens*, *Indigofera heterantha*, *Urtica dioica* and *Gerbera gossypina* are given as fodder to cattle.

Almost all the plant species of Tolipir landscape are being traditionally exploited by the local populace in different ways, i.e., medicine, fodder/ forage, timber and furniture wood, human food, fuel wood, construction material, ornamental, vegetable and flavouring agents, etc., to different degrees. Not all the flora has a fast growth rate and/or high population density; this puts the species readily cultivated for use at risk (Hunde et al., 2004). This is especially crucial where species with multiple known uses are concerned (Sarwat et al., 2012).

It is imperative that policies be made and implemented to secure rare species of plants. To reduce the consumption of firewood, the locals need to be presented with affordable alternatives.

Figure 1. *Aesculus indica*

Figure 2. *Albizia lebbek*

Figure 3. *Quercus incana*

Table 1. List of plants and use of their different parts

Name of Plant species	Leaves/Branches	Roots	Stem	Fruits	Bark	Seeds	Flwr	Resin/LateN/Gum	Whole plant	Ornamental	Food	Fodder	Fuel	Timber	Av Tools	Medicinal	Hedge/Fencing	Roof Thatching	Tea/Coffee substitute	Cosmetic/Beautification	Honey bee spp.	Other (Paper pulp, match industry, dye, tool handles, utensils, rope making, furniture, boats, carts, miswak, insect repellent, soil binder/ground cover palnt etc.)
<i>Abies pindrow</i>	N	N	N	N	N	N	N	N	Y	Y	N	N	Y	Y	N	Y	N	N	N	N	N	Y
<i>Acer pentapomicum</i>	Y	N	Y	N	N	N	N	N	N	N	N	Y	Y	Y	Y	N	N	N	N	N	N	N
<i>Aesculus indica</i>	Y	N	Y	Y	Y	N	N	N	N	N	N	Y	Y	Y	Y	Y	N	N	N	N	N	Y
<i>Albizia lebbeck</i>	Y	N	N	N	Y	Y	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Castanea sativa</i>	Y	N	Y	Y	N	N	N	N	N	N	Y	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Cedrella serrata</i>	Y	N	Y	N	Y	N	N	N	N	N	N	Y	Y	N	N	Y	N	Y	N	N	N	N
<i>Celtis caucasica</i>	Y	N	Y	Y	N	N	N	N	N	N	Y	Y	Y	N	Y	Y	N	Y	N	N	N	N
<i>Dalbergia sissoo</i>	Y	N	Y	N	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	Y
<i>Elaeagnus angustifolia</i>	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	Y
<i>Elaeagnus umbellata</i>	Y	N	N	N	N	Y	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Eriobotrya japonica</i>	Y	N	N	Y	N	N	Y	N	N	N	Y	N	N	N	N	Y	N	N	Y	N	N	Y
<i>Ficus carica</i>	N	N	N	Y	N	N	N	Y	N	N	Y	Y	Y	N	N	Y	N	N	N	N	N	N
<i>Ficus palmate</i>	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	Y	N	N
<i>Juglans regia</i>	Y	Y	Y	Y	Y	Y	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	Y
<i>Nerium oleander</i>	Y	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Pinus roNburgii</i>	Y	N	Y	N	Y	Y	N	Y	N	N	Y	N	Y	Y	N	Y	N	Y	N	N	N	Y
<i>Pinus wallichiana</i>	Y	N	Y	N	N	N	N	Y	N	N	N	N	N	Y	N	Y	N	N	N	N	N	Y
<i>Pistacia chinensis</i>	Y	N	Y	N	Y	N	N	Y	N	N	N	Y	Y	Y	N	Y	N	Y	N	Y	N	Y
<i>Populus ciliata</i>	Y	N	Y	N	N	N	N	N	Y	Y	N	Y	Y	Y	N	N	N	N	N	N	N	Y

<i>Prunus armeniaca</i>	Y	N	Y	Y	N	Y	N	N	N	N	Y	Y	Y	N	Y	Y	N	N	N	N	Y
<i>Prunus domestica</i>	Y	N	Y	Y	N	Y	Y	Y	N	N	Y	Y	Y	N	N	Y	N	N	N	Y	N
<i>Prunus persica</i>	Y	N	Y	Y	N	N	N	N	N	N	Y	Y	Y	N	N	Y	N	N	N	N	N
<i>Punica granatum</i>	Y	N	Y	Y	Y	Y	N	N	N	N	Y	Y	Y	N	Y	Y	Y	Y	N	N	N
<i>Pyrus malus</i>	N	N	Y	Y	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	Y	Y
<i>Pyrus pashia</i>	N	N	Y	Y	N	N	N	N	N	N	Y	N	Y	N	Y	Y	Y	N	N	Y	N
<i>Quercus baloot</i>	Y	N	Y	N	Y	N	N	N	N	N	N	Y	Y	Y	Y	Y	N	N	N	N	N
<i>Quercus dilatata</i>	Y	N	Y	Y	Y	N	N	N	N	N	N	Y	Y	N	N	Y	N	N	N	N	N
<i>Quercus glauca</i>	Y	N	Y	N	N	N	N	N	N	N	N	Y	Y	Y	N	N	N	N	N	N	N
<i>Quercus incana</i>	Y	N	Y	N	Y	N	N	N	N	N	N	Y	Y	Y	Y	Y	N	Y	N	N	N
<i>Robinia pseudo-acacia</i>	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	Y	N	N	N	N	N
<i>SaliN acmophylla</i>	Y	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N
<i>SaliN denticulate</i>	Y	N	Y	N	Y	N	N	N	N	N	N	Y	Y	N	Y	Y	N	N	N	N	N
<i>Sapindus mukorossi</i>	N	N	N	Y	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N
<i>Skimmia laureola</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N
<i>Ziziphus spp</i>	Y	N	Y	Y	N	N	N	N	N	N	Y	Y	Y	N	N	Y	N	N	N	Y	Y
<i>Berberis lycium</i>	Y	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N
<i>Clematis b Buchananiana</i>	Y	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N
<i>Clematis connata</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
<i>Clematis montana</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Debregeasia salicifolia</i>	Y	N	Y	Y	Y	N	N	N	N	N	Y	N	Y	N	N	Y	N	N	N	N	N
<i>Desmodium polycarpum</i>	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N
<i>Heracleum cachemirica</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Indigofera heterantha</i>	Y	N	Y	N	N	N	Y	N	N	N	N	Y	Y	N	N	N	N	N	N	N	N
<i>Isodon rugosus</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N
<i>Jasminum mesnyi</i>	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	Y	N	N	N	Y	N
<i>Lagustrum lucidam</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N
<i>Myrsine africana</i>	N	N	N	N	N	N	N	N	Y	N	Y	N	Y	N	N	Y	N	Y	N	N	N
<i>Rosa bruNnii</i>	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	Y	N	N	N	N	Y
<i>Rubus fruticosus</i>	Y	N	N	Y	Y	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N

<i>Rubus niveus</i>	Y	N	N	Y	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	Y
<i>RumeN hastatus</i>	Y	Y	N	N	N	N	N	N	N	N	N	Y	Y	N	N	Y	N	N	N	N	N	N
<i>Sarcococca saligna</i>	Y	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Sophora mollis</i>	N	N	Y	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	Y	N	N
<i>Viburnum nervosum</i>	N	N	N	N	N	N	N	N	Y	N	Y	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Viburnum cotinifolium</i>	N	N	N	N	N	N	N	N	Y	N	Y	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Viburnum grandiflorum</i>	Y	N	N	Y	N	Y	N	N	N	N	Y	Y	Y	N	N	Y	N	N	N	N	N	N
<i>ZanthoNylum armatum</i>	Y	N	N	Y	N	Y	Y	N	N	N	Y	N	Y	N	N	Y	Y	N	N	N	N	Y
<i>Achellia millefolium</i>	Y	N	N	N	N	N	Y	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Ajuga bracteosa</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Allium griffithianum</i>	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	N
<i>Alysicarpus bupleurifolius</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Anaphalis adnata</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Androsace rotundifolia</i>	Y	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Anemone tetrasepala</i>	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Aquilegia pubifoera</i>	N	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Artemisia absinthium</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	Y	N	N	N	N	N	Y
<i>Artemisia maritime</i>	Y	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	Y
<i>Artemizia dubia</i>	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Asparagus filicinus</i>	Y	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	N
<i>Bergenia ciliata</i>	N	N	N	N	N	N	N	N	Y	N	Y	N	N	N	N	Y	N	N	N	N	N	N
<i>Caltha alba</i>	N	N	N	N	N	N	N	N	Y	Y	N	N	Y	N	N	Y	N	N	N	N	N	N
<i>Campanula benthamii</i>	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Cirsium falconeri</i>	N	N	N	N	N	N	N	N	Y	N	Y	N	Y	N	N	N	N	N	N	N	N	N
<i>Conyza bonariensis</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Dicliptera bupleuroides</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Dioscorea bulbifera</i>	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	N	Y
<i>Dioscorea deltoidea</i>	N	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Duchesnea indica</i>	N	N	N	Y	N	N	N	N	N	N	Y	Y	N	N	N	Y	N	N	N	N	N	N
<i>Epilobium tibetanum</i>	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	Y	N	N	Y

<i>Euphorbia helioscopia</i>	N	N	N	N	N	Y	N	Y	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Euphorbia wallichii</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Fragaria nubicola</i>	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	N
<i>Fumaria indica</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Gallium aparine</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Gallium asperifolium</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Gerbera gossypina</i>	Y	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Heracleum candicans</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	Y	N	N	Y	N	N	N	N	N	Y
<i>Hypericum perforatum</i>	Y	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	Y
<i>Iris milesii</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N
<i>Launia secunda</i>	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
<i>Lespedeza juncea</i>	N	N	N	N	N	N	N	N	Y	N	Y	Y	N	N	N	Y	N	N	N	N	N	N
<i>Malvestrum coromendelianum</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Medicago minima</i>	N	N	N	N	N	N	N	N	Y	N	Y	Y	N	N	N	N	N	N	N	N	N	N
<i>Melilotus alba</i>	Y	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	Y	N	N	N	N	N	N
<i>Mentha royleana</i>	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N	N
<i>Nepeta erecta</i>	Y	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Nepeta laevigata</i>	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Nepeta nervosa</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	Y	N
<i>Oenothera rosea</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Parthenium hysterophorus</i>	N	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Pimpinella stewartii</i>	N	N	N	Y	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Pimpinella stewartii</i>	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Podophyllum emodi</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Polygonatum multiflorum</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Primula denticulata</i>	N	N	N	N	N	N	N	N	Y	Y	Y	Y	N	N	N	N	N	N	N	N	N	N
<i>Prunella vulgaris</i>	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N
<i>Ranunculus arvensis</i>	Y	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	N	N	N	N	N
<i>Ranunculus hirtellus</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	N

<i>Ranunculus muricatus</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>RumeN dentatus</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	Y	N	N	Y	N	N	N	N	N	N
<i>Salvia hians</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Salvia lanata</i>	N	N	N	N	N	N	N	N	Y	N	Y	N	Y	N	N	Y	N	N	N	N	N	N
<i>Salvia moorcroftiana</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Sambucus wightiana</i>	N	N	N	Y	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Saussurea candolleana</i>	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	Y
<i>Senecio chrysanthemoides</i>	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N	N	N	Y
<i>Seseli libaNtis</i>	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	N	N
<i>Swertia ciliata</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>TaraNacum officinale</i>	Y	Y	N	N	N	N	N	N	N	N	Y	Y	N	N	N	Y	N	N	N	N	N	N
<i>Thalictrum pedunculatum</i>	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
<i>Thymus liniaris</i>	Y	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Trichodesma indicum</i>	Y	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N	N
<i>Triufoium repens</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N
<i>Urtica dioica</i>	Y	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	N	N	N	N	Y
<i>Valeraina jatamansi</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Valeriana pyrolifolia</i>	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N
<i>Verbascum Thapsus</i>	N	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N
<i>Veronica beccabunga</i>	Y	N	Y	N	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Veronica biloba</i>	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	N
<i>Veronica melissifolia</i>	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
<i>VincetoNicum hirundinaria</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	Y	N	N
<i>Viola canscens</i>	Y	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Viola pilosa</i>	Y	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N
<i>Brachiaria spp.</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N
<i>Dactyloctenium aegyptium</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N
<i>Desmostachya bipinnata</i>	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	Y	N	N	N	Y
<i>Koeleria spp.</i>	Y	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	Y
<i>Oplismenus spp.</i>	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	N	N

Faiz et al. (2018). Plants and Their Uses by Tolipir Communities
J Biores Manag. 5(3): 1-11

<i>Phalaris miNr</i>	N	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	N	N	N	N	N	Y
<i>Phragmites karka</i>	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	Y	N	N	N	N
<i>Poa nepalensis</i>	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	Y	N	N	N	N	N
<i>Pogonatherum spp.</i>	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N
<i>Stipa sibirica</i>	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	N	N	Y
<i>Themeda spp.</i>	N	N	N	N	N	N	N	N	N	Y	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Adiantum capillus-veneris</i>	Y	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Adiantum incisum</i>	N	N	N	N	N	N	N	N	N	Y	Y	N	N	N	N	N	Y	N	N	N	N	N
<i>Adiantum venustum</i>	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	Y	N
<i>Athyrium tenuifrons</i>	N	N	N	N	N	N	N	N	N	Y	N	Y	N	N	N	N	Y	N	N	N	N	Y
<i>Coniogramme rosthornii</i>	N	N	N	N	N	N	N	N	N	Y	Y	Y	N	N	N	N	N	N	N	N	N	N
<i>Dryopteris juNtaposita</i>	Y	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	N	N
<i>Equisetum arvense</i>	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y
<i>Polystichum squarrosus</i>	Y	Y	N	N	N	N	Y	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y
<i>Pseudophagopteris pyrrhorhachis</i>	Y	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Pteris cretica</i>	Y	N	Y	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	Y
<i>Clematis grata</i>	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N	N
<i>Hedera nepalensis</i>	Y	N	N	N	N	N	N	N	N	N	N	N	Y	N	N	N	Y	N	N	N	N	N
<i>Momordica dioica</i>	N	Y	N	Y	N	N	N	N	N	N	N	Y	N	N	N	N	Y	N	N	N	Y	N
<i>SmilaN glaucophylla</i>	N	N	N	N	N	N	N	N	N	Y	N	N	N	N	N	N	Y	N	N	N	N	N

CONCLUSION

This study documents ethnobotanical uses of 156 plant species belonged to 93 genera and 51 families listed from five villages of Tolipir landscape (Ali Sojhal, Kanchi Kot, Khor Chana, Kahoo Kot and Noor Kot villages). It was found that most of the species were used for medical treatment.

REFERENCES

- Adeel M, Malik RN, Shinwari ZK, Mahmood A (2011). Ethnobotanical survey of plants from Neelum, Azad Jammu & Kashmir, Pakistan. *Pak. J. Bot.* 43: 105-110.
- Ahmad M, Sultana S, Fazl-I-Hadi S, Ben HT, Rashid S, Zafar M (2014). An ethnobotanical study of medicinal plants in high mountainous region of Chail Valley (District Swat-Pakistan). *J Ethnobiol Ethnomed.* 10: 36. doi:10.1186/1746-4269-10-36
- Etkin NL (1993). Anthropological methods in ethnopharmacology. *J Ethnopharmacol.* 38: 93-104.
- Faiz AH, Ghufuran MA, Mian A, Akhtar T (2014). Floral Diversity of Tolipir National Park (TNP), Azad Jammu and Kashmir, Pakistan. *Biologia.* 60 (1): 43 – 55
- Faiz AH, Abbas F, Zahra L (2015) Avifaunal diversity of Tolipir national park, azad Jammu and Kashmir, Pakistan. *J. Anim. Plant sci.* 25 (3supp2.): 404-409
- Faiz AH, Abbas F, Zahra L (2015) Community structure and diversity of butterflies in Tolipir National park, azad Jammu and Kashmir (AJK) Pakistan. 25(3 supp. 2): 355-358
- Faiz AH, Abbas F, Zahra L, (2015). Anthropogenic influences on the tolipir landscape, lesser Himalayas, Pakistan. *J. Bio res. manage.* 2(4): 20-27
- Faiz AH, Abbas F (2016). Mammalian diversity of Tolipir national park, Azad Jammu and Kashmir, Pakistan. 48(4): 1209-1212.
- Hunde, DZ, Asfaw, Kelbessa E (2004). Use and management of medicinal plants by indigenous people in ‘Boosaf’, Welenchiti area. *Ethiop. J. Biol. Sci.* 3: 113-132.
- Ishtiaq M, Mumtaz SA, Tanveer H, Ghani A (2012). Medicinal Plant Diversity in the Flora of Leepa Valley, Muzaffarabad (AJK), Pakistan. *Afr. J. Biotechnol.* 11(13): 3087-3098.
- Jordan SA, Cunningham DG, Marles RJ (2010). Assessment of herbal medicinal products: Challenges, and opportunities to increase the knowledge base for safety assessment. *Toxicol Appl Pharmacol.* 243: 198–216.
- Kharkwal G and Rawat YS (2010). Structure and composition of vegetation in subtropical forest of Kumaun Himalaya. *Afr. J. Plant Sci.* 4(4): 116-121.
- Luitel RD, Rokaya MB, Timsina B, Münzbergová Z (2014). Medicinal plants used by the Tamang community in the Makawanpur district of central Nepal. *J Ethnobiol Ethnomed.* doi: 10.5.1186/1746-4269
- Martin JG (1995). *Ethnobotany: A methods manual.* Chapman and Hall, New York.
- Khan MA, Ajab M. Mujtaba KG, Hussain M (2012). Ethnobotanical study about medicinal plants of poonch valley azad kashmir. *J. Anim. Plant Sci.* 22(2): 493-500.
- Sarwat, Shinwari ZK, Ahmad N (2012). Screening of potential medicinal plants from district Swat specific for controlling women diseases. *Pak. J. Bot.* 44(4): 1193-1198.

Watson A, Alessa L, Glaspell B (2003). The relationship between traditional ecological knowledge, evolving cultures, and wilderness protection in the circumpolar north. *Conserv Ecol.* 8:2.