Definition 3.2

A simple graph in which every pair of vertices adjacent is called a complete graph.

Theorem 3.1:

The complete graph K_n is a Zumkeller graph when $n \equiv 2 \pmod{4}$ and $n \leq 10$.

Proof:

Let $V = \{v_i \mid 1 \leq i \leq n\}$ be the vertex set and $E = \{e_{ij} \mid 1 \leq i \leq n, 1 \leq j \leq n \text{ and } i < j\}$ be the edge set of the complete graph K_n .

Define an injective function $f : V \rightarrow N$ such that

$$f(v_1) = \frac{n}{2} \left(\frac{n}{2} + 1 \right)$$

$$f(v_i) = \left(\frac{n}{2} + i - 1 \right) \left(\frac{n}{2} + i \right) \text{ for } 2 \leq i \leq n$$

Define an induced function $f^* : E \rightarrow N$ such that $f^*(e_{ij}) = f^*(v_i v_j) = f(v_i) f(v_j)$.

Let $f(v_i) = a \times b$ and $f(v_j) = c \times d$ where a, b and c, d are a pair of consecutive integers.

Claim: To prove $f(v_i) f(v_j) = abcd$ is a Zumkeller number.

Now,

$$f(v_n) = \left(\frac{n}{2} + n - 1 \right) \left(\frac{n}{2} + n \right)$$

Since $n \leq 10$,

$$\text{We have } f(v_n) \leq 14 \times 15$$

Here the integers in the product of two consecutive integers are less than or equal to 15. All odd numbers other than 9 and 15 are primes in this product and in each pair of consecutive numbers there is an odd number.

Therefore, the product $abcd$ of those numbers have the prime factorisation of the form $2^k p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ for which at least one k_i is odd.

Now we prove that the products of any two numbers in the set $\{8 \times 9, 9 \times 10, 14 \times 15\}$ are also the same form of prime factorisation.

$$(i) \quad 8 \times 9 \times 9 \times 10 = 2^4 3^4 5,$$

$$(ii) \quad 8 \times 9 \times 14 \times 15 = 2^4 3^3 5$$

$$(iii) \quad 9 \times 10 \times 14 \times 15 = 2^2 3^3 5 7$$

Therefore, all the edges in the complete graph K_n have the prime factorisation of the form $2^k p_1^{k_1} p_2^{k_2} \dots p_m^{k_m}$ in which at least one k_i is odd.

By the property of Zumkeller numbers, $abcd$ is a Zumkeller number.

Hence the complete graph K_n admits a Zumkeller labeling when $n \equiv 2 \pmod{4}$ and $n \leq 10$.

An illustration of this theorem is shown in the example 3.1.

Example 3.2

The graph shown in the Fig.3.1 is a Zumkeller labeling of K_6

Fig.3.1. Zumkeller labelling of K_6

Definition 3.3

The one-point union of two complete graphs is a graph $K_n^{(2)}$ ($n \geq 3$) consists of two copies of K_n sharing a common vertex. It contains $2n-1$ vertices and $n(n-1)$ edges.

Theorem 3.2:

The graph $K_n^{(2)}$ admits a Zumkeller labeling when $n \leq 6$.

Proof:

Let $V = \{v_i \mid 1 \leq i \leq n-1\} \cup \{u_i \mid 1 \leq i \leq n-1\} \cup \{v_0\}$

be the vertex set and

$$E = \{e_{ij} \mid 1 \leq i \leq n-1, 1 \leq j \leq n-1 \text{ and } i < j\}$$

$$\cup \{e'_{ij} \mid 1 \leq i \leq n-1, 1 \leq j \leq n-1 \text{ and } i < j\}$$

$$\cup \{e_{0i} \mid 1 \leq i \leq n-1\} \cup \{e_{0j} \mid 1 \leq j \leq n-1\}$$

be the edge set of the graph $K_n^{(2)}$.

Define an injective function $f : V \rightarrow N$ such that

$$f(v_0) = 8 \times 9$$

$$f(v_i) = (i+2) \times (i+3)$$

$$f(u_i) = (i+8) \times (i+9)$$

Define an induced function $f^* : E \rightarrow N$ such that

$$f^*(e_{ij}) = f^*(v_i v_j) = f(v_i) f(v_j)$$

$$f^*(e'_{ij}) = f^*(u_i u_j) = f(u_i) f(u_j)$$

$$f^*(e_{0i}) = f^*(v_0 v_i) = f(v_0) f(v_i) = 8 \times 9 f(v_i)$$

$$f^*(e_{0j}) = f^*(v_0 v_j) = f(v_0) f(v_j) = 8 \times 9 f(v_j)$$

All the edges receive the Zumkeller numbers. This can be proved as in Theorem 3.1.

Hence The graph $K_n^{(2)}$ admits a Zumkeller labeling when $n \leq 6$.

Theorem 3.2 is illustrated in the following example 3.2.

Example 3.3

The Fig.3.2 shows the existence of the Zumkeller labeling for the graph $K_6^{(2)}$.

13. Rosa, A.: On certain valuations of the vertices of a graph. In N.B. Gordan and Dunad, editors, Theory of graphs. International Symposium. Paris (1966) 349–359.
14. Yuejian Peng, Bhaskara Rao, K.P.S.: On Zumkeller numbers. Journal of Number Theory. 133(4) (2013) 1135–1155.
15. Zongheng Zhou, S. Das and H. Gupta.: Connected k-coverage problem in sensor networks, Proceedings. 13th International Conference on Computer Communications and Networks (IEEE Cat. No.04EX969), Chicago, IL, (2004) 373-378

AUTHORS PROFILE

Dr.B.J.Balamurugan, received his Ph.D degree in Mathematics from the University of Madras, Chennai, India. Currently, he is an Assistant Professor(Senior) of Mathematics in the School of Advanced Sciences at VIT University, Chennai Campus, Chennai, India. He has more than 22 years of teaching experience at Undergraduate and Postgraduate level courses.

Dr.B.J.Balamurugan has published more than 32 research papers in various journals and conference proceedings. His research interest includes graph theory, graph grammars, fuzzy logic and Petri nets.

R. Madura Meenakshi received her M.Sc and M.Phil degrees in Mathematics from Avinashilingam University, Coimbatore, Tamil Nadu, India. Currently, she is a research scholar at SSN College of Engineering, Chennai, India. Her research interest includes Graph Theory, Image Processing and Machine Learning.