No. 76, February 2019

Doi: 10.36071/clivar.76.2019

CLIVAR Exchanges

Special Issue: CLIVAR-FIO Summer Course on "Past, Present and Future Sea Level Changes"

CLIVAR (Ocean and Climate: Variability, Predictability and Change) is the World Climate Research Programme's core project on the Ocean-Atmosphere System

International
Science Council

Editorial

Jose Santos, Liping Yin

International CLIVAR Project Office, Qingdao, China

In order to fulfill CLIVAR's mission (to understand the dynamics, the interaction, and the predictability of the climate system with emphasis on ocean-atmosphere interactions). It is important to foster international collaboration among the global scientific community not only between established senior scientist but also identifying the next generation of researchers who will continue these endeavors in the years to come.

The next generation of scientists will play an important and even leading role in the future. Due to the global climate change, the society and environment in which we live are encountering huge threat, such as the sea level rise, and the scientific communities are meeting more formidable challenges on the climate research. One of the challenges is the urgent need to foster the future generation of researchers. To achieve this, the capacity building for the young scientists, which aims to promote their research ability and expertise, to increase their exposure to the scientific communities, to provide a platform for the data and methods sharing, and to enhance the international communication and cooperation, is now an indispensable part of CLIVAR mission.

With this in mind, CLIVAR is continuously looking for opportunities for young scientists with outstanding research potential to interact, to learn from their peers and senior scientists what is required to better understand aspects of the climate system and address research challenges of societal relevance, and to build lasting relationships and collaborations.

With the strong emphasis of CLIVAR on capacity building in particular geared towards Early Career Scientists (ECS), CLIVAR has made great efforts in all aspects, including the creating of the "Early Career Scientists Network" homepage, providing endorsement for activities and projects, co-organizing the WESTPAC Training Course and organizing the Early Career Scientists Symposium (ECSS); Especially, CLIVAR has established a plan to offer a Series of Summer Courses; on even years in collaboration with the First Institute of Oceanography (China), and on odd years with the International Centre for Theoretical Physics (Italy). The 2018 event was titled "Summer Course on Past, Present and Future Sea Level changes", and was held in June 2018 in Qingdao; with the purpose of providing early career scientists and engineers specializing in sea level research with an update in observations, knowledge, and understanding for the study of global and regional sea level changes and their impacts in coastal areas.

A total of 202 applications coming from 39 countries

for participation in the summer school were received. During the selection process, the Organizing Committee assessed the relevance of the applicants' potential contributions to the summer school. Throughout the selection process, the Committee considered potential for future knowledge transfer an important factor, the education background as well as the scientific expertise was also considered. We also aimed to achieve the widest possible geographical distribution of the trainees, and a gender balanced distribution. Improving and increasing the capability of developing countries was an additional aim of the summer school. With these considerations, 38 candidates from 25 countries (15 females and 23 males) were finally selected as trainees for the CLIVAR-FIO summer School in 2018.

All the trainees were required to give a presentation that covers their main research focus, their institutions' work areas and/or preliminary ideas for cooperation with other countries in the region. The trainee report contributes to establishing a research network and identifying possible areas of cooperation among the trainees. These reports were graded by all the participants including some of the invited lecturers, scores that were used to select participants that received the Best Trainee Award every year based on their performance.

Based on the scores received, we proceed to invite the participants receiving the highest scores to have their reports published in this special issue of Exchanges magazine, we want to acknowledge the trainees who accepted the challenge of working with the editorial team, to prepare the articles presented in this issue.

Introduction to the CLIVAR Summer School in China

Jing Li¹

1. International CLIVAR Project Office, Qingdao, China

1. Background

As a follow-up action after successfully hosting a CLIVAR Early Career Scientists Symposium in September 2016, which was held back to back with the CLIVAR Open Science Conference, and in line with the strong emphasis of CLIVAR on capacity building in particular geared towards Early Career Scientist, the First Institute of Oceanography (FIO) affiliated to the Minister of Natural Resources (MNR) has signed a 5-year agreement with the World Climate Research Programme (WCRP), agreeing to host a biannual 'CLIVAR Summer School' starting from 2018. The CLIVAR Summer School will be organised based on the existing resources of FIO, including but not limited to the scientific network of WCRP and CLIVAR, and the training platform of "UNESCO/IOC Regional Training and Research Center on Ocean Dynamics and Climate (UNESCO/IOC-ODC)", also hosted by FIO.

The objectives of the CLIVAR Summer School shall be to provide training and joint research opportunities for the next generation of climate and ocean scientists, while increasing the networking of early career scientists across different countries through:

- provision of a biannual training course in English, with recognized international experts invited as lecturers at appropriate times, to around 40-45 young scientists and doctoral /master students globally;
- organisation of training courses in English on related topics and promoting interactions between experienced oceanographers and students;
- fostering international joint research among lectures and trainees;
- exchange of visiting scholars and doctoral education.

2. The 2018 CLIVAR Summer School

2.1 Course Description

The first CLIVAR-FIO Summer School on "Past, Present and Future Sea Level Changes" was organized in Qingdao, China, from June 25 to 30, 2018, back to back with the eighth UNESCO/IOC-ODC Training Course on Ocean Forecast System, held during 2-7 July 2018. The objective of the 2018 CLIVAR Summer School was to provide early career scientists and engineers specializing in sea level research with an update in observations, knowledge, and understanding for the

study of global and regional sea level changes and their impacts in coastal areas. This Summer School covered a wide range of physical processes contributing to global and regional sea level change: from observations to modelling of the main physical processes of global and regional sea level rise and variability. In addition, there was a specific focus on impact studies in coastal areas.

A total of 38 trainees (15 females and 23 males) from 25 countries were selected out of the 202 applicants by considering the potential for future knowledge transfer, the education background, and the scientific expertise as important criteria, as well as the widest possible geographical distribution of the trainees, and a gender balanced distribution (Figure 1). During the summer course, 17 lectures were provided by eight experienced lecturers on sea level science from Germany, UK, France, Australia and China, who covered topics of observation and simulation of sea level changes, the contribution from thermal expansion, vertical land movement, glacier and ice sheet, the sea level projection in the past and in 2100 and beyond, as well as the coastal impacts of sea level rise and hazard management in coastal areas.

The school offered 42 hours of academic activities including keynote lectures, practical applications and field visits, spreading over 6 days (Figure 2). The Summer Course aimed to help establish research networks, and identify possible areas of cooperation among trainees and their institutions. To this end, each trainee was required to give a 10-minute presentation at the training course, which covered his/her main research focus, his/her institution's areas of work, and preliminary ideas for cooperation with other countries in the region. Trainee reports were scored by the instructors and participants, and the reports who obtained the highest scores were invited to submit an article to this special issue of CLIVAR Exchanges.

In order to encourage team work, all trainees were divided into groups (of 5-6 participants) with a coordinator elected by the members of each group with tasks to help organize their respective discussions and prepare group reports. The coordinator was required to chair one trainee report session. A group report was also required at the end of the training course, which summarizes the knowledge they have learnt from the

Figure 1: Geographical Distribution of participants accepted to attend the Summer Course

training courses, results of their assignments, and suggestions for cooperation in the future.

For evaluating the performance of trainees during the training course, a score was given to each trainee. The score was determined by attendance, motivation and activity performance during the training course and the quality of the trainee report and group report. A certificate of accomplishment was issued to the trainees who completed at least 85% of the course. Four trainees were selected as the best trainees according to their training scores.

A site visit was organised during one of the sessions of the summer school, to provide opportunities for both trainees and lecturers to visualize the China's progress on physical oceanographic researches, as well as the advancement of the high performance computing facilities (figure 3) and research vessels in FIO, the host of the summer course.

The CLIVAR Summer School also had the participation of supporting staff from the International CLIVAR Project Office (ICPO), the First Institute of Oceanography (FIO) and the UNESCO/IOC Regional Training and Research Center on Ocean Dynamics and Climate (ODC).

2.2 Reflection on the 2018 CLIVAR Summer School

Sea level research is a remarkable cross-disciplinary field, and demands an integral understanding of the global and regional ocean physical processes and their contributing factors. Modeling future sea level rise necessitates a working understanding of vertical land movement, ice sheet, glaciers and ocean dynamics, and observational and historical sea level records to put sea level oscillations in a paleoclimate context. Regional and global sea level research has huge societal and

economic implications for coastal communities, and should also be explicitly linked to coastal engineering and ocean forecasting research.

Several **significant outstanding questions** on sea level studies have been drawn from the course, including:

- One of the major challenges of sea level studies is the lack of high-resolution models and sea ice modelling;
- It is necessary to accurately estimate ocean heat content/thermal expansion;
- The need to understand the existence of potential positive feedback mechanisms in Antarctic melting;
- To determine threshold for melting of the Greenland Ice sheet
- To robustly attribute response to Green House Gases (GHG) and aerosol forcing, constrain projections
- To know how to better reconcile paleo sea level data with current climate models;
- To build a downscale model link to the contribution of hydrologists and other disciplines and decision-makers;
- The sea level rise demands to reduce the risk and to improve the resilience of coastal areas through identifying areas that are particularly vulnerable, monitoring for changes, and developing and implementing adaptation plans.

There were several **suggestions on future cooperation** coming out from group discussion during the course, including:

- To increase the knowledge and experience sharing on numerical modeling and data assimilation;

Figure 2: Trainees in class, group discussion

- To increase the knowledge and experience sharing on numerical modeling and data assimilation;
- To increase open access to code and data in published research;
- To increase communication channels among science disciplines;
- To increase the international network and collaboration amongst early career scientists in order to master and understand the current climate change constraints;
- To formulate better channels for communicating science to the general public;
- To develop earth science programs and resources targeted towards early career scientists in underrepresented countries, and countries that may experience dramatic climate effects.

2.3 Major outcomes

The CLIVAR-FIO Summer School has done an excellent job in drawing early career scientists from around the world and from a wide range of disciplines, including physical oceanography, atmospheric sciences, biogeochemistry, and coastal engineering, providing important opportunities to network with classmates from different scientific backgrounds. It also enabled the lecturers to know better about their audience, and make necessary adjustment about their lectures to be more easily digested by the trainees. Also, good interaction between lecturers and trainees was observed both in the class and during the breaks. According to the feedback from trainees, the content, quality and arrangement of the lectures are well designed and delivered. There were interrelations among lectures, and the key concepts and messages

on sea level changes have been reviewed and re-emphasized by several lecturers during their talks. However, some participants indicated that more practice should be expected for the training courses in the future. By the end of the training course, new ideas for future cooperation emerged from some of the groups, which are strongly encouraged by CLIVAR.

a. New ideas for future research and cooperation proposed by participants:

- Contribution of Sea Level Change and Interannual variability on Tropical Cyclones;
- To develop (or modify the existing) Ocean Forecasting System (OFS) covering the Bay of Bengal and South China Sea; and to enhancing the observation system in the region to increase the accuracy of the OFS;
- Close the surface wind fields on the ocean and run the ocean model to investigate the effects of wind on the western boundary currents in Western Pacific Ocean;
- To set up the wave model for Hawaii in addition to the general atmospheric model;
- Development or enhancement of ad hoc tools to produce a multi-hazards and risk assessment;
- Assessment of economic impacts of coastal flood risk and sea level rise scenarios;
- Preparation of a paper about long-term trends in sea level in each country of the group members, and their associated process that could explain this variation from a global and regional point of view.

Figure 3: Trainees and lecturers visiting FIO High Performance Computing Center

b. Enhanced international networking and interdisciplinary cooperation have been promoted among the trainees and/or with the lecturers;
c. Trainees will share information learned at the summer course with colleagues in home countries;
d. Due to the excellence performance during the summer school, one of the trainees, Kemgang Ghomsi Franck Eitel from Cameron, was nominated by one of the lecturers to join the CLIVAR Atlantic Region Panel, which has been approved by CLIVAR SSG.

- The use of social media and other communication tools will be further explored, in order to maintain the active involvement of trainees in CLIVAR activities in the future.

3. Follow up actions

CLIVAR would like to maintain the network formulated through the CLIVAR Summer School, and try to maximize the use of its platform and resources to provide opportunities for early career scientists to communicate with scientists in other countries. Several follow up actions have been taken, include:

- A special issue of CLIVAR Exchanges to publicize the high quality work of the trainees' is coordinated by ICPO;
- A special Channel called 'Spotlight for a Scientist' has been created on CLIVAR website (<http://www.clivar.org/ecs/spotlight-scientist>) to provide opportunities for Early Career Scientists to communicate with scientists worldwide through CLIVAR platform. This platform received positive feedback from Early Career Scientist, in particular from developing countries in Africa and South America. So far there are 16 profiles in total being published;
- ICPO consolidated the proposals from CLIVAR panels and research foci for the topic of the upcoming CLIVAR-FIO Summer School in 2020;

High-resolution projections of sea-level change processes on the Northwestern European Shelf

Tim Hermans^{1,3}, Aimée Slangen¹, Jonathan Tinker², Matthew Palmer², Bert Vermeersen^{1,3}

1. NIOZ Royal Netherlands Institute for Sea Research, Department of Estuarine & Delta Systems, and Utrecht University, PO Box 140, 4400 AC Yerseke, The Netherlands
2. Met Office Hadley Centre, FitzRoy Rd, Exeter, Devon EX1 3PB, United Kingdom
3. Faculty of Civil Technology and Geosciences, Geoscience and Remote Sensing, Delft University of Technology, Delft, The Netherlands

Introduction

Satellite altimetry shows that regionally sea-level change (SLC) can depart significantly from the global mean (Ducet et al., 2000). These regional deviations are the result of a number of different contributions, such as melt of land ice, terrestrial water storage, vertical land movement, changes in local sea water density and changes in the circulation of the oceans and atmosphere. Here, we focus on SLC due to changes in the circulation of the oceans and atmosphere. Due to salinity and temperature changes, the density of sea water changes and sea level will change as a result of expansion or contraction, also called steric SLC. Changes in water density and atmospheric forcing will also cause a different ocean circulation, leading to dynamic SLC. Patterns of steric and dynamic SLC can be simulated with process-based, global models of atmospheric and ocean circulation, for instance from the Coupled Model Intercomparison Project 5 (CMIP5) database (e.g. Slangen et al., 2012, 2014). These climate models simulate the response of ocean and atmospheric circulation under historical radiative forcing or under future scenarios, such as the Representative Concentration Pathways (RCPs; Meinshausen et al., 2011). However, the horizontal resolution of the ocean core of these CMIP5 models is often in the order of 100 by 100 km or more, which is too coarse to resolve the small-scale dynamics that matter on a regional to local scale.

Especially in areas such as the North Western European shelf (NWES), a high resolution is important to capture small scale (10-20 km) features in the bottom topography and circulation and possibly resolve the internal Rossby radius (2-5 km) on the shelf (Holt & Proctor, 2008), and to include tidal mixing processes. Since global climate models have a coarse horizontal resolution, do not include all processes important to shelf seas, and also often their vertical levels are limited in shallow regions, the resulting projections of sea-level change

might not be completely appropriate for areas such as the NWES (Ådlandsvik and Bentsen, 2007). In this study we investigate the importance of high-resolution simulations of SLC processes in the NWES area. We use high-resolution simulations for the period of 1980-2099 from dynamical downscaling experiments of the UK Met Office. Climate models of CMIP5 were dynamically downscaled with the limited-area, regional shelf seas model NEMO AMM7 (O'Dea et al., 2017), which has a nominal horizontal resolution of 7 by 7 km and terrain-following coordinates. The trends and variability of sea level simulated by the high-resolution downscaling experiments will be compared with the corresponding global climate model results and with observations. It will also be investigated which processes in the North Western European shelf seas are the main drivers behind the differences between the projections of the global and downscaled models. Preliminary results show that the high-resolution projections provide enhanced regional detail with respect to the low-resolution projections and that sub-annual variability is better represented.

Methodology

Simulations from 1980 to 2099 of sea level and several other parameters from two global climate models are dynamically downscaled using the regional shelf seas model AMM7 CO6. AMM7 CO6 is a shelf seas configuration of the primitive-equation Nucleus for European Modelling of the Ocean (NEMO). Previous versions AMM CO4 and AMM CO5 have been described extensively and were validated against historical data (O'Dea et al., 2012, 2017). The AMM7 domain extends from 20°E to 13°W, and from 40°N to 65°N, and comprises the eastern North Atlantic Ocean and several marginal seas (see Figure 1).

The regional shelf seas model is nested in two different global climate models from the CMIP5 database, HadGEM2-ES (Collins et al., 2011) and MPI-ESM-LR

Figure 1: Shallow shelf seas in the AMM7 domain and the NOOS bathymetry on the AMM7 domain; image obtained from O'Dea et al. (2017).

(Giorgetta et al., 2013) and driven by lateral ocean boundary conditions and atmospheric surface forcing derived from these global models (see Figure 2 for a schematic overview). The downscaling set-up is similar to that of Tinker et al. (2015), who showed that their modelling system is appropriate for providing centennial climate projections. Via lateral boundary conditions, temperature, salinity, sea surface height and barotropic currents are prescribed to the AMM7 domain (O'Dea et al., 2017). The atmosphere of the climate models over the NWES is dynamically downscaled using the regional atmospheric model RCA4 of the Rossby Centre prior to applying the atmospheric surface forcing. River run-off is simulated with the Total Runoff Integrating Pathways (TRIP) river routing model. Recognizing the importance of tides and surges on the NWES, the model has a non-linear free-surface and a time splitting approach for baroclinic and barotropic processes (O'Dea et al., 2017).

Figure 2: Schematic overview of dynamical downscaling set-up. The regional shelf seas model AMM7 is forced by lateral oceanic boundary conditions, surface atmospheric forcing and river runoff.

Preliminary results

We now show some preliminary results of sea-level trends and sub-annual sea-level variability. In Figure 3, linear sea-level trend anomalies for the 'future' period of 2006-2099 are shown for the global and downscaled climate models. Compared to the global climate models (Figures 3a and 3c), the downscaled simulations show enhanced detail (Figures 3b and 3c). The differences in the spatial patterns of the trends between the global and downscaled HadGEM2-ES models are larger than that for MPI-ESM-LR, which is most likely due to the fact that the spatial resolution of the global MPI-ESM-LR model (orthogonal curvilinear coordinates; approximately $3/4^\circ$ by $1/2^\circ$ in the North Sea) is closer to that of the regional model than the resolution of HadGEM2-ES (1° by 1°). In the North Sea, the trend anomalies obtained from the HadGEM2-ES global model (see Figure 3a) are over 1 mm/yr larger than the trends on the shelf west of the United Kingdom. This contrast is not observed in the trends of the regional model (see Figure 3b). This could be the result of the closed English Channel in HadGEM2-ES, which might lead to water piling up in the southern North Sea, but this warrants further investigation.

In Figure 4, we show a comparison of the sub-annual variability in the global and downscaled simulations of one of the models (HadGEM2-ES) from 1980 to 2016. Additionally, these are compared to the sub-annual variability of tide gauge records from the Permanent Service for Mean Sea Level (Holgate et al., 2013; PSMSL, 2018). The sub-annual variability is computed as the standard deviation of the de-trended time series of monthly mean sea level, in which the inverse barometer effect in the models was accounted for as well. Gaps of one month in the tide gauge data were filled by linear interpolation.

We find that the sub-annual variability of the global HadGEM2-ES model matches that of the downscaled results in a general sense (see Figure 4). However, at several tide gauge locations the downscaled results are closer to the observations than the results of the global climate model. This is especially the case along the coastlines of the southern part of the North Sea and southwest of the United Kingdom, where the variability simulated by the global model differs around 2 cm with that of the tide gauge records.

Figure 3: Projections of linear sea level trends for the period 2006-2099 (high-end scenario RCP8.5) for the a) global HadGEM2-ES and b) dynamically downscaled HadGEM2-ES models, and for the c) global MPI-ESM-LR and d) dynamically downscaled MPI-ESM-LR models. The trends are given with respect to the spatial mean trend of the NWES. Global climate model output was bilinearly interpolated onto the NEMO AMM7 grid, and the original land mask (red line) was filled with nearest-neighbor extrapolation.

Outlook

The preliminary results show that especially for HadGEM2-ES there can be significant differences (> 1 mm/yr) between projected sea-level trends in the global and downscaled models, for example in the southern North Sea, near the Norwegian Trench and at the Iceland-Faroe Ridge. Additionally, the sub-annual variability is closer to observed variability in the downscaled models, implying a more realistic atmospheric forcing. In the following steps, we will further investigate the sea-level trends and variability (also on longer time scales than sub-annual) and validate results against observations for the historical period. Additionally, we will investigate the drivers of differences in trends and variability, including the analysis of temperature and salinity profiles, investigating the changes in atmospheric forcing, and decomposing SLC into contributions due to steric changes and changes in bottom pressure (Landerer et al., 2006). Currently, relatively coarse-resolution (~ 100 km) global models are used for centennial projections of changes in steric and dynamic sea level. With the results of our study, we can quantify the uncertainties in projections of SLC due to the coarse resolution of global climate models and the parameterization and/or omission of small-scale processes that are important to shallow shelf sea regions. This will help to improve SLC projections in these regions.

Figure 4: Sub-annual variability of sea level from 1980-2016 for a) the global HadGEM2-ES model and b) the dynamically downscaled HadGEM2-ES model. The circles and their colour show the variability in tide gauge records over the same time period.

References

- Ådlandsvik, B., & M. Bentsen, 2007. Downscaling a twentieth century global climate simulation to the North Sea. *Ocean Dynamics*, (M), 453–466. <https://doi.org/10.1007/s10236-007-0125-2>
- Collins, W. J., Bellouin, N., Doutriaux-Boucher, M., Gedney, N., Halloran, P., Hinton, T., Hughes, J., Jones, C. D., Joshi, M., Liddicoat, S., Martin, G., O'Connor, F., Rae, J., Senior, C., Sitch, S., Totterdell, I., Wiltshire, A., and Woodward, S., 2011. Development and evaluation of an Earth-System model – HadGEM2. *Geosci. Model Dev.*, 4, 1051-1075, <https://doi.org/10.5194/gmd-4-1051-2011>, 2011.
- Ducet, N., P. Y. Le Traon,, & G. Reverdin, 2000. Global high-resolution mapping of ocean circulation. *Journal of Geophysical Research*, 105(C8), 19477–19498.
- Giorgetta, M. A., et al., 2013, Climate and carbon cycle changes from 1850 to 2100 in MPI-ESM simulations for the Coupled Model Intercomparison Project phase 5. *J. Adv. Model. Earth Syst.*, 5, 572–597. doi: 10.1002/jame.20038.
- Holgate, S. J., A. Matthews,, P. L. Woodworth, L. J. Rickards, M. E. Tamisiea, E. Bradshaw, P. R. Foden, K. M. Gordon, S. Jevrejeva, & J. Pugh, 2013. New Data Systems and Product at the Permanent Service for Mean Sea Level. *Journal of Coastal Research*, 29, 3, 493-504. doi:10.2112/JCOASTRES-D-12-00175.1.
- Holt, J., and R. Proctor, 2008. The seasonal circulation and volume transport on the northwest European continental shelf: A fine-resolution model study, *J. Geophys. Res.*, 113, C06021. doi:10.1029/2006JC004034
- Landerer, F. W., J. H. Jungclaus, , & J. Marotzke, 2006. Regional Dynamic and Steric Sea Level Change in Response to the IPCC-A1B Scenario. *Journal of Physical Oceanography*, 37, 296–312. <https://doi.org/10.1175/JPO3013.1>
- Meinshausen, M., S. J. Smith, K. V. Calvin, J. S. Daniel, M. L. T. Kainuma, J.-F. Lamarque, K. Matsumoto, S. A. Montzka, S. C. B. Raper, K. Riahi, A. M. Thomson, G. J. M. Velders and D. van Vuuren, 2011. The RCP Greenhouse Gas Concentrations and their Extension from 1765 to 2300. *Climatic Change* (Special Issue), doi:10.1007/s10584-011-0156-z
- O'Dea, E., A. K. Arnold, K. P. Edwards , R. Furner, P. Hyder, , M. J. Martin, ...H. Liu, 2012. An operational ocean forecast system incorporating NEMO and SST data assimilation for the tidally driven European North-West shelf. *Journal of Operational Oceanography*, 5(1), 3–17.
- O'Dea, E., R. Furner, S. Wakelin , J. Siddorn, J. While, P. Sykes, ...H. Hewitt, 2017. The CO5 configuration of the 7 km Atlantic Margin Model: large-scale biases and sensitivity to forcing , physics options and vertical resolution. *Geoscientific Model Development*, 10, 2947–2969.
- Permanent Service for Mean Sea Level (PSMSL), 2018, "Tide Gauge Data", Retrieved 29 March 2018 from <http://www.psmsl.org/data/obtaining/>.
- Slangen, A. B. A., C. A. Katsman , R. S. W. Van de Wal, L. L. A. Vermeersen, & R. E. M. Riva, 2012. Towards regional projections of twenty-first century sea-level change based on IPCC SRES scenarios. *Climate Dynamics*, 38(5–6), 1191–1209. <https://doi.org/10.1007/s00382-011-1057-6>
- Slangen, A. B. A., M. Carson, C. A. Katsman, R. S. W., van de Wal, A. Kohl, L. L. A. Vermeersen, & D. Stammer, 2014, Projecting twenty-first century regional sea-level change. *Climate Change*, 124, 317-332. doi:10.1007/s10584-014-1080-9.
- Tinker, J., J. Lowe, J. Holt , A. Pardaens, & A. Wiltshire, 2015. Progress in Oceanography Validation of an ensemble modelling system for climate projections for the northwest European shelf seas. *Progress in Oceanography*, 138, 211–237. <https://doi.org/10.1016/j.pocean.2015.07.002>

Understanding the Recent Hawaii High Sea Level Event

Xiaoyu Long¹, Matthew Widlansky¹

1. University of Hawaii, Sea Level Center

Introduction

During the years of 2016 and 2017, the Hawaii islands experienced record-breaking high sea level across the island chain (Fig.1), which caused nuisance flooding in vulnerable coastal communities and exasperated beach erosion, especially around the times of highest tides. Near-real-time analyses of tide gauge observations and satellite-based altimetry measurements of sea surface heights around Hawaii allow the non-tidal above normal sea levels to be broken down into four primary contributions: (1) long-term sea level rise, (2) oceanic planetary waves, (3) oceanic meso-scale eddies, and (4) an inverse-barometer effect of low atmospheric pressure on the ocean. Long term sea level rise is unavoidable, and under which the occurrence of high sea level events will become more extreme and have longer duration. Oceanic meso-scale eddies are known to have impact on Hawaii sea level (Firing and Merrifield, 2004). However, it is still challenging to fully resolve these eddies using current generation of ocean model. Inverse barometer effect generally plays a small role in Hawaii latitude. In this study, we will focus on the large scale oceanic dynamics, by using both observation and ocean general circulation model (OGCM), which are the oceanic planetary waves that contribute significantly to the high sea level events during last two years.

Data and model setup

In this study, we use ECMWF ocean reanalysis version 4 (ORAS4), satellite altimetry data from the SSALTO/DUACS multi-mission dataset and the tide gauges data archived in University of Hawaii Sea Level Center. ORAS4 assimilates satellite altimetry in its assimilation system and thus we were able to resemble the observed sea level height better than other reanalysis dataset.

For the model experiments, we use CESM pop2, which is originally developed at LANL and is the ocean component of NCAR earth system model CESM. The ocean model is forced by atmospheric data from JRA55, the Japanese 55-year Reanalysis. The model is configured nominally in 1-degree resolution. The model is spun-up with climatological forcing for 50 years, then forced with real-time forcing from 1979 to 2017. A few experiments are designed to isolated possible mechanisms that are responsible for the high sea level event, and the experiment details are

introduced in the following sections.

Results

Previous studies identified possible relation between high sea level events in Hawaii and ENSO events (Yvonne et al., 2004), though they did not discuss the underline dynamical mechanism. And also the relationship between Hawaii high sea level and ENSO is fragile. For example, the 1997 ENSO has comparable strength as the 2015 ENSO event, but no extreme high sea level was observed in Hawaii (Fig.1). The zonal propagation of the signal proposes the possible role of oceanic Rossby wave (Fig.2). The coast trapped Kelvin wave signals have similar strength in 1997 and 2015. However, the downstream propagation of the signal is so different in 1997 and 2015, and the propagation of the high sea level signal is the cause of the long duration of high sea level observed in Hawaii during 2016-2017. The propagation of high sea level may be explained by long Rossby wave, and the wave source can be either boundary signal (coastal trapped kelvin wave) or the wind forced. Two experiments are conducted to isolate the different source of Rossby wave. The subtropic experiment uses full sub-tropic winds to force the ocean while keeping winds in other region as climatology, and this experiment only include the role of wind forcing in the basin. The tropic experiment only uses full winds in the Tropics, which can reproduce the ENSO and coast trapped Kelvin wave at the eastern boundary. The results (not shown) indicate that most of the variability of Sea surface height (SSH) in Hawaii is contributed by the wind forcing in the basin while the boundary signal only contributes about 10% of the variance.

A natural question arises: why the SSH anomaly is so different in 1998-1999 compared to that in 2016-2017. We integrate the 1.5 layer long Rossby wave model (Qiu, 2002) with reanalysis atmospheric forcing. The 2016-2017 high sea level is well reproduced, while the model produces a similar SSH response as in 2016-2017, which did not happen in the real world. The 1.5 layer model represents nothing but the potential vorticity (PV) conservation of the mode-1 baroclinic Rossby wave. Other factors that might be important to SSH are absent in this simple model. In order to understand the different SSH responses to ENSO,

Figure 1: SSH anomaly at (a) Hawaii and (b) Santa Cruz. For Hawaii, the gauge data (blue) is average 5 stations on different islands, and the ORAS4 and satellite altimetry (red) is averaged over a 5x6 degree box around Hawaii. The Santa Cruz tide gauge (0.7S 90.3W) here is used as an indicator of ENSO signal. The green curve in (a) is the 11-year low-pass filtered time series of blue curve. The anomaly is calculated by subtracting the climatology at each location. The group photo of 2016 cruise.

we conduct another two experiments which make the 2016-2017 case starts like (exp3) or ends like (exp4) 1998-1999. In exp3, the winds in 2013-2014 are replaced by winds in 1995-1996, and in exp4, the winds in 2016-2017 are replaced with winds in 1998-1999.

Figure 2: Hovmöller diagram of SSH anomaly (shading, units in meter) and Ekman pumping (hatching) average over 14-23N from (a) Year 1996 to 1999 and (b) Year 2014 to 2017. Green hatching is Ekman suction and black is Ekman pumping, only contours of $\pm 2e-7$ m/s are plotted.

As is shown in Fig.4, letting the ENSO 2015 starts like ENSO 1997 does not noticeably change the SSH response, which is consistent with the results that the wind forcing in the basin is more important. In contrast, the exp4 kills the high sea level events in the control run. The 1997 ENSO quickly switched to La Niña condition, and in the wind field, the prevailing westerly during ENSO also switched to easterly (Fig.3).

Figure 3: SSH (shading), Ekman pumping velocity (contour) and surface wind stress (vector) averaged over the two years period indicated in the plot.

Figure 4: 4 SSH anomaly averaged between 14-23N for the three OGCM experiments. Left: control simulation forced with real atmospheric flux; Middle: the atmospheric forcing during 2013-2014 is replaced by 1995-1996; Right: the atmospheric forcing during 2016-2017 is replaced by 1998-1999.

4 Future work

Here we only present some preliminary results about the cause of 2016-2017 Hawaii high sea level, and why it is different from the SSH response in 1998-1999. During these two ENSO years and after, the SSH anomaly is well correlated with steric height. Steric height variability is usually controlled by the temperature variation in the first few hundred meters. Our future work is to understand the cause of the different temperature anomaly in 1998-1999 and 2016-2017. One possible reason is the enhanced evaporation cooling in the La Niña condition in 1999. One model experiment is ongoing, in which we prescribe the sensible heat flux and evaporation flux in 1998-1999 the same as in 2016-2017, while the momentum flux (Ekman pumping) is calculated automatically in the model. If the surface cooling is the reason of the negative SSH anomaly, then this experiment should generate SSH anomaly similar to 2016-2017.

References

Magdalena Alonso Balmaseda, Kristian Mogensen, and Anthony T Weaver. Evaluation of the ECMWF ocean reanalysis system ORAS4. *Quarterly Journal of the Royal Meteorological Society*, 139(674):1132–1161, 2013.

Firing and Merrifield. Extreme sea level events at Hawaii: Influence of mesoscale eddies. *Geophysical Research Letters*, 31(24), dec 2004.

Yvonne L Firing, Mark A Merrifield, Thomas A Schroeder, and Bo Qiu. Interdecadal Sea Level Fluctuations at Hawaii. *Journal of Physical Oceanography*, 34(11):2514–2524, nov 2004.

Bo Qiu. Large-Scale Variability in the Midlatitude Subtropical and Sub- polar North Pacific Ocean: Observations and Causes. *Journal of Physical Oceanography*, 32(1):353–375, jan 2002.

Assessment of the intraseasonal Kelvin wave activity during Central Pacific and Eastern Pacific El Niño events

Jonathan Cedeño^{1,3,*}, Óscar Pizarro^{1,2,3}, Aldo Montecinos^{2,3}

1. Departamento de Oceanografía, Universidad de Concepción, Concepción
2. Departamento de Geofísica, Universidad de Concepción, Concepción
3. Instituto Milenio de Oceanografía, Concepción

Introduction

The intraseasonal oceanic oscillations in the tropical Pacific are mainly driven by equatorial Kelvin waves (EKW) forced by the winds in the western and central equatorial Pacific (Enfield, 1987). They peak at periods near 60-75 days, and are tied to intraseasonal atmospheric waves known as Madden Julian Oscillation (MJO) (Kessler et al., 1995). The equatorial Pacific also influences the intraseasonal variability in the west coast of South America through two processes: poleward propagation of coastal-trapped waves (CTW) excited by EKW (Shaffer et al., 1997), and local wind forced oscillations associated with tropical Pacific warming via atmosphere teleconnections (Hormazábal et al., 2002).

El Niño-Southern Oscillation (ENSO) is closely related to MJO. It modulates the signal of MJO (Kessler et al., 1995), and in turn this intraseasonal oscillation plays a key role on the development of ENSO events, either as a trigger mechanism or as a medium to sustain El Niño (Johnson and McPhaden, 1993).

Additionally, there is evidence of some interdecadal fluctuations that affects El Niño, determining its flavour (Choi et al., 2011). Yu and Kao (2007) and Ashok et al. (2007) are the first to describe El Niño types. They found one mode of variability on sea surface temperature (SST) anomalies, with a maximum amplitude in the central Pacific, calling it “date line”, “Modoki”, or “central Pacific” (CP) El Niño. By contrast, “canonical” El Niño events (Rasmusson and Carpenter, 1982) develop larger SST anomalies near the west coast of South America during the peak phase. This mode is also known as “eastern Pacific” (EP) El Niño.

Several studies have analyzed the characteristics of these events from the perspective of Kelvin wave dynamics. Using different model outputs, Dewitte et al. (2012a, b) found that Kelvin waves play different roles during the evolution of El Niño. On CP events, there is a decoupling between baroclinic modes, being the first mode participate in the (heat content) recharge process; meanwhile the second mode is more associated to the ocean adjustment mode of the delayed oscillator mechanism (Jin, 1997), participate on the discharge

process. On EP events, both modes have a comparable role in the recharge-discharge process of El Niño. In the intraseasonal band, dissipative processes seem to inhibit propagation of Kelvin waves in the eastern Pacific at CP El Niño events (Mosquera-Vásquez et al., 2014). It has been associated with the reinforcement of the zonal gradient in equatorial stratification, leading scattering of energy and partial reflection of Kelvin waves through intraseasonal Rossby waves.

In this work, the equatorial Kelvin wave activity is studied through the intraseasonal variance of sea level due to the relevance between these two processes on the Pacific region (Luther, 1980). In addition, based on the definition of El Niño years from Yu et al. (2012), composites for EP and CP El Niño events were prepared to assess the differences between different types of events.

Data and Methods

Sea level from three stations along the equatorial Pacific: Christmas (157°W), Santa Cruz (90°W) and La Libertad (81°W), were used in this study. The data were downloaded from the University of Hawaii Sea Level Center (UHSLC). The intraseasonal variance was obtained firstly by band-pass filtering using a cosine-Lanczos filter with half amplitudes at 30 and 120 days, and then by estimating the mean variance on 120-days consecutive segments.

Zonal changes in the sea level intraseasonal variance along the equatorial Pacific

Fig. 1 shows the intraseasonal variance of sea level from the three stations. It was observed from the longest record (La Libertad) that during 1950-1979, the variance was typically 5 cm², with relatively small peaks. In contrast, the next 34 years were characterized by large peaks in 1982-83, 1986-87 and 1997-98. All these periods correspond to El Niño years, with particularly large events in 1982-83 and 1997-98. The change in mean variance before and after 1980 at La Libertad could be related to the PDO phase shift of 1977 (Mantua et al., 1997).

Figure 1: Intraseasonal variance of sea level for equatorial stations. The values in parentheses show the mean variance over the corresponding period.

The variance average at Santa Cruz during the 1980-2012 period was 4 cm^2 , being somewhat lower than La Libertad and Christmas (8 cm^2). Yoon (1981) found that the effects of the Galápagos islands in the energy propagation and amplitude of Kelvin waves is negligible due to the relatively smaller meridional scale ($\sim 100 \text{ km}$) of the archipelago, compared with the typical baroclinic mode 1 internal deformation radius ($\sim 485 \text{ km}$) of the wave. For the 1997-98 El Niño event, the variance was comparable between Christmas and Santa Cruz, but far from the peak of almost 50 cm^2 at La Libertad. This pulse, occurred around May 1998, and was not observed at Santa Cruz sea level. On the contrary, it raised the sea level and slightly deepened the thermocline at the Ecuadorian coast (Zambrano et al., 2000).

The sea level variance composites for EP and CP El Niño events are shown on Fig. 2 (left panels). Generally, they describe a two-peak pattern. The first is centered in autumn (0) (summer (0)) on EP (CP) events. The second peak occurs in summer (+1) on both events, persisting by a couple of months.

The differences in variance levels between composites are evident, especially over the second peak. In Christmas, the CP event was more energetic than EP by a factor of 1.6. By contrast, on eastern Pacific stations, the EP composite had a larger variance than CP. The difference reach $+10 \text{ cm}^2$ in Santa Cruz, but is greatest on La Libertad ($+15 \text{ cm}^2$). Also, at this station, the second peak is wider than that at Christmas and Santa Cruz.

We first tested the differences between the EP composite (including 1982-83 and 1997-98 events) and the CP composite (Fig. 2, right panels). Because of these two extreme El Niño events, statistically significant differences were observed during peak and decay phases

of El Niño at La Libertad and Santa Cruz stations, peaking on Apr. (+1). At Christmas, the statistically significant differences were limited to a few days of Jun. (0).

In order to unmask a possible bias introduced by the great variance of the 1982-83 and 1997-98 El Niño events, we also analyzed EP composites excluding those two events. The results (not shown) indicate that the differences were still significant between CP and EP El Niño years at La Libertad and Santa Cruz stations, on summer (+1) peak.

It is worth noting that the intraseasonal variance on CP El Niño events experience a decrease from central to eastern Pacific, which is more evident in the second peak of El Niño (Fig. 2a). These findings evidence the Kelvin wave dissipation process on CP El Niño events, previously documented by Mosquera-Vásquez et al. (2014) based on observed and modeled subsurface data.

Discussion and conclusion

Herein, we studied the intraseasonal Kelvin wave activity during El Niño traced by their surface manifestation on sea level. Our results suggest some distinctive characteristic patterns in which Kelvin waves are present in the equatorial Pacific during both EP and CP El Niño events. The intraseasonal Kelvin wave activity peaks in austral summer, when generally El Niño reach their maxima amplitude. Its variance is higher during EP rather than CP events, and their differences are statistically significant in the peak of the event. The decrease in EKW energy levels in CP events is tied with an observed attenuation of this signal on sea level height altimetry, from central to eastern equatorial Pacific.

During the onset phase of El Niño EP, the variance levels in the equatorial stations were slightly higher than in CP years. Nevertheless, their differences are not statistically significant (90% level). This increased variability of the sea level during the onset could be a characteristic of some extreme events (i.e. 1997-98 El Niño), a closer examination of the data suggests this, but definitively not of the CP events.

The low intraseasonal activity of sea level during CP years observed in the eastern equatorial Pacific seems to be determined more by scattering (due to a strong thermocline slope) rather than by the wind, despite the changes in its spatial pattern. In an extension of this study, Cedeño (2015), using a simple equatorial Kelvin wave model, shows that the intraseasonal variance of modeled sea level for both types of El Niño events does not show significant differences in the eastern Pacific during the onset, peak and decaying phases (meaning that, despite the wind forcing characteristics of each type of El Niño events, the sea level response is similar in the eastern Pacific). This is consistent with the description of Dewitte et al. (2012a), who used data from a coupled global circulation model. These authors found that during CP years, the Kelvin wave amplitude responds in a lesser extent to the forcing characteristics of zonal wind stress, and more to a condition of equatorial thermocline that promotes a wave dispersion in the eastern Pacific.

Figure 2: EP and CP El Niño composites of sea level intraseasonal variance for equatorial stations (a) and their differences (b). EP composite included the extraordinary 1982-83 and 1997-98 El Niño events. The organization and time notation of the composite follows the work of Larkin and Harrison (2002). The central year for which the event takes place is designed as year “0”, while “-1” and “+1” years refer to the annual periods that precede and succeed the central event, respectively. The red dots represent the segments which were significant at $\alpha = 0.10$ using a Monte Carlo significance test.

What controls the thermocline depth in the eastern Pacific during CP events? Zhang and McPhaden (2006) proposed that the local variations of zonal wind stress in the eastern equatorial Pacific could work together or against with the remote forcing to determine the thermocline depth at this region, relativizing the well-known ENSO mechanism of thermocline feedback. Particularly, they referred that during 2002-03 El Niño (CP) event, observed easterly winds were unusually high in this basin, producing both a shallower thermocline and intensification of local upwelling. In our study, nevertheless, the variance composite of zonal wind stress does not show a maximum of intraseasonal activity in the eastern Pacific during CP years. It is clear that a more detailed analysis of this variable is required, as well as the quantification of their effect in the variations of the local thermocline that determine, ultimately, the attenuation degree of Kelvin waves in central Pacific El Niño events.

Acknowledgment

This research was funded by the Ecuadorian state through the Secretaría Nacional de Educación Superior, Ciencia, Tecnología e Innovación (SENESCYT) scholarship (052-ARG-2013) granted to JC. Complementary funds were obtained from Instituto Milenio de Oceanografía (IMO), Chile (Ministerio de Economía/Iniciativa Científica Milenio 2012 No.

IC120019). OP and AM also thank to FONDECYT (Fondo Nacional de Desarrollo Científico y Tecnológico de Chile) projects No. 1121041 and 1140845.

* Now at Facultad de Ingeniería Marítima, y Ciencias del Mar (FIMCM), Escuela Superior Politécnica del Litoral (ESPOL).

References

Ashok K., S.K. Behera, S.A. Rao, H. Weng, and T. Yamagata (2007). El Niño Modoki and its possible teleconnection. *J. Geophys. Res.* 112(C11007), 1:27.

Cedeño J. (2015). Interannual Variability of Intraseasonal Kelvin Waves in the Eastern Equatorial Pacific. M.Sc. Thesis. Oceanography program. Universidad de Concepción. Concepción.

Choi J., S.-I. An, J.-S. Kug and S.-W. Yeh (2011). The role of mean state on changes in El Niño’s flavor. *Clim. Dyn.*, 37, 1205–1215.

Dewitte B., J. Vásquez-Cuervo, K. Goubanova, S. Illig, K. Takahashi, G. Cambon, S. Purca, D. Correa, D. Gutierrez, A. Sifeddine and L. Ortlieb (2012a). Change in El Niño flavours over 1958-2008: implications for the long-term trend of the upwelling off Peru. *Deep-Sea Res. II* 77-80, 143:156.

Dewitte B., J. Choi, S.-L. An, and S. Thual (2012b). Vertical structure variability and equatorial waves during Central Pacific and Eastern Pacific El Niño in a coupled general circulation model. *Clim. Dyn.* 38, 2275:2289.

Enfield D.B. (1987). The Intraseasonal oscillation in Eastern Pacific Sea Level: how is it forced? *J. Phys. Oceanogr.* 17, 1860:1876.

Hormazábal, S., G. Shaffer, and O. Pizarro (2002). Tropical Pacific control of intraseasonal oscillations off Chile by way of oceanic and atmospheric pathways, *Geophys. Res. Lett.*, 29 (6).

Jin F.-F. (1997). An equatorial ocean recharge paradigm for ENSO. Part I: Conceptual model. *J. Atmos. Sci.* 54, 811:829.

Johnson, E.S., and M.J. McPhaden (1993). Structure of intraseasonal Kelvin Waves in the Equatorial Pacific. *J. Phys. Oceanogr.* 23, 608:625.

Kessler, W., M. McPhaden, and K. Weickmann (1995). Forcing of intraseasonal Kelvin waves in the equatorial Pacific, *J. Geophys. Res.*, 100, 10,613–10,631.

Mantua, N., S. Hare, Y. Zhang, J. Wallace, and R. Francis (1997). A Pacific interdecadal climate oscillation with impact on salmon production, *Bull. Amer. Meteor. Soc.*, 78 (6), 1069–1079.

Mosquera-Vásquez K., B. Dewitte and S. Illig (2014). The Central Pacific El Niño intraseasonal Kelvin wave. *J.*

Geophys. Res. Oceans 119, 6605–6621.

Larkin, N., and D. Harrison (2002), ENSO warm (El Niño) and cold (La Niña) event life cycles: ocean surface anomaly patterns, their symmetries, asymmetries, and implications, *J. Climate*, 15 (10), 1118–1140.

Luther D.S. (1980). Observations of long period waves in the tropical oceans and atmosphere. Ph.D. thesis. MIT-Woods Hole Oceanographic Institution, Joint Program in Oceanography. Woods Hole, MA.

Rasmusson, E.M. and T.H. Carpenter (1982). Variations in Tropical Sea Surface Temperature and Surface Wind Fields Associated with the Southern Oscillation/El Niño. *Mon. Wea. Rev.* 110, 354–384.

Shaffer G., O. Pizarro, L. Djurfeldt, S. Salinas and J. Rutllant (1997). Circulation and Low-Frequency Variability near the Chilean Coast: Remotely Forced Fluctuations during the 1991–92 El Niño. *J. Phys. Oceanogr.*, 27, 217–235.

Yoon, J.-H. (1981), Effects of islands on equatorial waves, *J. Geophys. Res.*, 86 (C11), 10,913–10,920.

Yu J.-Y. and H.-Y. Kao (2007). Decadal changes of ENSO persistence barrier in SST and ocean heat content indices: 1958–2001. *J. Geophys. Res.* 112, D13106.

Yu J.-Y., Y. Zou, S.T. Kim and T. Lee (2012). The changing impact of El Niño on US Winter temperatures. *Geophys. Res. Lett.* 39, L15702.

Zambrano, E., R. Martínez, and J. Garcés-Vargas (2000), Variabilidad oceanográfica en Ecuador asociada con el evento ENSO 1997–98, *Acta Oceanogr. Pac.*, 10 (1), 9–20.

Zhang X. and M.J. McPhaden (2006). Wind stress variations and interannual sea surface temperature anomalies in the eastern equatorial Pacific. *J. Climate*, 19, 226–241.

The Nigerian's Gulf of Guinea coastal evolution and response to multi-scale forcing

Dada Olusegun Adekoyejo¹

1. Department of Marine Science and Technology, Federal University of Technology, Akure, Nigeria

Abstract

The ocean warming and sea-level rise will significantly impact coastal zones, displace ecosystems, alter geomorphological configurations and their related sediment dynamics, and increase the vulnerability of man-made infrastructure; they are also expected to intensify the coastal flooding occurrence on low-lying coasts in the next few decades. The West African coast, a low-lying coastal area, has been declared as one of the most vulnerable areas to experience annual or more frequent flooding, implying an increase in flood frequency that will require improved protection and migration in some extreme cases. This challenge requires the improvement of understanding on the possible trends as well as contributions of different forcing to the local coastal sea-level change and subsidence at various time-scales and locations. These aforementioned situations formed the fulcrum of my research interest, that is, coastline change, sea level change, coastal flooding, coastal subsidence and other coastal processes along the Nigerian coast.

of global climate change, especially acceleration in sea-level rise, further rise in sea temperature, more extreme weather events and storms surges, and altered precipitation and ocean acidification; and in addition the human intervention (Vitousek et al., 2017; Borman et al., 2016; Church et al., 2011; Nicholls and Cazenave, 2010). For coastal management, it implies that low-lying areas could potentially be subject to increased erosion and flooding in the future. A predicted increase in frequency and magnitude of storms would add the issue of sea level rise, which not only has an impact on coastal morphodynamics but affects the economy and population of the low-lying coastal countries.

Research Region

This research focuses on the Nigerian coast of the Gulf of Guinea (GoG), North Atlantic, where I have been carrying out research for the past 5 years. The Nigerian sector is the most fascinating sector of the GoG coast for littoral research. This sector is truly superlative because (i) it houses the world's third largest delta, (ii) it is one of the world's most prolific hydrocarbon (oil and gas) provinces on which the economy of the region solely depends, and (iii) it's morphologically defined by barrier-lagoon, transgressive mud, arcuate delta and flanks and strand coasts (Fig. 1). Due to the low elevation and relief of Nigerian coast, a large number of coastal dwellers and multi-billion dollars economic infrastructures situated in this region are tend to subject to coastal margin subsidence and rising sea levels, which may lead to seawater incursions and inundation of sensitive areas. The combined eustatic sea-level rise and subsidence will have a direct negative impact on more than 40 percent of the Nigerian population and could also have severe consequences for Nigeria's entire economy. In spite of this uniqueness, it is faced with many environmental challenges and the spatial and temporal variability of the governing coastal processes and their implications in this sector remains poorly studied.

Previous Research

My previous research along the Nigerian coast focused on understanding shoreline evolution, on long-term (century) and short-term (seasonal) scales and the dominant processes responsible for these changes on different timescales (Dada et al., 2015; Dada et al.,

Figure 1: Map showing different sections of the Nigerian coast.

Research Background

Coastal zones are the interface that occurs between atmospheric, oceanic and land systems. Aside from being key geomorphological archives, a large proportion of the world's population, critical economic infrastructures and diverse ecosystems are found in these regions. However, they face an uncertain future, because they are highly vulnerable to the effects

2016a, b; Dada et al., 2018; Dada et al., 2019; Dada et al., under review).

Current Research

My current research focuses on investigating the recent trends in coastal sea level variations and coastal flooding, as well as land subsidence in the Nigerian Transgressive Mud coast, where accelerated coastline retrogradation (as high as -39 m/yr; Figs 2-3) and coastal flooding (Fig. 4) are taking place. However, the mechanism causing accelerated coastal retrogradation in this region remains unidentified (Dada et al., 2019; Ebisemiju, 1987). Hence the objectives of my current research are to:

- 1. investigate the recent trends in coastal sea level and its relative contributions to the coastal flooding events on the low-lying Nigerian Transgressive Mud coast from altimetry-derived sea level, tides and wave run-up intra- and inter-annual variability.
- 2. investigate the recent status and future potential of land subsidence in the Nigerian sections of the Transgressive Mud coast and equally the Niger Delta, unravel and quantify the different drivers of subsidence (whether natural or anthropogenic processes), and to understand how they relate to loss of ecosystem services under a changing climate using remote sensing technique InSAR.

Figure 2: Shoreline change along the Nigerian Transgressive Mud coast (Dada et al., 2019).

Research Collaborations

Interested collaborators are welcome to join our research team from any part of the world. Previous works had been carried out in cooperation with investigators from China and France.

Figure 3: (a-f) Coastline retrogradation along the Mahin Mud coast. Most of the houses along the Mahin mud coast at Ayetoro and other areas have been overthrown by these gullies (Dada et al., 2019).

Figure 4: Coastal flooding events at Ayetoro Coastal Community along the Nigerian Transgressive Mahin Mud coast (Dada et al., under review).

References

Borman T.G., Schmidt, J., Adams J.B., Mfikili, A.N., Farre R.E., Smit A.J., 2016. Relative sea-level rise and the potential for subsidence of the Swartkops Estuary intertidal marshes, South Africa. *South Africa Journal of Botany* 107, 91-100. <https://doi.org/10.1016/j.sajb.2016.05.003>

Church, J., Clark, P. U., Cazenave, A., Gregory, J., Jevrejeva, S., Levermann, A., Merrifeld, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D. and Unnikrishnan, A. S., 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, chap. Sea Level Change, Cambridge University Press, Cambridge, United Kingdom.

Dada, O.A., Qiao, L.L., Ding, D., Li, G.X., Ma, Y.Y and Wang, L.M., 2015. Evolutionary trends of the Niger Delta Shoreline during the last 100 years: Responses to rainfall and river discharge. *Marine Geology* 367, pp. 202-211.

Dada, O.A., Li, G.X., Qiao, L.L., Ding, D., Ma, Y.Y and Xu J.S., 2016a. Response of wave and coastline evolution to global climate change off the Niger Delta during the past 110 years. *Marine Systems*, 160, pp. 64-80.

Dada, O.A., Li, G.X., Qiao, L.L., Ding, D., Ma, Y.Y and Xu J.S., 2016b. Seasonal shoreline behaviours along the arcuate Niger Delta coast: Complex interaction between fluvial and marine processes. *Continental Shelf Research*, 122, pp. 51-67.

Dada, O.A., Li, G.X., Qiao, L., Asiwaju-Bello, Y.A., Anifowose, A.Y.B., 2018. Recent Niger Delta shoreline response to Niger River hydrology: Conflict between forces of Nature and Humans. *J. African Earth Sciences* 139, 222-231

Dada, O.A., Agbaje, A.O., Adesina, R.B., and Asiwaju-Bello, Y.A., 2019. Effects of coastal land use on coastline dynamics along the Nigerian Transgressive Mud Coast. *J. Ocean and Coastal Management* 158, 251-264.

Dada, O.A., Almar R., and Oladapo, M.I. Recent coastal sea level variations and flooding events in the Nigerian Transgressive Mud coast of Gulf of Guinea. *J. Marine Systems* (under review).

Nicholls, R. J., and Cazenave, A., 2010. Sea-level rise and its impact on coastal zones, *Science*, 328, 1517-1520.

Vitousek, S., Barnard, P.L., Fletcher, C.H., Frazer, N., Erikson, L. and Storlazzi, C.D., 2017. Doubling of coastal flooding frequency within decades due to sea-level rise. *Scientific Reports* 7: 1399.

Formation and types of thermal inversion in the Bay of Bengal

K M Azam Chowdhury^{1,2,4}, Wensheng Jiang^{1,3*}, Guimei Liu⁴, Md. Kawser Ahmed², Sirajuddin Md. Babar Chowdhury⁵, Nuruddin Md. Khaled Chowdhury⁶

1. College of Oceanic & Atmospheric Sciences, Physical Oceanography Laboratory, Ocean University of China, Qingdao, China
2. Department of Oceanography & International Centre for Ocean Governance (ICOG), University of Dhaka, Dhaka, Bangladesh
3. Laboratory of Marine Environment and Ecology, Ocean University of China, Qingdao, Shandong, China
4. National Marine Environmental Forecasting Center, Beijing 100081, China
5. University of Chittagong, Bangladesh
6. Helen DeVitt Jones Fellow, Department of Geosciences, Texas Tech University, Lubbock, Texas, USA

Abstract

The study of vertical layers concerns the hydrodynamics, biogeochemistry, climate system and cyclogenesis of an ocean. The thermal inversion (TI) formation and isothermal layer depth (ILD) in the Bay of Bengal (BoB) were analyzed from 2009 to 2013 using Argo profile float data. A new numerical methodology is devised to determine the isothermal layer depth when two categories of thermal inversions are encountered from temperature profiles. The TI of the 1st category was higher during the December-March, but that of the 2nd category was higher during May-September. In the case of 0.06°C threshold, April-November temperature profiles display comparatively less TI. The lack of observational data hinders the more extensive study of the TI formation in this bay.

1. Introduction

The coastal area of Bangladesh is a well-known region for extreme cyclones and is strongly affected by this occurrence. The upper layer heat content, known as cyclonic heat potential, is the vital energy required for the intensification and movement of tropical cyclones of this basin (Bishnu et al., 2011). There exist a significant relation between surface temperature and the layer thickness of thermal inversion (TI). So the research on the variation of TI is critical for the nation, as the intensification rate of cyclones is responsive to the initial ILD distribution and TI variation. Thadathil et al. (2002) pointed out that throughout winter, strong TI occurs along the northern Bay of Bengal (BoB). Seasonal change in TI in the north Indian Ocean was studied by Bijoy et al. (2006), and they found that the inversion peak in February is approximately 1.57°C in the BoB. Up to now very few studies are available on the formation and spatio temporal variation of

TI (Thadathil et al., 1992 & 2002 & 2007 & 2016; Bijoy et al., 2006; Girishkumar et al., 2013) in the BoB, furthermore, those studies are not basin-wide, some focus on part of the bay; and some pay attention to short periods of time. In the present study two categories of TI are defined and their spatiotemporal distribution in the BoB is described briefly.

2. Materials and Method

2.1 Data

The temperature and salinity data of Argo profiling floats for the duration of January 2009 to December 2013 are used here. The quality control procedures have reduced the number of Argo profiles from 10626 to 9057. Each profile is interpolated by following the cubic spline methodology for 1-meter vertical intervals starting from 5 meters depth.

2.2 Method

Argo float cannot provide correct data at the surface of the ocean. In this study, the temperature and salinity at 5 meters depth are considered to be the surface temperature and salinity, respectively.

In this study Mixed-layer depth (MLD) is calculated by using fixed density criteria (i.e., 0.2 kgm⁻³ increase in sigma-t from surface value) (Narvekar et al., 2014). The ILD is calculated using the following mathematical equations:

- i. $T_{ILD} = (T^2 - 1) \text{ } ^\circ\text{C}$ (Thadathil et al., 2007 & 2016) [When TI is absent]
- ii. $T_{ILD} = T_{MLD} \text{ } ^\circ\text{C}$ (Thadathil et al., 2007) [When the TI of the 1st category is present]
- iii. $T_{ILD} = [T_{MLD} \cdot (1 - TI^m)] \text{ } ^\circ\text{C}$ [When the TI of the 2nd category is present]

Where

T_{ILD} = ILD temperature

T_{MLD} = MLD temperature

T^2 = 5 meters depth temperature (surface/initial value)

TI^m = mean temperature increase within thermal inversion layer.

Figure 1: Vertical profiles of temperature (red line) and sigma-t (green line) depicting thermal inversion (TI) of the 1st category (left) and the 2nd category (right) in the Bay of Bengal. MLD (upper rectangle), ILD (upper and middle rectangle) and BLT (middle rectangle) are shown in Fig and star represent maximum temperature increase within thermal inversion profile.

Barrier layer thickness (BLT) is calculated as the depth difference between isothermal layer depth and mixed-layer depth (i.e., $BLT = ILD - MLD$).

When TI is absent, the general definition of ILD is defined as the depth where the temperature of sea water is 1°C lower than that at 5 meters depth (considering the value at 5 meters depth as the surface/initial value). The calculation of ILD is quite complex, while the temperature profile displays TI. Sprintall et al. (1992) has compared ILD values with the top of the thermocline depth. He has also compared the MLD based on a sigma-t criterion that was undertaken in three equatorial regions of the tropical oceans. A shallower mixed layer (ML) compared to an isothermal layer (IL) is profound in the above-mentioned study. TI exposed different stages such as growth stage, mature stage and decay stage at certain times. The general definition of ILD (i.e., when TI is absent) can cause overestimation in calculating ILD for those stages of the TI profile case. In the present research, while assessing the ILD, two different categories of TI are noted, one that stays deeper than MLD (termed as the TI of the 1st category) and the other inversion that stays within the MLD or MLD is greater than the highest inversion temperature depth (termed as TI of 2nd category) (see Fig. 1). In the case of the 1st category of TI, MLD is shallower than the highest TI temperature depth, but in the case of the 2nd category of TI, MLD is deeper than the highest TI temperature depth.

The TI of the 1st category is similar to what defined by Thadathil et al. (2007), i.e., ILD is the maximum depth where the MLD temperature is repeated below the ML layer. In the case of the 2nd category, ILD is defined in such a way that it can balance the overestimation in the calculation of ILD compared to using the general definition of ILD (definition while TI is absent). As MLD is greater than the highest inversion temperature depth in the case of the 2nd category, there is no scope for

the MLD temperature to repeat below the MLD layer. Therefore, the definition of the TI for the 1st category cannot be used in the 2nd category. For that rare case, we need to devise a new methodology to calculate the TI of the 2nd category. In the case of the 2nd category, ILD is the maximum depth where the MLD temperature is reduced by 1°C minus the mean temperature increase (TI^m) within the inversion layer.

3. Formation of TI

TI is a very common phenomenon in the Bob. The details of TI distribution (1st category and 2nd category) compared with two different threshold temperatures (0.002°C and 0.06°C) are shown in Table 1. Argo temperature profiles have an error of $\pm 0.002^{\circ}\text{C}$. Using a threshold temperature of 0.002°C , it can be noted that out of a total of 7442 profiles, 45% (3319 in no.) of profiles have shown the TI of the 1st category, while 33% (2463 in no.) of profiles have shown the TI of the 2nd category, but only 22% (1660 in no.) of profiles have shown the absence of TI. The average temperature difference in the TI of the 2nd category is 0.06°C . When using a threshold temperature 0.06°C , it can be noted that out of a total of 7442 profiles, 40% (3002 in no.) of profiles have shown the TI of the 1st category, while 13% (972 in no.) of profiles have shown the TI of the 2nd category, but 47% (3468 in no.) of profiles have shown the absence of TI. In the case when threshold temperature is 0.002°C , the highest number of the TI of the 1st category is noted throughout January within 74% of profiles, which is consistent with the result of Thadathil et al. (2002), stating that strong temperature inversions were prominent throughout winter; and the lowest is noted throughout April and May. The highest number of the TI of the 2nd category is noted throughout June with 62% of profiles, and the lowest was displayed throughout March with 14% of profiles. In the case when threshold temperature is 0.06°C , the highest number of the TI of the 1st category is noted throughout January with 70% of profiles, and the

lowest is noted throughout April and May with 11% of profiles. The highest number of the TI of the 2nd category is noted throughout June and July with 22% of profiles, and the lowest was displayed throughout February with 5% of profiles. If we omit less than a 0.06°C increase in subsurface temperature, then 47% of profiles indicate no TI throughout the year (Table 1).

MONTH	THRESHOLD 0.002°C			THRESHOLD 0.06°C		
	TI1 %	TI2 %	NTI %	TI1 %	TI2 %	NTI %
Jan	74	20	5	70	11	19
Feb	63	17	19	60	5	35
Mar	55	14	30	52	6	42
Apr	15	37	49	11	16	73
May	15	58	27	11	20	69
Jun	19	62	19	15	22	63
Jul	23	61	16	18	22	60
Aug	20	52	28	17	18	65
Sep	38	41	21	35	16	49
Oct	41	24	35	35	7	58
Nov	49	28	23	38	13	49
Dec	61	29	9	57	17	25
Average	45%	33%	22%	40%	13%	47%

Table 1: Comparison of the thermal inversion (TI) profiles (1st category, 2nd category and absence of TI) using two different threshold temperatures(in percentages)

Note: TI1= thermal inversion of the 1st category, TI2= thermal inversion of the 2nd category, NTI= absence of thermal inversion, Boldface Italic numbers show distinguishing features.

The month wise distribution of several parameters in different categories of TI is described in Tables 2&3. Within the 1st category of TI, the average temperature increase is 0.6°C, average maximum TI is 2.1°C, average depth of TI starts at 9.9 meters, average depth of maximum temperature increase is 35.4 meters, average MLD is 25.7 meters, and average ILD is 49.1 meters. Within the 2nd category of TI, the average temperature increase is 0.1°C, average maximum TI is 0.8°C, average depth of TI starts at 8.3 meters, average depth of maximum temperature increase is 26.6 meters, average MLD is 39.1 meters, and average ILD is 54.2 meters. From Table 2, it is noted that maximum TI depth is shallower than MLD in the 2nd category of TI, which is opposite the case in the 1st category. It is also remarkable that MLD is deeper in the 2nd category compared to 1st category of TI. Note: DT= Difference in temperature during inversion in °C, MDT=Maximum difference in temperature in a profile during inversion, DTIS= Depth of TI start, DMTI=Depth of maximum TI, MLD= Mixedlayer depth, ILD=Isothermal layer depth.

4. Conclusion

TI is a phenomenon that occurs most prominently along the northern BoB. The TI layer leads to intensified ILD in the respective periods. A new numerical methodology (i.e., 2nd category of TI) is devised to determine ILD when two categories of TI were encountered from temperature profiles. The upper layer variation is related to several

MONTH	TI OF 1 ST CATEGORY					
	DT	MDT	DTIS	DMTI	MLD	ILD
Jan	2.0163	4.6709	10.41	56.58	33.68	78.66
Feb	1.7333	4.4366	11.64	56.32	27.08	79.12
Mar	0.7993	2.4473	15.04	49.91	20.43	66.97
Apr	0.2119	0.9293	7.35	14.07	14.55	28.3
May	0.5564	1.1779	9.6	25.07	25.34	37.09
Jun	0.1865	0.8233	8.74	33.03	31.54	43.53
Jul	0.1905	1.1324	10.02	40.71	37.99	48.04
Aug	0.2655	0.8396	8.56	27.43	22.38	39.29
Sep	0.2444	0.9644	7.36	23.82	20.96	32.61
Oct	0.2617	1.1691	10.48	27.50	23.71	41.06
Nov	0.3717	2.7300	9.82	29.72	21.78	41.14
Dec	0.7485	4.2280	9.48	40.81	28.41	53.77
Averages	0.6	2.1	9.9	35.4	25.7	49.1

Table 2: Month wise distribution of TI profiles (1st category) with different parameters

MONTH	TI OF 2 ND CATEGORY						TI ABSENT	
	DT	MDT	DTIS	DMTI	MLD	ILD	MLD	ILD
Jan	0.0949	1.0123	12.11	48.93	58.03	69.69	51.12	67.30
Feb	0.0773	1.1764	8.44	28.77	49.67	74.92	41.63	73.05
Mar	0.0991	0.7577	7.02	13.18	23.19	53.40	23.82	49.63
Apr	0.0886	0.8166	6.43	11.9	21.83	33.05	20.85	32.92
May	0.0723	0.6761	8.01	19.36	30.74	39.92	28.59	38.67
Jun	0.0664	0.7312	7.62	29.08	41.45	49.91	34.06	46.94
Jul	0.0642	0.5893	8.49	31.11	43.46	53.01	39.47	53.19
Aug	0.0727	0.5141	7.8	31.22	46.18	57.88	36.54	55.20
Sep	0.0629	0.4521	7.28	30.52	45.37	59.58	27.85	56.13
Oct	0.0470	0.4511	8.28	21.18	37.59	53.37	32.76	52.84
Nov	0.0732	0.9066	8.66	20.57	27.96	48.69	26.21	46.63
Dec	0.1308	1.0800	8.96	33.94	43.21	56.97	29.73	57.44
Averages	0.1	0.8	8.3	26.6	39.1	54.2	32.7	52.5

Table 3: Month wise distribution of the TI of the 2nd category and TI absent with different parameters

other physical processes in the BoB. The intensification of cyclones can be extensively examined from the perspective of TI heat content. Detailed studies should be done in the future work.

Acknowledgement

This study was supported by the NSFC Shandong Joint Fund for Marine Science Research Centers (Grant U1606402). We express gratitude to the people who support us various ways to complete this study. We are also grateful to UNESCO/IOC-ODC and CLIVAR for their overall facilities.

References

Bijoy, T., C. Gnanaseelan, and P. S. Salvekar (2006). Seasonal evolution of temperature, inversions in the north Indian Ocean.Current Science, Vol. 90, No. 5.

B. Kumar, & A. Chakraborty (2011). Seasonal variability of heat content and isothermal layer in the Bay of Bengal. Journal of Oceanography and Marine Science Vol.2(3), pp. 63–77.

G. Kumar, M. S., M. Ravichandran, & M. J. McPhaden (2013). Temperature inversions and their influence on the Mixed Layer heat budget during the Winters of 2006–2007 and 2007–2008 in the Bay of Bengal. *Journal of Geophysical Research: Oceans*, 118, 2426–2437. <http://doi.org/10.1002/jgrc.20192>.

J. Narvekar and S. Prasanna Kumar (2014). Mixed layer variability and chlorophyll a biomass in the Bay of Bengal. *Biogeosciences*, 11, 3819–3843. <http://doi:10.5194/bg-11-3819-2014>.

Kara AB, Rochford PA, Hurlburt HE. 2000. An optimal definition for ocean mixed layer depth. *Journal of Geophysical Research* 105: 16 803–16 821.

Madhupratap, M., Gauns, M., Ramaiah, N., Kumar, S.P., Muraleedharan, P.M., De Sousa, S.N., Sardesai, S., & Muraleedharan, U. (2003). Biogeochemistry of the Bay of Bengal: physical, chemical and primary productivity characteristics of the central and western Bay of Bengal during Summer monsoon 2001. *Deep Sea Research Part II*, 50(5), pp.881–896. [https://doi.org/10.1016/S0967-0645\(02\)00611-2](https://doi.org/10.1016/S0967-0645(02)00611-2).

P.Thadathil, & Aravind, K.G. (1992). Surface Layer Temperature Inversion in the Arabian Sea during Winter. *Journal of Oceanography*, Vol. 48, pp. 293 - 304. <https://doi.org/10.1007/BF02233989>.

P.Thadathil, V. V. Gopalakrishna, P. M. Muraleedharan, G. V. Reddy, N. Araligidat, & S. Shenoy, (2002). Surface layer temperature inversion in the Bay of Bengal. *Deep Sea Research Part I*, 49, 1801–1818. [https://doi.org/10.1016/S0967-0637\(02\)00044-4](https://doi.org/10.1016/S0967-0637(02)00044-4).

P.Thadathil (2007). Observed seasonal variability of BL in the BoB. *Journal of Geophysical Research*, (C: Ocean):112(2). <http://doi:10.1029/2006JC00365>.

P.Thadathil, I. Suresh, S. Gautham, S. Prasanna Kumar, M. Lengaigne, R. R. Rao, S. Neetu, and A. Hegde (2016), Surface layer temperature inversion in the Bay of Bengal: Main characteristics and related mechanisms, *J. Geophys. Res. Oceans*, 121, 5682–5696, doi:10.1002/2016JC011674.

Sprintall, J., & Tomczak, M. (1992). Evidence of Barrier Layer in the surface layer of the tropics. *Journal of Geophysical Research*, 97, 7305–7316. <https://doi.org/10.1029/92JC00407>.

Modeling of the Biogeochemical Cycle and Dispersion of Nitrogen off Lake Idku Outlet, Abu Qir Bay, Egypt

Amr T. Salama¹, Samir M. Nasr¹, Ahmed E. Rifaat²

1. Institute of Graduate Studies and Research, Alexandria University, Egypt
2. National Institute of Oceanography and Fisheries, Egypt

Introduction

The Egyptian Mediterranean coast hosts five lakes which are collectively called the Northern Lakes. These lakes receive their water mainly from agricultural drainage systems. Over the years, the water quality of these drains has continually been deteriorating, and the levels of pollutants have increased inside the lakes (Siam and Ghobrial, 2000; Okbah and El-Gohary, 2002; Saad et al., 2003; Ali and Khairy, 2016;) (1,2,3,4), which in turn could affect adversely on the water quality of the Mediterranean coastal water, as the northern lakes discharge their water directly to the coastal water through natural channels.

One of these lakes is Idku Lake, which suffered from a significant increase of nitrogen levels over the last few decades (Saad and Younes, 2006; EEAA, 2015) (5,6), and affected negatively on the water quality of Abu Qir bay (Emam et al., 2013; Hemaida, 2016; Ismail and Osman, 2016;) (7,8,9), which is connected directly to the lake through an outlet (Figure 1).

Figure 1: Map illustrating the study area (Coordinates are in UTM).

In this study we try to investigate the dynamics of the nitrogen components that disperse from the lake into the bay, by taking into account the impact of hydrodynamics on the dispersion patterns of nitrogen forms as well as the biogeochemical cycle.

Materials and Methods

A coupled high resolution hydrodynamic -

biogeochemical model was developed on the basis of Delft3D over the period from November 2015 to May 2016. The model comprised most of the nitrogen cycle components and processes (Figure 2a), as well as the hydrodynamics controlling factors within the model domain including (water levels at the bay and the Lake Outlet, wind direction and speed, bottom roughness, and water viscosity).

Delft3D is a modeling suite was developed by the Deltares Institute (The Netherlands), Delft3D consists of a number of modules, which are linked to and integrated with one another. The most common modules for estuarine and coastal ecological modeling approaches, are (Delft3D-FLOW and Delft3D-WAQ). The FLOW module is a multi-dimensional (2D or 3D) hydrodynamic (and transport) simulation program which calculates non-steady flow and transport phenomena resulting from tidal and meteorological forcing on a curvilinear, boundary fitted grid or spherical coordinates (Delft3D-FLOW, 2016) (10). Whilst, Delft3D-WAQ simulates the far and mid-field water and sediment quality due to a variety of transport and water quality processes. It includes several advection diffusion solvers and an extensive library of standardized process formulations with user-selected substances (Delft3D-WAQ, 2016) (11).

Figure 2a: The components and processes of the developed biogeochemical model.

The hydrodynamic model grid was developed via RGFRID which is an extension tool in Delft3D suite. The grid consisted of 405 grid points in horizontal direction and 201 grid points in vertical direction. As presented in (Figure 2b), the grid cells were intensified particularly around the outlet. Moreover, the grid cell resolution was (33*30 m), while in the intensified area, the grid cell resolution was (12.14 * 14.05 m).

After ending the simulation of the hydrodynamic model, a coupling process was operated, for integrating the outputs of the hydrodynamic model with the biogeochemical model processes and components.

Figure 2b: The grid of the developed hydrodynamic model.

Results and Discussion

The model results showed that the water level in the lake was higher than those in the bay (Figure 3), leading the dispersion of nitrogen toward the bay (Figure 4). In addition, a hypothetical scenario has been set to investigate the impact of sea level rise at the bay on the fate of nitrogen components in Idku Lake. The results pointed out that the increase of water level at Abu Qir bay will lead to seawater intrusion toward the lake, which in turn may sequester the nitrogen forms inside the lake, and consequently increase the eutrophication rate in Idku lake.

Figure 3: Water levels variation at the outlet and the bay over the simulation time.

Figure 4: Spatial dispersion of Total Nitrogen (mg/l-N) from Idku Lake outlet to Abu Qir bay over different times of the simulation.

Regarding the nitrogen cycle, the dissolved oxygen levels have played a key role in regulating the cycle processes.

As the study area is a shallow coastal area, and the water column there was turbulent oxygenated, this resulted in the increase of the nitrification rate as well as the inhabitation of the denitrification process. Detrital nitrogen in algal cells was transferred to sediments by settling. Also, the sedimentary nitrogen mineralization process had a pivotal role in releasing more nitrogen fluxes back to the water column (Figure 5).

Figure 5: Schematic overview of the systematic nitrogen transformations over the model domain. Negative fluxes (gN) refers to the loss of N-fluxes by processes, while the positive fluxes refer to the gain of N-fluxes.

Conclusion

Most of nitrogen forms that derived from Idku lake outlet, were dissipated and transported over the open boundaries of the model domain. In addition, the residence time of water masses was very low over the simulation time, decreasing the rate of retention of nitrogen fluxes over the study area.

Acknowledgement

The Authors of this work are grateful to the Egyptian Science & Technology Development fund (STDF) for funding this work, under the project number (25445).

References

Siam, E. and Ghobrial, M. (2000). Pollution influence on bacterial abundance and chlorophyll-a concentration: case study at Idku Lagoon, Egypt. *Scientia Marina*. 64(1): 1-8.

Okbah, M. and El-Gohary, S. (2002). Physical and chemical characteristics of Lake Edku water, Egypt. *Mediterranean Marine Science*. 3(2): 27-40.

Saad, M.A., Abdel-Moati, A.R., and Abdel-Atti, N.A. (2003). Effect of Anthropogenic Pollution on Nutrient Specis in an Egyptian Wetland. In: *The International Conference on Water and Environment*. Bhopal, India. 136-148 p.

Ali, E.M. and Khairy, H.M. (2016). Environmental assessment of drainage water impacts on water quality and eutrophication level of Lake Idku, Egypt. *Environmental Pollution*. 216: 437-449.

Saad, M.A. and Younes, W.A. (2006). Role of phosphorus and nitrogenous species in water quality of a coastal Egyptian heavily polluted Mediterranean basin. *International Journal of Oceans and Oceanography*. 1(1): 1-19.

EEAA. (2017). Idku Lake (Field Campaigns Reports). Accessed: (3 October 2017). Available from: <http://www.eeaa.gov.eg/en-us/topics/water/lakes.aspx>.

Emam, W., Saad, A., El-Moselhy, K.M., and Owen, N. (2013). Evaluation of water quality of Abu Qir Bay, Mediterranean coast, Egypt. *International Journal of Environmental Science and Engineering (IJESE)*. 4: 47-54.

Hemaida, H.A. (2016). Impact of Drainage Effluents on the water quality of Abu-Qir Bay, Alexandria, Egypt. *Journal of High Institute of Public Health*. 37(1): 155-168.

Ismail, M.M. and Osman, M.E. (2016). Seasonal fluctuation of photosynthetic pigments of most common red seaweeds species collected from Abu Qir, Alexandria, Egypt. *Revista de Biología Marina y Oceanografía*. 51(3): 515-525.

Delft3D-FLOW. (2016). *Delft3d-Flow User Manual (Simulation of Multi-Dimensional Hydrodynamic Flows and Transport Phenomena, Including Sediments)*. Deltares, The Netherlands. 702 p.

Delft3D-WAQ. (2016). *Delft3d Water Quality Manual (Versatile Water Quality Modelling In 1d, 2d or 3d Systems Including Physical, (Bio)Chemical and Biological Processes)*. Deltares, Delft, The Netherlands. 394 p.

Sediment Transport and Coastal Erosion Processes in Southern Spain Under Climate Change Scenarios

Andrea Lira-Loarca¹, Manuel Cobos¹, Agustín Millares¹, Asunción Baquerizo¹, Miguel A. Losada¹

1. Andalusian Institute for Earth System Research (IISTA), University of Granada, Spain

Abstract

This report summarizes the work being carried out by the Environmental Fluid Dynamics Group of the University of Granada (UGR) within the European project AQUACLEW (Advancing quality of climate services for European water). The overall goal of AQUACLEW is to use innovative research techniques and integrated co-development with users to advance the quality and usability of several climate services for a number of water related sectors under climate change scenarios.

Within AQUACLEW-UGR we will analyze the changes in physical processes such as sea waves, fluvial discharges and sediment transport, which interact and control the dynamics of the study zones as well as the integrity of the physical environment and ecologic condition. The analysis will be made under different climate change and sea level rise scenarios. The case studies are the Guadalfeo and Adra rivers, located in Granada and Almería (Spain) respectively, and their deltas and adjacent coasts.

This work aims to have a better understanding of climate change impacts and to improve the adaptive capability of fluvial, coastal and transition zones in Mediterranean environments that have been altered by human activities. More precisely, the specific objectives of the project are (1) to develop advanced feedback loops along the whole chain of information production of the climate services, to better understand the needs of the users, (2) to contribute to the quality and usability of climate services at fluvial, coastal and transition zones of semiarid watersheds in the Mediterranean Andalusian coast and (3) to evaluate if and how the co-developed work facilitates decision-making by evaluating two real-world case studies.

Introduction

Deltaic systems are unique landscapes in continuous transformation due to the sculpting action of marine and fluvial dynamics. In the last two centuries, the occupation of these areas due to the growth of tourism and the industrial development has favored the irrational use of their resources. The non-integrated management of these systems leads to important economic costs and does not meet the goals of the

EU framework directives of application for rivers and coastal zones. In particular, it deviates from the principles of the Water Framework Directive 2000/60 and the Directive 2014/89 for maritime spatial planning.

Mediterranean deltas are especially vulnerable to sea level rise, which is one of the most important causes of delta retreat around the globe (Duong et al., 2016). Therefore, the present issues found in deltaic systems and the erosion in the adjacent coasts will become aggravated in a climate change scenario that includes sea level rise and changes in the frequency and persistence of storms.

The evidence of climate change has increased the demand of data and favoured the creation of climate services (CS) that provide projections at different time and spatial scales. In AQUACLEW-UGR, we propose to use innovative research techniques and integrated co-development with users to advance the quality and usability of several CSs in coastal areas. This work will contribute to the Environmental Fluid Dynamics Group's ongoing research into how climate change impacts fluvial sediment dynamics and how this in turn affects coastal processes.

The Environmental Fluid Dynamics Group (<http://gdfa.ugr.es>) of the Andalusian Institute for Earth System Research (IISTA) of the University of Granada, Spain is an interdisciplinary group whose expertise comprises a wide range of fields, such as:

- Integrated management of ports and coasts and risk analysis in engineering.
- Hydro- and morpho- dynamics of coastal and fluvial environments, including rivers, beaches, deltas and estuaries.
- Interaction between nature and infrastructures.
- Theoretical, numerical and experimental analysis of fluvial, marine and coastal structures.
- Characterization and simulation of multivariate maritime climate data.

AQUACLEW

AQUACLEW (<http://aquaclew.eu>) is part of the ERA4CS (European Research Area for Climate Services) project,

Figure 1: Project methodology for user-driven design, development and evaluation of climate services

an initiative by JPI Climate (Joint Programming Initiative), co-funded by the European Commission (Grant 690462) with a timeframe of 2017-2020. It brings together nine complementary organizations from across Europe, with different experience and expertise in developing climate services, providing data and collaborating with users.

The overall goal of AQUACLEW is to use innovative research techniques and integrated co-development with users to advance the quality and usability of several climate services for a number of water related sectors under climate change scenarios.

The methodology will be tested in seven diverse real-world local case-studies for climate adaptation decision-making across Europe (from southern Spain to northern Sweden) which includes:

- Flash flood risks in pre-alpine regions.
- Flash flood risks in urban areas.
- Drought and water resource allocation for industry, tourism, agriculture and energy.
- Hydropower production.
- Biodiversity decline.
- Agricultural production.
- Sediment transport and coastal erosion.

AQUACLEW-UGR

In the case of the University of Granada (AQUACLEW-UGR), the project will focus on the issue of sediment transport and coastal erosion by studying fluvial and coastal interactions under Mediterranean climate conditions and climate change scenarios. This research will lead to the development of innovative new climate services for the sustainable management of these systems at a regional scale. For this, two case studies have been proposed: the Guadalfeo and Adra rivers, their deltas and adjacent coasts.

The Guadalfeo and Adra rivers are located in Granada and Almería, respectively, both flowing into the Alborán sea in the south of Spain. Their basins are semiarid, have a small size (1250 km² and 746 km², respectively) and show significant gradients with altitude that greatly condition the erosive dynamics and sediment transport to their adjacent coasts.

Guadalfeo River basin

In the Guadalfeo River basin (southern Spain, Figure 1), the same as in other rivers of the Mediterranean coast, the gravel nature and the huge availability of sediment boost the bed load sediment transport (Millares et al., 2014a; 2014b), being the main contributor to the supply of coastal sediments.

The river was dammed 19 km upstream from the mouth in 2004, regulating 85% of the basin runoff. As a consequence, the delta currently presents erosion problems and severe coastline retreat (Figures 3-4).

Figure 3: Guadalfeo river mouth before (left) and after (right) river damming

Figure 2: Location of the study site

Figure 4: Aerial images of the Guadalfeo river mouth with highlighted shoreline

This area has become particularly vulnerable not only to human-driven changes but also to the effects of global warming with respect to the increase of the frequency and intensity of storms and sea-level rise giving place to flooding events and coastal erosion (Figure 5).

Figure 5: “Poniente beach” before and during a storm event in 2017

This stretch of beach is occupied by an exclusive hotel complex, golf fields, restaurants, farming settlements and residential properties that are primarily summer homes. Therefore, the coast has high environmental and recreational value. For this reason, artificial nourishment projects have been frequently conducted since the construction of the dam (Figure 6). However, the success of these interventions has been very limited partly due to the too fine sediment used for the nourishment, the morphology of the nourished coastline (Bergillos et al., 2018) and the increasing sea-level and climate change impacts.

This project will focus on the sediment transport and coastal erosion problem and aims to bring research closer to coastal managers, providing them with an overview and detailed characterization of the morphodynamics in the study sites and management alternatives under climate change and sea level rise scenarios along with an uncertainty assessment.

References

Bergillos, R. J., López-Ruiz, A., Principal-Gómez, D., Ortega-Sánchez, M. (2018). An integrated methodology to forecast the efficiency of nourishment strategies in eroding deltas. *Science of the Total Environment*, 613-614, 1175–1184.

Duong T.M., Ranasinghe R., Walstra D., Roelvink D., (2016). Assessing climate change impacts on the stability of small tidal inlet systems: Why and how? *Earth-Science Reviews*, 154, 369-380.

Millares, A., M. Polo, A. Moñino, J. Herrero, M. Losada. (2014a), Bedload dynamics and associated snowmelt influence in mountainous and semiarid alluvial rivers, *Geomorphology*, 206, 330–342.

Millares, A., M. Polo, A. Moñino, J. Herrero, M. Losada (2014b). Reservoir sedimentation and erosion processes in a snow-influenced basin, in Southern Spain, in *Reservoir Sedimentation—Special Session on Reservoir Sedimentation of the 7th International Conference on Fluvial Hydraulics, River Flow*, pp. 91–98, Taylor and Francis Group, London, U. K.

Figure 6: “Playa Granada” before and after nourishment works

Research on the Energy and Carbon Dioxide Fluxes in the Atmosphere-Ocean Interface of the Tropical Coastal Ocean

Yusri Yusup^{1,2}, John Stephen Kayode¹, Abbas F.M. Alkarkhi³

1. Environmental Technology, School of Industrial Technology, Universiti Sains Malaysia, USM 11800, Pulau Pinang, Malaysia.
2. Centre for Marine & Coastal Studies (CEMACS), Universiti Sains Malaysia, Pulau Pinang, Malaysia.
3. Malaysian Institute of Chemical & Bioengineering Technology, Universiti Kuala Lumpur, 78000 Melaka, Malaysia.

Abstract

The tropical coastal ocean is a vital jigsaw-piece to understanding exchanges of energy, moisture, and carbon between the atmosphere and the ocean biospheres. Carbon dioxide, latent and sensible heat fluxes, microclimate variables, and surface water temperature data were collected using an instrumented platform at the north-eastern end of a tropical island (5°28'6"N, 100°12'1"E), which is within the intertidal zone on the continental shelf of the Malacca Straits in the southern South China Sea. The instruments were installed on a stainless-steel platform extending the pre-existing pier. Starting from November 2015, data was collected for more than two years, during which all cycles of the seasons or Monsoons were encompassed. To determine the carbon dioxide and energy fluxes, the eddy covariance method was used. The results of the analyses are briefly discussed. The data and results have important applications in improving numerical weather models. Our website at <http://atmosfera.usm.my> periodically displays the recent 24-hour fluxes and microclimate variables measured.

Introduction

The coastal ocean is a major component of the biogeochemical carbon cycle (Bauer et al., 2013; Chen et al., 2013; Gruber et al., 2009). The coastal ocean contributes 30% of the total carbon dioxide (CO₂) sink of the global ocean (Chen & Borges, 2009). Some researchers found that this contribution could be even larger. Tsunogai et al. (1999) reported that the continental shelves can take up to 1 Pg of CO₂ annually (i.e., 50% of the global ocean CO₂ uptake).

The high heat capacity of the ocean enables it to regulate the weather and climate through the exchanges of energy and mass. The tropical coastal ocean remains a vital jigsaw-piece to understand the global atmosphere-ocean exchanges. With the establishment of the eddy covariance station, two years of continuous observations of the energy and mass fluxes, and microclimate variables are made possible, which can be used to elucidate the atmosphere-ocean processes

of the tropical coastal ocean.

Locations of the highest exchange of carbon, moisture, and energy have yet to be discovered but prove important to further improvement of climate and weather predictions. Therefore, the purpose of the work is to understand the role of the tropical coastal ocean as the exchangers of energy and mass fluxes and their links to the microclimate and seasonal drivers.

Site description and instrumentation

CO₂, energy fluxes, microclimate variables, as well as the sea water temperature, were collected at a tropical coastal ocean at the north-western end of the island of Pulau Pinang, Malaysia (Figure 1). The station, which is named the "Muka Head Station", is located within the intertidal zone and enclosed strait. The wave heights there are relatively low, which is approximately 0.25 m. The mean water depth beneath the station is 4 m. The location experiences tropical Monsoons: The Northeast Monsoon from December to March, the Spring Transitional Monsoon from April to May, the Southwest Monsoon from June to September, and the Fall Transitional Monsoon from October to November.

The CO₂ and energy fluxes were measured using an eddy covariance system, which was installed on a stable stainless-steel platform extending a pre-existing pier so that the system would be directly over the tropical coastal ocean. The eddy covariance system included a 3-D sonic anemometer (model 81000V, RM Young, USA) and an open-path CO₂/H₂O gas analyzer (model LI-7500A, LI-COR, Inc., USA) installed 4.1 m above the sea water surface. The gas analyzer and sonic anemometer were factory-calibrated before deployment. A data logger (model LI-7550 Analyzer Interface Unit, LI-COR, Inc., USA) was used to record the eddy covariance data at a frequency of 20 Hz. The sonic anemometer was also used to measure wind speed and wind direction. The flux data were averaged in 30-min blocks.

Figure 1: Location of the eddy covariance and Biomet system at the Muka Head station denoted as a black circle ($5^{\circ}28'6''\text{N}$, $100^{\circ}12'1''\text{E}$).

The “Biomet” system, which consisted the slow-response sensors, measured the microclimate variables. The sensors were the atmospheric temperature and relative humidity (RH) sensor (model HMP155, Vaisala, Finland; the accuracy of TA is $\pm 0.7^{\circ}\text{C}$ and the accuracy of RH is $\pm 1.7\%$), the pyranometer (RG) (model LI-200SL, LI-COR, Inc., USA; error was $<5\%$), and the net radiometer (RN) (model NR LITE 2, Kipp & Zonen, Inc., USA; sensitivity was $13.6 \mu\text{V W}^{-1} \text{m}^{-2}$). These sensors complemented the eddy covariance system. Sea water temperatures at depths of 0.5 m (TS1) and 2.5 m (TS2) were measured using two thermistors (LI-COR, Inc., USA) with the accuracy of $\pm 1^{\circ}\text{C}$. Since TS1 was positioned near the water surface, the temperature measurements were assumed to be the sea surface temperature. An additional data logger recorded the Biomet data (model 9210b Xlite, Sutron Corporation, USA) at a sampling frequency of 1 min and averaged in 30-min blocks. All slow-response sensors were factory-calibrated before installation.

Results and Discussion

The CO_2 flux reveals the dynamical nature of the CO_2 exchange of the coastal tropical ocean on monthly to seasonal time scales. Overall, the tropical coastal ocean was a strong CO_2 sink with an annually-averaged mean of $-0.07 \mu\text{mol m}^{-2} \text{s}^{-1}$ (or $-2.10 \text{ mol C m}^{-2} \text{yr}^{-1}$). Figure 2 shows the averaged CO_2 flux reported at other locations in the South East Asia and East Asia regions (Bates, 2006; Shim et al., 2007; Wang et al., 2000; Zhai & Dai, 2009; Zhai et al., 2005; Zhai et al., 2007; Zhu et al., 2009).

CO_2 flux was regulated by precipitation and the Monsoons. Larger amount of precipitation led to higher CO_2 flux. The daily-averaged CO_2 flux time series (Figure 3) showed that it absorbed 60% more CO_2 during the Northeast Monsoon and the Spring Transitional Monsoon than during the Southwest Monsoon and Fall Transitional Monsoon. Monthly cumulative precipitation was large and peaked in September 2016, which possibly led to the increasing and sustaining daily-averaged CO_2 flux in October-December 2016. The persistent CO_2 emission could be due to the large riverine carbon input brought by the increased precipitation. While positive CO_2 flux was

Figure 2: Sources and sink of carbon dioxide fluxes (unit in $\text{mol C m}^{-2} \text{yr}^{-1}$) in the South East Asia and East Asia.

intermittent and sudden in other monthly intervals, the October-December months were the only interval that features constant positive CO_2 flux.

Figure 3: Daily-averaged CO_2 flux ($\mu\text{mol m}^{-2} \text{s}^{-1}$) from November 2015 to January 2017; the red curve is the smoothed averaged CO_2 flux values and the shaded area is the 95% confidence interval.

With regards to the overall energy budget, the results show that the southern South China Sea water heats up at the beginning of the year and cools as the year progresses (Figure 4). The three measured parameters of the solar radiation and energy deficit (or energy input and storage), i.e., global (R_G), net (R_N), and deficit (ϵ), peaked in April, and thereafter, the values decreased steadily throughout the year.

The energy stored in the ocean water intensified evaporation (LE), whereas the released sensible heat (H) increased during the cooling periods. During the Monsoons periods between the Monsoons, the change in LE recorded was -30% while that in H was $+18\%$. The inter-annual variations in LE (Figure 4) steadily decreased from

January to May, thereafter the values were considerably constant until September, from when the sensible heat steadily increased towards the maximum. The plateau-like shape produced a near-flat trend from May to September. In contrast to LE, H (Figure 4) increased steadily from January to May and decreased between May and June before increased again until August when it became constant through to December.

Figure 4: Annually-average of the monthly variation in: (a) energy fluxes, which consists of R_G , R_n , and ϵ ; (b) LE, and (c) H (W m^{-2}); vertical dash lines separate the Northeast Monsoon, Spring Transitional Monsoon, Southwest Monsoon, and Fall Transitional Monsoon.

Conclusions

This is the first few high-resolution eddy covariance measurement of CO_2 and energy fluxes in the intertidal zone of the tropical coastal ocean in the southern South China Sea. This location is severely under-sampled in the global network of flux measurements and unrepresented in the climate and weather models. The results show that the tropical coastal ocean is a carbon sink regulated by precipitation and seasonal changes. The results also show that the seasons affected the energy (LE and H) exchanges between the ocean and the atmosphere. The reader can also refer to our research website at <http://atmosfera.usm.my> to view the past 24-hour fluxes and microclimate variables measured.

Opportunities for collaboration

This study demonstrates a need to further collect more data that span multiple years, to fully describe the effects of small and large-scale changes brought by the weather and climate on the exchanges of mass and energy between the tropical coastal ocean and atmosphere. Further work can also be pursued by the “closing” of the energy and mass budgets. This can be achieved by accounting the tide, the storage, and the input of mass as well as water into the mass and energy budgets of the tropical coastal ocean system. More studies and analyses are also needed at other locations to generalize the process-based understanding of the atmosphere-ocean interactions for parameterization purposes in numerical weather and climate models.

References

- Bates, N.R. (2006). Air-sea CO_2 fluxes and the continental shelf pump of carbon in the Chukchi Sea adjacent to the Arctic Ocean. *Journal of Geophysical Research: Oceans*, 111(C10), C10013.
- Bauer, J.E., Cai, W.-J., Raymond, P.A., Bianchi, T.S., Hopkinson, C.S., & Regnier, P.A.G. (2013). The changing carbon cycle of the coastal ocean. [Insight]. *Nature*, 504(7478), 61-70. 10.1038/nature12857
- Chen, C.T.A., & Borges, A.V. (2009). Reconciling opposing views on carbon cycling in the coastal ocean: continental shelves as sinks and near-shore ecosystems as sources of atmospheric CO_2 . *Deep Sea Research Part II: Topical Studies in Oceanography*, 56(8-10), 578-590. 10.1016/j.dsr2.2009.01.001
- Chen, C.T.A., Huang, T.H., Chen, Y.C., Bai, Y., He, X., & Kang, Y. (2013). Air-sea exchanges of CO_2 in the world's coastal seas. *Biogeosciences*, 10(10), 6509-6544. 10.5194/bg-10-6509-2013
- Gruber, N., Gloor, M., Mikaloff Fletcher, S.E., Doney, S.C., Dutkiewicz, S., Follows, M.J., et al. (2009). Oceanic sources, sinks, and transport of atmospheric CO_2 . *Global Biogeochemical Cycles*, 23(1), GB1005. 10.1029/2008gb003349
- Shim, J., Kim, D., Kang, Y.C., Lee, J.H., Jang, S.-T., & Kim, C.-H. (2007). Seasonal variations in pCO_2 and its controlling factors in surface seawater of the northern East China Sea. *Continental Shelf Research*, 27(20), 2623-2636. 10.1016/j.csr.2007.07.005
- Tsunogai, S., Watanabe, S., & Sato, T. (1999). Is there a “continental shelf pump” for the absorption of atmospheric CO_2 ? *Tellus B*, 51(3), 701-712.
- Wang, S.-L., Chen, C.-T.A., Hong, G.-H., & Chung, C.-S. (2000). Carbon dioxide and related parameters in the East China Sea. *Continental Shelf Research*, 20(4), 525-544.
- Zhai, W., & Dai, M. (2009). On the seasonal variation of air-sea CO_2 fluxes in the outer Changjiang (Yangtze River) Estuary, East China Sea. *Marine Chemistry*, 117(1), 2-10.
- Zhai, W., Dai, M., Cai, W.-J., Wang, Y., & Wang, Z. (2005). High partial pressure of CO_2 and its maintaining mechanism in a subtropical estuary: the Pearl River estuary, China. *Marine Chemistry*, 93(1), 21-32.
- Zhai, W., Dai, M., & Guo, X. (2007). Carbonate system and CO_2 degassing fluxes in the inner estuary of Changjiang (Yangtze) River, China. *Marine Chemistry*, 107(3), 342-356.
- Zhu, Y., Shang, S., Zhai, W., & Dai, M. (2009). Satellite-derived surface water pCO_2 and air-sea CO_2 fluxes in the northern South China Sea in summer. *Progress in Natural Science*, 19(6), 775-779. <https://doi.org/10.1016/j.pnsc.2008.09.004>

Heat storage variability in the Bay of Bengal: Importance of using altimetry data

Gayan Pathirana^{1,2,3}, Gengxin Chen¹, P. N. Ranasinghe³, Zhanlin Liang¹, Dongxiao Wang^{*,1,2}

1. State Key Laboratory of Tropical Oceanography, South China Sea Institute of Oceanology, Chinese Academy of Sciences, Guangzhou, China.
2. University of Chinese Academy of Sciences, China
3. Department of Oceanography and Marine Geology, Faculty of Fisheries and Marine Sciences & Technology, University of Ruhuna, Matara, Sri Lanka.

*dxwang@scsio.ac.cn

Abstract

Monthly average AVISO altimeter data with Argo and OFES (OGCM for the Earth Simulator) data are utilized to examine the heat storage (HS) variability in the Bay of Bengal (BoB). The preliminary analysis using Argo and OFES data suggested that the HS anomaly (ΔH) in the BoB indicates an increasing trend. Further, it is noted that the surface (upper 5m) salinity (freshening) is decreasing (increasing) while mean temperature and salinity in the upper 100 m is increasing. In general, we refer the temperature change from the surface to some reference depth level (either mixed-layer depth or a deep layer) during the estimation of the heat content based on observations (ex: Argo) or model simulations (ex: OFES) in the ocean. As there exists an uncertainty in the estimated HS by using a reference depth level due to the complexity in the BoB, it brings the importance of using the observed change of sea level for the estimation of HS. The relationship between the sea level change and the HS variability has been discussed in previous studies and the HS derived from the sea level will provide an accurate results due to: 1) its sensitivity to changes in the heating throughout the water column; 2) no assumption about a mixed-layer or a reference depth level. We have noted in the results from the preliminary analysis that the magnitude, seasonality, and trend of ΔH varies with the method utilized to compute ΔH in the BoB. Further, the monthly sea surface temperature (SST) from Extended Reconstructed SST (ERSST) over 100 years (1891-2016) highlights, that the mean SST in the BoB has increased up to $\sim 1^\circ\text{C}$ since the 1950s. Thus the noted variations in ΔH and evidence for the heat buildup in the BoB highlights the necessity of examining the variations of ΔH utilizing long-term datasets. This study is in progress and expects that the findings will provide a positive contribution to solve/understand the heat storage variability in the BoB.

1 Introduction

The Bay of Bengal (BoB) a unique region in the global ocean due to the influences from Asian monsoon, freshwater flux (precipitation + river runoff), strongly stratified surface layer, seasonally reversing monsoon

currents, and seasonal winds, is a semi enclosed basin located in the northern Indian Ocean (Figure 1a). The BoB rim-countries experiences natural hazards, such as tropical cyclones, floods, land-slides, storm surges, tsunami (not frequent), coastal erosion, and sea level rise due to global warming which brings negative impacts on ecology/economy of the countries. The region is quite complex due to its dynamics and its influence (ecological and economical) on the rim-countries brings the relative importance of scientific research studies in order to understand its dynamics.

Figure 1a: Bathymetry of the BoB, region marked in dashed square represents the area selected for the study (5 – 20° N; 80 – 95° E).

The importance of understanding the role of ocean on weather and climate under a warming environment has been pointed out in previous studies while the long term changes in SST is been considered as one of the good indicators which provides evidence of ocean warming and heat buildup. The SST in the BoB indicates an increasing trend since 1950s and it is clearly evident from the estimations of ERSST (Figure 1b). The noted increasing trend of SST in the BoB is linked with global warming and highlights the potential of heat buildup

in the region. Though the increasing trend of SST in the BoB has been well examined in previous studies, the studies has been limited to a certain depth level (ex: upper 700 m). However, the heat storage variability and its long term changes in the entire basin have not been examined in detail before. Therefore in the present study, we have examined the heat storage variability in the BoB using monthly mean data from Argo, OFES (Masumoto et al., 2004), and sea surface height anomalies (SSHA) from AVISO (January 2004 to December 2011). The heat storage anomalies (ΔH) estimated using the SSHA, is capable of producing a clear picture for the entire basin in terms of thermal expansion of the water, without limitations of depth. Further, we have considered the effect from stratification on upper-ocean heat content.

Figure 1b: Monthly sea surface temperature (SST) variability estimated from ERSST (1891 – 2016) for the study area. Dashed black line represents the threshold SST (28° C) for the tropical cyclone genesis.

2 Data and Methods

2.1 Estimation of mixed-layer ΔH

We utilize OFES and gridded Argo data to estimate the monthly mixed-layer ΔH in the BoB from 2004 to 2010. Mixed-layer depth (MLD) is defined as the depth where the density change is equivalent to 0.8 °C temperature change from the surface value at 1 m (Rao and Sivakumar, 2000), which can be estimated using Eq. (1). This criterion is based on temperature with a fixed threshold ($dT=0.8^\circ\text{C}$) and considers both temperature and salinity effects on stratification.

$$\text{MLD} = \text{depth where } [\theta = (\theta_{(T,S,P)} + \theta_{(T+dT,S,P)})] \quad (1)$$

Where S is salinity, T is temperature at 1 m depth, P is water pressure and θ is the water density. The heat storage anomaly in the BoB is estimated using (Eq. 2), which is based on mixed-layer temperature and MLD.

$$\Delta H = \rho C_p h \Delta T \quad (2)$$

Where ΔH is the stored heat anomaly per unit area (Jm^{-2}), ρ is the sea water density ($\rho = 1024 \text{ kgm}^{-3}$), C_p is the specific heat capacity of sea water ($C_p \approx 4000 \text{ Jkg}^{-1} \text{ K}^{-1}$), ΔT is the vertically averaged temperature anomaly, which is estimated by removing the 7 year mean ($\Delta T = T - T_{\text{mean}}$), and h is the reference depth. We have considered two reference depth levels as, $h = \text{MLD}$ and $h = 250\text{m}$ for the study in order to examine the sensitivity of ΔH to the reference depth level.

2.2 Estimation of ΔH using SSH data

If there is a change in sea level due to warming of the oceans (thermal expansion), there is a possibility to estimate the heat content in the ocean in terms of observed sea level. Gill and Niiler (1973), pointed out that the changes in the ocean heat storage will influence the ocean topography due to the change in water density in terms of thermal expansion. Thus it leads for a direct relationship between sea level change and heat storage considering the relationship between heat storage and density change (Chambers and Tapley, 1997). Therefore the ΔH is estimated using Eq. (3).

$$\Delta H = (\rho C_p) / \alpha \Delta \eta \quad (3)$$

Where α is the thermal expansion coefficient of sea water and $\Delta \eta$ is the change in sea level. This relation provides a better ΔH result than the ΔH estimated using Eq. (2), as the sea level is sensitive to the changes in the heat content throughout the water column and can detect heat storage changes in the deepest layers.

However, the sea level change ($\Delta \eta$) recorded in the satellite observations is not solely due to the thermal expansion of sea water.

$$\Delta \eta = \Delta \eta_{\text{AVISO}} + \epsilon \quad (4)$$

Therefore the $\Delta \eta$ recorded in altimeter products is consisted with errors (ϵ) such as: instrument errors, tides, inverted barometer effect, change in $\Delta \eta$ due to the variability of salinity, and changes due to the barotropic response of the ocean to wind (propagating waves, surface currents, meso-scale eddies...etc.). Thus it is important to remove the other influencing factors from altimetry data except the thermal expansion impact in order to enhance the accuracy of the results. In the present study we utilize AVISO L4 monthly data from 2004 to 2010 obtained from Physical Oceanography Distributed Active Archive Center (PODAAC) (AVISO, 2011). The dataset is corrected for instrument errors, tides, and inverted barometer effects during the data processing and we apply the standard corrections to remove other the errors mentioned above.

2.3 Estimation of stratification / freshwater influence

The stratified upper-layers in the tropical oceans tends to trap more heat energy and maintain a warmer surface-layer than the deeper layers. The stratification in the BoB is heavily depending on the vertical distribution of salinity rather than the temperature. The freshwater influx to the BoB during summer to post-summer monsoon causes a less dense surface water layer. During this time there is subsurface water intrusion from the Arabian Sea (AS) into the BoB which brings high salinity waters and influence the increase of subsurface salinity. Thus it leads to the strengthening of the vertical stratification, which increases the potential for the heat accumulation in the upper waters in the BoB. Hence we estimate the influence of stratification utilizing available CTD data from National Oceanographic Data Center (NODC) with OFES and Argo data sets. The freshwater content (H_{FW}) in the BoB, is calculated using Eq. (5)

with respect to a reference salinity (S_{ref}), following Mahadevan et al., (2016).

$$H_{FW} = \int_{Z(S_{ref})}^0 \frac{(S(z) - S_{ref})}{S_{ref}} dz \quad (5)$$

where, $Z(S_{ref})$ is the depth of the isohaline S_{ref} . The selected isohaline for S_{ref} must lie well below the freshwater influenced layer (ex: $S_{ref}=35$ psu). The water stability (E) in terms of the buoyancy frequency is estimated using Eq. (6)

$$E = \frac{g}{\rho^*} \left[\frac{d\rho}{dz} - \frac{g\rho^*}{C^2} \right] \quad (6)$$

where g is the acceleration due to gravity (9.8 ms^{-2}), ρ is the sea water density (kgm^{-3}), ρ^* is the mean density in the water column dz , z is the depth (m), and C is the sound velocity (ms^{-1}) in the sea water (Chen and Millero, 1977).

3. Results and Discussion

Figure 2 illustrates the average MLD and standard deviation in the BoB, estimated from Argo and OFES data (2004 – 2010). The mean MLD varies between 15 – 45 m and indicates the presence of a deeper (shallow) mixed-layer in the southeastern (northern) part of the BoB (Figure 2a and b). However the estimated MLD from OFES data indicates relatively larger values compared to that estimated from Argo. The differences between the estimated MLDs is evident in the standard deviation (Figure 2c and d) and it suggests the presence of a larger MLD variability in the southern BoB from the Argo estimation while the northwestern BoB from the OFES estimation.

Figure 2: Average MLD (a) Argo, (b) OFES, and its standard deviation (c) Argo, (d) OFES in the BoB.

The difference between the estimated MLD from two datasets is evident in the correlation values (r), while it is noted the presence of a strong positive correlation in the central BoB (Figure 3a). Though there are differences in the estimated MLDs, the ΔH

calculated from two different datasets indicates a strong positive correlation for the entire BoB with $r > 0.6$ at 95% significance level (Figure 3b). However, the accuracy of the estimated MLD depends on the choice of MLD criterion and the vertical resolution of the data (Pathirana et al., manuscript submitted, 2018).

Figure 3: Correlations between (a) MLDs estimated from Argo and OFES, and (b) ΔH estimated from Argo and OFES data.

Seasonality of ΔH is evident from both datasets and indicates the evidences of positive ΔH (negative ΔH / releasing) during monsoon transition (winter) periods (Figure 4). The presence of an increasing trend in the ΔH is noted from 2004 to 2010 and points out the existence of a potential for the heat buildup in the BoB. Thus the noted increasing trend in SST after 1950s (Figure 1b) and in ΔH estimated considering $h=MLD$ (Figure) provide clear evidences of heat buildup in the upper-layers of the BoB and highlights the necessity of examining the long term changes in basin scale ΔH in order to explain the heat buildup in the region with higher accuracy.

Figure 4: Seasonal variability of fJH in the BoB during 2004 – 2010 from Argo (red line) and OFES (blue line). The dashed black line represents the linear trend of ΔH .

Though there are uncertainties in the estimated ΔH using Argo and OFES data due to the coarse vertical resolution and choice of MLD, still it provides enough evidences to understand the mixed-layer heat storage variability in the BoB. However it depends on the criteria we utilized to estimate the MLD and also on the dataset. Due to such uncertainties, the estimated ΔH varies according to the referred depth level and it does not provide a clear picture for an entire basin as we do not have observations below 2000m depth in the ocean.

In order to examine the variability of ΔH due to the referred depth level in the BoB, we have estimated ΔH using $h=MLD$, $h=250\text{m}$, and using SSHA and the

results are presented in Figure 5. It is evident in the results that the magnitude, seasonality, and trend varies with the method utilized to compute ΔH in the BoB. The ΔH based on MLD clearly follows the seasonal cycle of SST and MLD with peaks during pre-summer (April-May) and post-summer (October-November) in the BoB. Though the peak of ΔH during pre-summer is evident at any reference depth considered to compute ΔH , the peak of ΔH during post-summer shifts to winter with increasing the reference depth level. However, as we have considered only 7 years, here we do not conclude the reasons for the noted change in ΔH in the BoB.

Figure 5: Variability of ΔH in the BoB: (a) ΔH based on h =MLD from Argo (red line) and OFES (blue line), (b) ΔH based on h =250m from Argo, and (c) ΔH estimated from SSHA. In the figure dashed lines (black and red) represent the noted trend. The shaded area (light blue) indicates the standard deviation of ΔH .

4 Summary

In this study we are focusing on the heat storage variability in the BoB and the importance of using altimetry data to estimate the ΔH . The heat changes in the ocean are directly linked with the changes in the density (thermal expansion) and enables us to derive the heat storage from the observed sea level change. We have noted in the results that the magnitude, seasonality, and trend varies with the method utilized to compute ΔH in the BoB. The relation between the heat storage and the sea level anomalies has been discussed in the literature, but few studies have focused on the Indian Ocean. In the present study, we have selected the BoB which is more complex due to its unique characteristics and we expect to comparatively examine the variations in the heat storage in the BoB by using SSHA (AVISO) along with Argo and OFES data. Then discuss how accurately they represent the heat buildup in the BoB. Further we will examine the role of

freshwater layer in terms of stratification and sea level change in seasonal timescales.

The noted variations in ΔH and evidences for the heat buildup in the BoB highlights the necessity of examining the variations in ΔH utilizing long-term datasets. This study is in progress and expect that the findings will provide a positive contribution to solve/understand the dynamics in the BoB. Our goal is to find a method for the accurate estimation of heat storage variability (seasonal / long-term) in the BoB. The full research paper will be available under the same title in this article or can be accessed through writing to the corresponding author.

Acknowledgement

This work is supported by the Major State Research Development Program of China (2016YFC1402603), NSFC (grants 41476011 and 41521005), XDA (grant 11010102) and the Youth Innovation Promotion Association of CAS (2017397). The Argo data were collected and made freely available by the International Argo Program and the national programs that contribute to it (www.argo.ucsd.edu, argo.jcommops.org). Authors would like to thank China-Sri Lanka Joint Center for Education & Research (CSL-CER) for their support.

References

Antonov, J. I., 2002: Steric sea level variations during 1957-1994: Importance of salinity. *Journal of Geophysical Research*, 107(C12), doi:10.1029/2001JC000964.

AVISO. 2011. AVISO Level 4 Absolute Dynamic Topography for Climate Model Comparison. Ver. 1. PO.DAAC, CA, USA. Dataset accessed [2018-02-15] at <http://dx.doi.org/10.5067/DYNT0-1D1M1>.

Chambers, D. P., Tapley, B. D. and Stewart, R. H., 1997: Long-period ocean heat storage rates and basin-scale heat fluxes from TOPEX. *Journal of Geophysical Research*, 10525-10533.

Chen, C. T. and Millero, F. J., 1977: Speed of sound in seawater at high pressures. *Journal of the Acoustical Society of America*, 62, 1129-1135, doi:10.1121/1.381646.

Mahadevan, A., Jaeger, G. S., Freilich, M., Omand, M., Shroyer, E. L. and Sengupta, D., 2016: Freshwater in the Bay of Bengal: Its fate and role in air-sea heat exchange. *Oceanography*, 29(2), 72-81, doi:10.5670/oceanog.2016.40.

Masumoto, Y., Sasaki, H., Kagimoto, T., Komori, N., Ishida, A., Sasai, Y., Miyama T., Motoi, T., Mitsudera, H., Takahashi, K., Sakuma, H., and Yamagat, T., 2004: A fifty-year eddy-resolving simulation of the world ocean—Preliminary outcomes of OFES (OGCM for the Earth Simulator). *Journal of Earth Simulator*, 35-56.

McDougall, T. J., 1987. Neutral Surfaces. *Journal of Physical Oceanography*, 17, 1950-1964.

Murty, V. S. N., Sarma, Y. V. B. and Rao, D. P., 1996: Variability of the oceanic boundary layer characteristics in the northern Bay of Bengal during MONTBLEX-90. *Earth Planet Science*, 105(1), 41-61.

Pathirana, G., Wang, D., Abeyratne, M.K., Priyadarshana, T., Yun, Q., and Chen, G., 2018: Effect of seasonal barrier-layer on mixed-layer heat budget in the Bay of Bengal. *Ocean Dynamics*, in review.

Rao, R. R. and Sivakumar, R., 2000: Seasonal variability of near-surface thermal structure and heat budget of the mixed layer of the tropical Indian Ocean from a new global ocean temperature climatology. *Journal of Geophysical Research*, 105, 995-1015, doi:10.1029/1999JC900220.

Rebert, J. P., Donguy, J. R. and Eldin, G., 1985: Relations Between Sea Level, Thermocline Depth, Heat Content, and Dynamic Height in the Tropical Pacific Ocean. *Journal of Geophysical Research*, 90(C6), 11719-11725.

Thompson, K. R. and Demirov, E., 2006: Skewness of sea level variability of the world's oceans. *Journal of Geophysical Research*, 111(C05005), doi:10.1029/2004JC002839.

Is the reconstruction of the storm surge in 1906 in the German Bight with reanalyses possible?

Elke M. I. Meyer¹, Ralf Weisse¹

1. Helmholtz-Zentrum Geesthacht Zentrum für Material- und Küstenforschung GmbH, Max-Planck-Straße 1, 21502 Geesthacht, Germany

Abstract

Long-term atmospheric reanalyses provide an opportunity to reconstruct historical storm surge events. In this study, the historical storm surge from March, 13th 1906 in the German Bight is investigated. For this event, a tide surge model was run, forced by tidal constituents from the FES2004 tide model and by wind and pressure fields from ensemble members of the 20th Century Reanalysis project and the ERA-CLIM project. Initial results show that while winds from a few ensemble members produce surges close to the observations, generally wind fields appear to be too weak to reproduce observed water level extremes during this event.

Introduction

On March, 13th 1906, a severe storm surge hit the German North Sea coast. For many East Frisian Islands in the southern part of the German Bight, it still represents one of the highest ever observed extreme (Deutsches Gewässerkundliches Jahrbuch, 2014). So far, a detailed analysis was not possible because of missing atmospheric data to run corresponding storm surge simulations. Recently atmospheric data from long reanalysis projects became available (e.g. the Twentieth Century Reanalysis Project version 2c (20CR-V2c, Compo et al., 2011) or the ERA-CLIM & ERA-CLIM2 projects (Buizza et al 2017, Laloyaux et al. 2016) that provide an opportunity to model historical storm surge events. Here we assess the capability of using such reanalysis data in reproducing the extreme storm surge event in 1906.

The projects assimilate at least historical surface pressure in their global models. Data for assimilation are for example from records by weather stations, light vessels or from logbooks from ship cruises and collected within the International Surface Pressure Databank (ISPD, Cram et al. 2015). The 20CR-V2c reanalysis data, starting in 1851, is composed of 56 ensemble members. It provides mean sea level pressure and wind speed at a 6-hourly interval with a horizontal resolution of about 200km. The ERA-CLIM-project provides the 3-hourly intervals products of ERA-20c and CERA-20c with 10 ensemble members of which the horizontal resolution is approximately 125 km since 1900 (Buizza et al 2017, Laloyaux et al. 2016).

Weather situation

A low-pressure system moved from England over the central North Sea to Scandinavia. In the night from March, 12th to March 13th 1906, the core of the system was located over central Sweden and an anticyclone was established over Southwest Europe. Strong north-westerly winds were prevailing over most of the North Sea pushing waters towards the German Bight and the East Frisian Islands.

Model-run

Data from the FES2004 global tide model (Lyard et al .2006) were used as lateral boundary conditions for the tide-surge model. Wind and pressure fields from reanalyses discussed above were used to drive the model.

The tide-surge model used is the hydrodynamic model TRIM-NP (Pätsch et al, 2017) from which hourly tide-surge levels were stored. The TRIM-NP model is a three-dimensional finite difference Tidal Residual Intertidal Mudflat (TRIM) model developed by Casulli and Cattani (1994) and then nested and parallelised by Kapitza (2008). The model run is a three-level nested integration with spatial resolutions of 12.6, 6.4, 3.2 and 1.6 km. The model covers the area from 20°W to 30°E and from 42°N to 65°N, with a resolution of 12.6km. The higher resolution grids are located in the Southern North Sea. All model runs were used in a barotropic mode and started from February, 1st 1906.

Results

Water levels from Norderney (East Frisian Island; 53° 42'N, 7° 9'E) and Cuxhaven (Elbe estuary; 53°52'N, 8°42'E) were analysed. The observed water level was 3.95m above sea level (ASL) for Norderney (Figure 1) and 4.51m ASL for Cuxhaven (Figure 2) on March, 13th 1906. During the storm surge all ensemble member runs simulate water levels less than the observed maxima for Norderney and Cuxhaven. A few ensemble members using winds from the 20CR-V2c run are close to the observed water levels. The model run without wind forcing shows a high tide during the storm event (blue line). The analysis of the astronomical tides indicates a very high tide during this event.

The analysis of the wind speeds in the North Sea (latitude: 54°, longitude: 6°) show only moderate wind speeds in the reanalyses. Non-digitized weather records from the German Weather service suggest that wind speeds in the area probably have exceeded 10 Beaufort (pers. comment Birger Tinz, German Weather Service).

Figure 1: Simulated water levels at Norderney using wind fields from different reanalysis products (colours) and observed highest level during the event (black).

Figure 2: Simulated water levels at Cuxhaven using wind fields from different reanalysis products (colours) and observed highest level during the event (black)

During the event, ensemble members from the 20CR-V2c run simulate higher wind speed than the ensemble members from ERA-20c and CERA-20c (Figure 3). The wind speed from all ensemble members is lower than 10 Bft (black line in Figure 3).

Summary

While long atmospheric reanalyses provide an opportunity to study historic storm surge events none of the available wind fields were able to reconstruct the observed water levels that occurred during an event in March 1906 along parts of German North Sea coast.

Figure 3: Simulated wind speed at the location 54°N and 6°E from all reanalysis products (colours) and 10Bft-threshold (black).

Comparison with observations suggest, that wind speeds in the reanalyses were too moderate to recover the unusual character of this extreme event. Additionally, other factors such as insufficient spatial resolution, neglecting of gusts, or unresolved bathymetric effects may further contribute to the observed discrepancies. Assimilating additional pressure data that had not been digitized during the time of the production of the reanalyses may improve the reanalyses products. As quite a few additional data have now been introduced into the ISPD it may well be that with newer reanalysis products the situation may substantially improve.

Acknowledgement

"Support for the Twentieth Century Reanalysis Project dataset is provided by the U.S. Department of Energy, Office of Science Innovative and Novel Computational Impact on Theory and Experiment (DOE INCITE) program, and Office of Biological and Environmental Research (BER), and by the National Oceanic and Atmospheric Administration Climate Program Office."

Reference

Buizza, R., S. Brönnimann, L. Haimberger, P. Laloyaux, M.J. Martin, M. Fuentes, M. Alonso-Balmaseda, A. Becker, M. Blaschek, P. Dahlgren, E. de Boisseson, D. Dee, M. Doutriaux-Boucher, X. Feng, V.O. John, K. Haines, S. Jourdain, Y. Kosaka, D. Lea, F. Lemarié, M. Mayer, P. Messina, C. Perruche, P. Peylin, J. Pullainen, N. Rayner, E. Rustemeier, D. Schepers, R. Saunders, J. Schulz, A. Sterin, S. Stichelberger, A. Storto, C. Testut, M. Valente, A. Vidard, N. Vuichard, A. Weaver, J. While, and M. Ziese, 2018: The EU-FP7 ERA-CLIM2 Project Contribution to Advancing Science and Production of Earth System Climate Reanalyses. Bull. Amer. Meteor. Soc., 99, 1003–1014, <https://doi.org/10.1175/BAMS-D-17-0199.1>

Casulli V, Cattani E (1994) Stability, accuracy and efficiency of a semi-implicit method for three-dimensional shallow water flow. Comput Math Appl 27(4):99–112

Compo, G.P., J.S. Whitaker, P.D. Sardeshmukh, N. Matsui, R.J. Allan, X. Yin, B.E. Gleason, R.S. Vose, G. Rutledge, P. Bessemoulin, S. Brönnimann, M. Brunet, R.I. Crouthamel, A.N. Grant, P.Y. Groisman, P.D. Jones, M. Kruk, A.C. Kruger, G.J. Marshall, M. Maugeri, H.Y. Mok, Ø. Nordli, T.F. Ross, R.M. Trigo, X.L. Wang, S.D. Woodruff, and S.J. Worley, 2011: The Twentieth Century Reanalysis Project. *Quarterly J. Roy. Meteorol. Soc.*, 137, 1-28. DOI: 10.1002/qj.776.

Cram, T. A., Compo, G. P., Yin, X. , Allan, R. J., McColl, C. , Vose, R. S., Whitaker, J. S., Matsui, N. , Ashcroft, L. , Auchmann, R. , Bessemoulin, P. , Brandsma, T. , Brohan, P. , Brunet, M. , Comeaux, J. , Crouthamel, R. , Gleason, B. E., Groisman, P. Y., Hersbach, H. , Jones, P. D., Jónsson, T. , Jourdain, S. , Kelly, G. , Knapp, K. R., Kruger, A. , Kubota, H. , Lentini, G. , Lorrey, A. , Lott, N. , Lubker, S. J., Luterbacher, J. , Marshall, G. J., Maugeri, M. , Mock, C. J., Mok, H. Y., Nordli, Ø. , Rodwell, M. J., Ross, T. F., Schuster, D. , Srncic, L. , Valente, M. A., Vizi, Z. , Wang, X. L., Westcott, N. , Woollen, J. S. and Worley, S. J. (2015), The International Surface Pressure Databank version 2. *Geosci. Data J.*, 2: 31-46. doi:10.1002/gdj3.25

Deutsches Gewässerkundliches Jahrbuch, Küstengebiet der Nordsee Landesamt für Landwirtschaft, Umwelt und ländliche Räume Schleswig-Holstein Flintbek 2014, ISSN 2193-6374, www.dgj.de

Kapitza H (2008) Mops: a morphodynamical prediction system on cluster computers. In: Laginha JM, Palma M, Amestoy PR, Dayde M, Mattoso M, Lopez J (eds) High performance computing for computational science-VECPAR 2008, Lecture Notes in Computer Science, Springer, pp 63-68

Laloyaux, P., Balmaseda, M., Dee, D., Mogensen, K. and Janssen, P. (2016), A coupled data assimilation system for climate reanalysis. *Q.J.R. Meteorol. Soc.*, 142: 65-78. doi:10.1002/qj.2629

Lyard, F., Lefèvre, T., Letellier and O. Francis. Modelling the global ocean tides: a modern insight from FES2004, *Ocean Dynamics*, 56, 394-415, 2006.

Pätsch, J. ; Burchard, H.; Dieterich, C.; Grawe, U. ; Groger, M.; Mathis, M.; Kapitza, H.; Bersch, M.; Moll, A.; Pohlmann, T.; Su, J.; Ho-Hagemann, HTM; Schulz, A.; Elizalde, A.; Eden, C.. "An evaluation of the North Sea circulation in global and regional models relevant for ecosystem simulations". *Ocean Modelling* 116. (2017): S. 70-95. doi: 10.1016/j.ocemod.2017.06.005

This page intentionally left blank

This page intentionally left blank

Editorial

Jose Santos, Lipin Yin.....	1
Introduction to the CLIVAR Summer School in China	
Jing Li.....	2
High-resolution projections of sea-level change processes on the Northwestern European Shelf	
Tim Hermans, Aimée Slangen, Jonathan Tinker, Matthew Palmer, Bert Vermeersen.....	6
Understanding the Recent Hawaii High Sea Level Event	
Xiaoyu Long, Matthew Widlansky.....	10
Assessment of the intraseasonal Kelvin wave activity during Central Pacific and Eastern Pacific El Niño events	
Jonathan Cedeño ¹ , Óscar Pizarro, Aldo Montecinos.....	13
The Nigerian's Gulf of Guinea coastal evolution and response to multi-scale forcing	
Dada, Olusegun Adekoyejo.....	17
Formation and types of thermal inversion in the Bay of Bengal	
K M Azam Chowdhury, Wensheng Jiang, Guimei Liu, Md. Kawser Ahmed, Sirajuddin Md. Babar Chowdhury, Nuruddin Md. Khaled Chowdhury.....	20
Modeling of the Biogeochemical Cycle and Dispersion of Nitrogen off Lake Idku Outlet, Abu Qir Bay, Egypt	
Amr T. Salama, Samir M. Nasr, Ahmed E. Rifaat.....	24
Sediment Transport and Coastal Erosion Processes in Southern Spain under Climate Change Scenarios	
Andrea Lira-Loarca, Manuel Cobos, Agustín Millares, Asunción Baquerizo, Miguel A. Losada.....	27
Research on the Energy and Carbon Dioxide Fluxes in the Atmosphere-Ocean Interface of the Tropical Coastal Ocean	
Yusri Yusup, John Stephen Kayode, Abbas F.M. Alkarkhi.....	30
Heat storage variability in the Bay of Bengal: Importance of using altimetry data	
Gayana Pathirana, Gengxin Chen, P. N. Ranasinghe, Zhanlin Liang, Dongxiao Wang.....	33
Is the reconstruction of the storm surge in 1906 in the German Bight with reanalyses possible?	
Elke M. I. Meyer, Ralf Weisse.....	38

The CLIVAR Exchanges is published by the International CLIVAR Project Office ISSN No: 1026-0471

Editor: Jose Santos

Guest editor: Liping Yin

Layout and Designing: Harish J. Borse, ICMPO at IITM, Pune, India

Note on Copyright

Permission to use any scientific material (text as well as Figs) published in CLIVAR Exchanges should be obtained from the authors. The reference should appear as follows: Authors, Year, Title. CLIVAR Exchanges, No.pp. (Unpublished manuscript).

Production, Printing and Mailing of this issue by ICMPO with the support of IITM and Ministry of Earth Sciences, India.

WCRP is sponsored by the World Meteorological Organization, the International Council for Science and the Intergovernmental Oceanographic Commission of UNESCO.

Contact:

Executive Director, ICPO
First Institute of Oceanography, SOA,
6 Xianxialing Road, Laoshan District, Qingdao 266061, China
icpo@clivar.org
http://www.clivar.org

Ministry of Earth Science

Please recycle this newsletter by passing on to a colleague or library or disposing in a recognised recycle point