To the Editor:

The authors describe an advanced vertebroplasty technique in a very complex patient. This scenario provides an opportunity to discuss relevant issues regarding osteoporotic compression fractures (OCF) and subsequent management. Why undertake the challenge for "Rescue" in this medically compromised patient population? Recent literature testifies to an impaired pulmonary status (12), loss of function (3), and increased mortality rate versus age matched controls (4). Furthermore, even "silent" fractures lead to compromised function (5). These facts presented behind the safety of a medical journal still do not match the dramatic intensity of the singular patient presentation in our offices. This pain described as "stabbing, sharp" with any transitional motion leads to significant debilitation and demoralization (6). "I would rather die than live like this" is a commonly expressed sentiment. Documentation of this patient's clinical condition with Verbal Pain Scores, medication record, and functional impairment would "paint" a more accurate picture of the argument for a highly invasive treatment. I wasn't aware that treatment of painful vertebral fractures with multivitamins and calcium constitutes "aggressive" non-surgical management.

The other pertinent issue is the actual OCF treatment in the fact of neurologic compromise. What are the risks of performing vertebroplasty in a fracture with demonstrated retropulsion and "cord compromise?" The potential for spinal cord or nerve injury is well-publicized with more frequent case reports (7-9). Cement extravasation may occur by various routes, including basivertebral veins, segmental veins, or cortical defects (10). Therefore, the injectate can cause local and/or distant complications depending on the carrier pathway. Recent dramatic case reports have highlighted the devastating results of "cement emboli" (11, 12) and leads me to question the wisdom of intervertebral "air" injection. While retropulsion is not specifically addressed in the literature, other identifiable risk factors for cement migration include: target level above T-7, larger injectate volume, and "injector" use (13). The authors wisely immediately sought Neurosurgical opinion. However, the examining surgeon obviously felt the risks of a surgical approach outweighed the potential benefits. The authors

then scheduled the procedure with the Neurosurgical availability demonstrating optimal perioperative planning.

The anatomic challenge of vertebroplasty in the vertebrae plana is well-documented and initial reports advised against treatment in this setting. More recently, the literature has given some guidance in these challenging fractures (13, 14). The authors describe a thoughtful ingenious approach with the guide-wire technique. In addition, the utilization of a bilateral approach highlights due caution with angle entry in placing the vertebroplasty needles. The wrong angle would lead to cord violation! Needle placement was optimally placed in this case in the anterior third of the vertebra. Next, "gradual" cement injection was performed with "live" fluoroscopy. PLEASE NOTE GRADUAL! Most complications from vertebroplasty are "cement" related! These include: undesirable migration (vascular-emboli!) or posterior (cord compromise) and allergic reaction.

Lane et al (15) retrospectively examines the prognostic importance of fluid and gas-filled clefts on vertebroplasty and concludes that greater pain relief is accomplished with "cleft-filling." The peer-reviewed analgesic efficacy of vertebroplasty has been well established and lead me to question the introduction of a potentially dangerous variable (air injection) in a high-risk situation. Had other more conservative measures been explored? We have had good experience with the Gray Ramus nerve block and denervation at our institution and this procedure does not carry the same potential risk of cord injury (16). Neuroaxial narcotic therapy might also be a less invasive option. The intervention planned should be balanced against the operator's skill, procedural risk, and expected outcome. The authors are to be commended for their technical skill, ingenuity, and prudence for neurosurgical back up.

REFERENCES

1. Leech J, Dulberg C, Kellie S, Pattee L et al. Relationship of lung function to severity of osteoporosis in women. *Am Rev Respir Dis* 1990; 141:68-71.
2. Schlaich C, Minne HW, Bruckner T et al. Reduced pulmonary function in patients with spinal osteoporotic fractures. *Osteoporos Int* 1998; 8:261-267.
3. Lyles KW, Gold DT, Shipp KM et al. Association of osteoporotic vertebral compression fractures with impaired functional status. *Am J Med* 1993; 94:595-601.
4. Kado D, Browner WS, Palermo L et al. Vertebral fractures and mortality in older women: a prospective study. Study of Osteoporotic Fractures Research Group. *Arch Intern Med* 1999; 159:1215-1220.
5. Nevitt M, Ettinger B, Black DM et al. The association of radiographically detected vertebral fractures with back pain and function: A prospective study. *Ann Intern Med* 1998; 128:793-800.
6. Silverman SL. The clinical consequences of vertebral compression fracture. *Bone* 1992; 13:S27-S31.
7. Harrington K. Major neurologic complications following percutaneous vertebroplasty with methylmethacrylate. A case report. *JBJS* 2001; 83:1070-1073.
8. Lee BJ, Lee SR, Yoo TY. Paraplegia as a complication of percutaneous vertebroplasty with polymethylmethacrylate: A case report. *Spine* 2002; 27:E419-E422.
9. Ratliff J, Nguyen T, Heiss J. Root and spinal cord compression from methylmethacrylate vertebroplasty. *Spine* 2001; 26:E300-E302.
10. Yeom JS, Kim WJ, Choy WS et al. Leakage of cement in percutaneous transpedicular vertebroplasty for painful osteoporotic compression fractures. *Bone Joint Surg Br* 2003; 85:83-89.
11. Scroop R, Eskridge J, Britz GW. Paradoxical cerebral arterial embolization of cement during intraoperative vertebroplasty: A case report. *AJNR Am J Neuroradiol* 2002; 23:868-870.
12. Chen HL, Wong CS, Ho ST et al. A lethal pulmonary embolism during percutaneous vertebroplasty. *Anesth Analg* 2002; 95:1060-1062.
13. Ryu KS, Park CK, Kim MC et al. Dose-dependent epidural leakage of polymethylmethacrylate after percutaneous vertebroplasty in patients with osteoporotic vertebral compression fractures. *J Neurosurg* 2002; 96:S56-S61.
14. Peh WC, Gilula LA, Peck DD. Percutaneous vertebroplasty for severe osteoporotic vertebral body compression fractures. *Radiology* 2002; 223:121-126.
15. Lane JI, Maus TP, Wald JT et al. Intravertebral clefts opacified during vertebroplasty: pathogenesis, technical implications, and prognostic significance. *AJNR Am J Neuroradiol* 2002; 23:1642-1646.
16. Chandler GS. Gray ramus communicans nerve block: Novel treatment approach for painful osteoporotic vertebral compression fracture. *SMJ* 2001; 94:387-393.

Gilbert Chandler III, MD
 Specialty Clinics of Georgia
 1240 Jesse Jewell Parkway, Suite 300
 Gainesville GA 30501
 E-mail: sewell87@aol.com