In Response: Glaser et al

We appreciate the letter to the editor by Dr. Ruan and colleagues. They respect the gravity of this issue. We have received worldwide correspondence echoing similar concerns. Unfortunately, ongoing dissemination of the "unsafe triangle" technique will be challenging to stop. In the interim, cases of paraplegia will continue to accumulate. In just a few short months following our publication, one of the authors has received records regarding 3 other cases of paraplegia following "unsafe triangle" transforaminal epidural steroid injections performed at the following levels: right L5-S1, left L2-3 and left T12-L1. All 3 of these are recent, dating from spring, 2008, and the other 2 occurring in the spring of 2010. In fact, the author was alerted of the second complication by an email from the interventional pain physician

who performed the procedure. He was complimentary of the article and went on to state "I wish I would paid attention to it earlier. Today, using the safe triangle approach, I had an unfortunate complication of paralysis" The "unsafe" triangle TFESI has garnered rapid growth and market share as an interventional procedure. This acceptance among interventionalists is unprecedented and is now a "bread and butter procedure" as noted by Dr. Ruan. Dr. Ruan, however, aptly states that this "unsafe triangle" technique is "dangerous and harmful." We applaud his courage and willingness to change his practice. This is a particularly difficult process for any seasoned and experienced practitioner. Our hope is that other physicians take similar notice and action, as Dr. Ruan. Practitioners must understand that the TFESI

was introduced as a conceptual improvement over the interlaminar ESI. This “unsafe triangle” approach gained traction among providers, before the procedure was fully vetted and before solid efficacy studies were available. The “unsafe triangle” TFESI went from a “drawing on a napkin” to “primetime” rapidly. Since the procedure was perceived as ‘safe’, complications were blamed on the provider. This was a “Black Swan event” and early complications did not lead practitioners to rethink this procedure. Despite this “Black Swan event,” most practitioners continue to shrug their shoulders and carry on with a “business as usual” approach.

A delayed understanding of the risks of the “unsafe triangle” TFESI may have occurred due to a low base rate of complications. When the volume of procedures is low, each complication appears rare and isolated with a low base rate. In this situation, practitioners were held accountable for malpractice. Now, thousands of practitioners worldwide are performing this procedure millions of times annually. This large volume multiplied by any base rate yields a sizable number of catastrophic complications. As the number of cases of paraplegia accumulate, practitioners can no longer only blame each other for negligence. We have argued that these events fundamentally point to the “unsafe triangle” TFESI as the root cause of the problem.

A recent article by Murthy et al, evaluating the location of radicular feeder vessels on angiography confirms the risks of the “unsafe triangle.” The International Spine Intervention Society continues to teach the “unsafe triangle” technique. A recent flyer for their bioskills labs suggests that they are limiting the “unsafe triangle” approach to the L4, L5, and S1 spinal nerves/nerve roots. This modest paradigm shift is not sufficient to reduce complications. Endorsing the “unsafe triangle” approach for the L4, L5, S1 levels will not mitigate the dangers of this approach: the vast majority of TFESIs are performed at L4, L5, and S1.

“Unsafe triangle” TFESIs can harm a patient, if this patient has an Arteria Radicularis Magna in the lower lumbar foramina or if the procedure is performed at a higher level. Paraplegia is a devastating singular event, caused by an interruption of the blood supply through this important vessel. Interruption can occur with embolization, cannulation, intimal flaps, vasospasm, and arterial injury. This complication can only be avoided by complete abandonment of the “unsafe triangle” approach.

These catastrophic complications were “Black Swan events” and should have prompted further inquiry into the root cause. We believe the root cause is the “unsafe triangle.” This approach should no longer be taught or

practiced. We contend that a better approach would be to use the Kambin triangle approach. This location, according to Murthy et al, is free of the ARM. Conceptually, the risk of paraplegia should be negligible and targeting should be preserved.

Dr. Ruan raises a valid point about reducing radiation exposure, by using bony landmarks. The posterior elements are important landmarks for gauging depth in Kambin triangle TFESIs. The needle can rest on the infero-lateral portion/base of the superior articular process dorsally bounding the foramen. This bony landmark can be used in anteroposterior, oblique, and lateral projections. The consistent location of this bony landmark may obviate the need for multiple lateral images. This bony landmark may be helpful, even when parallax exists. The most important aspect of this bony landmark in Kambin triangle TFESIs is that the needle avoids the radicular feeder vessels and the nerve root.

In conclusion, we condemn the efforts to salvage a technique that clearly puts patients in harm’s way when there are safe and effective alternatives. These alternatives include most promisingly, the Kambin triangle approach, which spatially places the bevel of a sharp needle as far as possible from the arterial supply to the spinal cord while still allowing access to the foramen. Our hope is that experienced individual practitioners abandon the “unsafe triangle,” as has Dr. Ruan. We also hope to bring awareness of this risk to fellows and residents to speed the adaptation and acceptance. We have been asked to present this compelling evidence at grand rounds of one prominent interventional pain management fellowship. There was great interest in the root cause of these complications and our approach to mitigate, if not eliminate, the risk. Interestingly, one of the fellows asked who would support him in court if he had a complication after a Kambin triangle TFESI. We hope that in the near future the same question will be asked about the “unsafe triangle” TFESI.

Scott E. Glaser, MD
Pain Specialists of Greater Chicago
7055 High Grove Blvd. Ste. 100
Burr Ridge, IL 60527
E-mail: sglaser@painchicago.com

Rinoo V. Shah, MD
Guthrie Clinic, Interventional Pain Management
31 Arnot Road
Horseheads, New York 14845
E-mail: rinoo_shah@yahoo.com