COMPARATIVE ANALYSIS OF THE FUNCTIONAL OUTCOME OF ARTHROSCOPIC ANTERIOR CRUCIATE LIGAMENT RECONSTRUCTION USING QUADRUPLED HAMSTRING GRAFT FIXED WITH BIO ABSORBABLE INTERFERENCE SCREW AGAINST TITANIUM INTERFERENCE SCREW

Orthopaedics

Dr. Sandeep Kumar	M.B.B.S., M.S. (Ortho.), Senior Resident, Department Of Orthopaedics, Narayan Medical College And Hospital, Jamuhar, Rohtas, Bihar.
Dr. Kumari Rashmi*	M.B.B.S., M.D. (Patho.), Tutor, Department Of Pathology, Jawaharlal Nehru Medical College, Bhagalpur, Bihar. *Corresponding Author
Dr. Kumar Anshuman	Professor, Department Of Orthopaedics, Narayan Medical College And Hospital, Jamuhar, Rohtas, Bihar.
Dr. Debarshi Jana	Young Scientist (DST), Institute Of Post-Graduate Medical Education And Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India.

ABSTRACT

Anterior cruciate ligament injury is one of the most common injuries around knee and poses quite a lot management controversies. Anterior cruciate ligament has a pivot role in function and stability of the knee joint along with all other ligaments, being a prime stabilizer preventing the anterior translation of tibia over femur. Acute anterior cruciate ligament injury causes recurrent episodes of instability, pain and decreased motion. Anterior cruciate ligament reconstruction allows return to pre injury levels even in athletes, delays development of early osteoarthritis and reestablish the stability of the joint. Graft fixation during ACL reconstruction can be achieved with use of either metal screws or bio absorbable screws. Bio absorbable screws usage provide better visibility in postoperative MRI and also avoid removal at later stage. However there are controversies regarding the ideal graft, ideal fixation device, ideal time and technique of reconstruction. In this study we have compared the metal and bio absorbable screw for fixing the graft and reported the result.

KEYWORDS

Anterior Cruciate Ligament Reconstruction, Bio Absorbable Interference Screw, Titanium Interference Screw

INTRODUCTION

The high number of ACL injuries is a growing problem with serious consequences for the patient and society. An acute ACL injury is seldom isolated (15%), and is usually associated with concomitant injuries to the menisci (60%), cartilage (20%) and collateral ligaments.

Though Arthroscopic intra articular reconstruction has become gold standard in ACL reconstruction in this century there is still debate regarding the choice of graft, fixation methods, single or double bundle and trans portal or trans tibial technique.

More recent studies have proved quadrupled Hamstring is superior in strength but time for healing, probable loss of strength during healing and minimal hamstring weakness post operatively are considerations.

Recent studies have proved endobutton and bone mulch screw have a very high yield load than any other fixation device in view of soft tissue graft fixation. The trans portal technique has been widely used nowadays but the trans tibial technique is easily reproducible and gives comparable functional outcomes though tunnel placement is not more accurate in trans tibial technique.

Bio absorbable implants in orthopaedic surgery was introduced by Rokkanan *et al* and bostman *et al* for use in surgery of the ankle. In 1987, Kurosaka introduced the current concepts of interference screw fixation.

AIM

To do comparative analysis of the functional outcome of Arthroscopic Anterior Cruciate Ligament Reconstruction using quadrupled hamstring graft with endobutton as femoral fixation device and bio absorbable interference screw against titanium interference screw as tibial fixation devices respectively.

MATERIALS AND METHODS

In this study 60 patients treated with arthroscopic anterior cruciate ligament reconstruction with quadrupled hamstring graft with endobutton as the femoral fixation device and titanium interference screw (no=30) and bio absorbable interference screw (no=30) as tibial fixation device respectively between May 2019 and November 2020 at Department of Orthopaedics, Narayan Medical College and Hospital, Jamuhar, Rohtas, Bihar.

Inclusion Criteria

- Patients with closed growth plate

- Primary ACL surgery
- No evidence of multiple ligament injury
- No previous knee surgeries
- No ligamentous injury to contralateral knee

Exclusion Criteria

- Additional ligamentous laxity in affected knee
- Previous ACL surgery of either knee
- Chronic muscle disorders
- Any co-existing local conditions in the form of
 - o Active articular infection
 - o Inflammatory joint disease
- Metabolic bone disease
- Neoplastic disease

Evaluation

All the patients are subjected for post-operative antero posterior and lateral radiographs to determine the tunnel placement and position of endobutton in femur and interference screw in the tibia. Patients are followed at 4 weeks, 8 weeks, 3months, 6 months and once in 6 months thereafter.

Statistical Analysis

Data reported as mean and significant difference between the two groups was studied using Yate's corrected Chi-Square test.

OBSERVATION AND RESULTS

60 Cases of arthroscopy assisted Anterior cruciate ligament reconstruction with quadrupled hamstring tendon graft using endobutton as the femoral fixation device and titanium interference screw (no=30) and bio absorbable interference screw (no=30) as tibial fixation device respectively was followed for 6 months to 1.5 years. The mean follow up was 10.5 months

Age Distribution

Minimum age was 20 years and maximum age was 55 with a mean age of 31.6 (Table 1).

Table 1 : Age Distribution

Age in years	No. of Patients	Percentage
15-20	3	5
21-25	16	26.66
26-30	14	23.33
31-35	11	18.33

36-40	6	10
41-45	7	11.66
46-50	2	3.33
51-55	1	1.66
Total	60	100

Sex Distribution

In this study, 51 patients were males and 9 patients were females (Table 2)

Table 2 : Sex Distribution

Sex	No. of Patients	Percentage
Male	51	85
Female	9	15
Total	60	100

Side Involved

In this study, 40 patients had injury in the right knee and 20 patients had injury in the left knee (Table 3)

Table 3 : Side Involved

Side Involved	No. of Patients	Percentage
Right	40	66.66
Left	20	33.33
Total	60	100

Table 4 : Mode of Injury

Mode of Injury	No. of Patients	Percentage
Sports	10	16.6
Fall	22	36.66
RTA	28	46.66
Total	60	100

Table 5 : Duration of Injury

Duration of Injury	No. of Patients	Percentage
<6 weeks	12	20
1-3 months	10	17
3-6 months	14	23
6-12 months	14	23
>12 months	10	17
Total	60	100

Table 6 : Associated Injury

Associated Injury	No. of Patients	Percentage
Medial meniscus tear	13	22
Lateral meniscus tear	3	5
Both	2	3
Nil	42	70
Total	60	100

Observation

- Greater number of our patients was seen in the younger age group of 20-40 years.
- Male preponderance was noticed in our study
- Right side was involved more commonly than left side
- Road traffic accident was the most common cause accounting for ACL injury.
- Medial meniscus injury was involved more than the lateral meniscus.
- Most of the patients returned to their pre-functional level at 4 months.

Scoring Analysis

60 patients of arthroscopic ACL reconstruction with quadrupled hamstring graft was followed for a minimum period of 6 months and maximum period of 1.5 years. All patients are evaluated with Lysholm and Gillquist scoring at the end of 6 months. The maximum score achieved was 100 and minimum score was 56. The scores were graded as :

Outcome	Points
Good	84-100
Fair	65-84
Poor	<65

Two patients in titanium interference group and one patient in bio absorbable interference screw group lost to follow up.

Outcome	Titanium screw group, No. of patients (28)	Percentage	Bio-abs screw group, No. of patients (29)	Percentage
Good	23	82.14	24	82.75
Fair	3	10.71	3	10.34
Poor	2	7.14	2	6.89

By Yates corrected Chi-Square Test,
 $\chi^2 = 0.06$ $P = 0.97$

The clinical outcome was nearly equal in both the groups.

DISCUSSION

Our study is to evaluate the functional outcome of arthroscopic single bundle ACL reconstruction with quadrupled Hamstring graft with

transtibial and transportal techniques using endobutton as femoral fixation device and titanium interference screw in 30 patients and bio absorbable interference screw in 30 patients as tibial fixation device.

In our study fall and road traffic accidents predominated as the cause of injury accounting for 37% and 47% respectively. Sports injuries accounted for only 10% in contrary to all international studies. D W Lewis reported 58% meniscal injury associated ACL tear at presentation. Medial meniscus was involved more than the lateral meniscus in his study and he also proposed meniscal repair or resection did not alter the outcome and chondral lesions are a better predictor of functional outcome. Stephen Lyman reported more than 50 % meniscal procedures with ACL reconstructions in 2009. In our study 30% of patients had meniscal injury at presentation and medial meniscus injury predominated lateral meniscus injury like other studies. None of our patients had significant chondral damage at diagnostic arthroscopy.

The fixation of the graft has been proved to be the site of failure rather than the graft itself irrespective of the type of graft especially in the early rehabilitation phase when the graft integration has not taken place and the fixation is of little significance after 8 to 12 weeks when graft has integrated with the bone as proposed by Dawn T Gulick.

In our study we used transtibial or transportal single bundle reconstruction with quadrupled hamstring graft placing the femoral tunnel between 10 30 and 11'o clock position in the right knee and between 1'o clock and 1 30 position in the left knee. John Paul proposed that placing graft at 10 30 position and 1 30 position in single bundle reconstruction reconstructs portions of anteromedial and posterolateral bundles. Masayoshi Yagi showed that anatomic reconstruction allowed better rotatory stability than nonanatomic placements of graft. Asheesh Bedi showed that trans portal placement of tunnel achieved more lateral placement than the trans tibial drilling and trans tibial approach to achieve lateral tunnel placements resulted in over reaming of tibia. Though double bundle reconstructions have gained attraction and studies have shown double bundle reconstruction to be superior in providing stability in high demand patients. Adachi, Ochi and Uchio showed no significant advantage of double bundle reconstruction than anatomic single bundle reconstruction in factors of stability and proprioception in general population.

The metallic screws distort the knee MRI wherein bio absorbable screw avoids impairment of imaging. Apart from this metallic screws have to be removed during surgical revision wherein bio absorbable screws would have been degraded. The major disadvantages are screw breakage at the time of insertion and postoperative inflammatory reaction causing synovitis. We did not come across such problems in our study.

In our study, functional outcome evaluated by Lysholm and Gillquist scoring was nearly equal in both titanium interference screw study group and bio absorbable interference screw study group and it is statistically insignificant with P value of 0.97. Our study shows that there is no significant difference in the outcomes associated with the use of titanium and bio absorbable interference screws used for anterior cruciate ligament reconstruction.

Since our study was a short term follow up we could not comment about the arthritic changes post operatively. Fox *et al'* reported 3 to 17% incidence of anterior knee pain, compared to 13% in our study, Apostolopoulos reported 10% of anterior knee pain. Kurt Spindler stated regular exercise can lead to increased outcomes in 2005. Our patients are put on home based physiotherapy programme insisting on knee flexion and quadriceps strengthening and mean flexion achieved was 135 degree. J A Grant concluded that home based physiotherapy is cost effective and not significantly inferior to supervised programmes. As overall conclusion several factors influence the functional outcome in arthroscopic ACL reconstruction. Factors like graft choice, graft fixation, tunnel placement and graft tensioning play a vital role in altering the final outcomes.

CONCLUSION

The results of our study were comparable with already published reports of comparative study done using bio absorbable versus metal interference screws. Our study shows that there is no difference in functional outcome whether bio absorbable or titanium interference screw was used.

The success of ACL reconstruction depend on the correct technique used for the surgery, precise placement of graft and rehabilitation methods than on type of graft fixation device used, neither titanium nor bio absorbable screws.

The blunt metal or titanium screw has been the de facto standard in graft fixation. Since the alternate bio absorbable screw overcomes some of the potential drawbacks, it should become the de facto standard in the future.

REFERENCES

1. A Cadaveric Study Relating Transtibial Lateralized Femoral Tunnel Position to the Anteromedial and Posterolateral Bundle Femoral Origins of the Anterior Cruciate Ligament Biomechanical Analysis of an Anatomic Anterior Cruciate Ligament Reconstruction Masayoshi Yagi, Eric K. Wong, Akihiro Kanamori, Richard E. Debski, Freddie H. Fu and Savio L-Y. Woo Am J Sports Med 2002; 30:660.
2. Adachi N, Ochi M, Uchio Y, Iwasa J, Kuriwaka MY. Ito Reconstruction of the anterior cruciate ligament SINGLE-Versus Double-Bundle Multi-stranded hamstring Tendons. J Bone Joint Surg [Br]. 2004; 86:515-20. Received, 2003.
3. Alwynsmit. Anatomy, biomechanics, diagnosis and treatment clinical ortho - page 172 - 180. Alexander A sapoga anatomical and biomechanical consideration. JBJS. 1983, 1990, 250.
4. Asheesh Bedi MD, Volker Musahl MD. Volker Steuber MD. Transtibial Versus Anteromedial Portal Reaming in Anterior Cruciate Ligament Reconstruction: An Anatomic and Biomechanical Evaluation of Surgical Technique Arthroscopy: The Journal of Arthroscopic and Related Surgery. 2011; 27(3):380-390.
5. Bonnet A. Traite Des Maladies Articulaires -2nd edition: Baillire, Paris.pp 1853; 354-357.
6. Chae-Gwan Kong, MD1, Yong In, MD2, Geon-Hyeong Kim, MD1 Cross Pins versus Endobutton Femoral Fixation in Hamstring Anterior Cruciate Ligament Reconstruction: Minimum 4-Year Follow-Up Knee Surg Relat Res. 2012; 24(1):34-39.
7. David Greenberg MD, Michael Robertson MD, Santaram Vallurupalli MD, Richard A, White MD, William C. Allen, MD Allograft Compared with Autograft Infection Rates in Primary Anterior Cruciate Ligament Reconstruction. J Bone Joint Surg Am. 2010; 92:2402-8.
8. Dawn T. Gulick Anterior Cruciate Ligament Reconstruction: Clinical Outcomes of Patella Tendon and Hamstring Tendon Grafts Journal of Sports Science and Medicine. 2002; 1:63-71.
9. Fu FH, Bennett CH, Ma CB. Current trends in anterior cruciate ligament reconstruction: Operative procedures and clinical correlation. Am J of Sports Med. 2000; 28:124-130.
10. Galen C. On the usefulness of the parts of the body. Ithaca Cortnet Univ, 100.
11. Howe, Johnson, Kaplan. ACL reconstruction using quadriceps patellar tendon graft. Part I. Long term follow up. Am J of Sports Med 1991; 19:447-57.
12. Kurosaka M, Yoshiya S, Andrich JT. Biomechanical comparison of different surgical techniques of graft fixation in anterior cruciate ligament reconstruction Am. J Sports Med. 1987; 15:225-229.
13. Petteri Kousa MD, Teppo L. N. Ja'rvinen,†§ MD, PhD, Mika Vihavainen,† Pekka Kannus,† MD, PhD, and Markku Ja'rvinen,† MD, PhD The Fixation Strength of Six Hamstring Tendon Graft Fixation Devices in Anterior Cruciate Ligament Reconstruction The American Journal Of Sports Medicine. American Orthopaedic Society for Sports Medicine, 2003.
14. Reconstruction Using Biodegradable Interference Fit Fixation A Prospective Matched-Group Analysis Am J Sports Med 2005; 33:1327.
15. Robert G. Marx, M.D., M.Sc., F.R.C.S.C., and Christopher R. Spock, B.A. Complications Following Hamstring Anterior Cruciate Ligament Reconstruction With Femoral Cross-Pin Fixation Arthroscopy: The Journal of Arthroscopic and Related Surgery. 2005; 21(6):762.e1-762.e3.
16. Satku K, Kumar VP, Ngoi SS. ACL injuries. To counsel or to oper-ate? J Bone Joint Surg Br 1986; 68:458-6.
17. Segond PF. Recherches cliniques et experimentales sur les epandements sanguins du genou par entorse. Prog med 1879; 16: 297-421.
18. Steiner ME, Hecker AT, Brown CH Jr, Hayes WC. Anterior cruciate ligament graft fixation: comparison of hamstring and patellar tendon grafts. Am j sports Med 1994; 22:240-6.
19. Thomas D Rosenberg Extensor Mechanism after Patellar Tendon Harvest in Acl Reconstruction Ajsm. 1992; 20(5).