CLINICOPATHOLOGICAL CORRELATION OF OVARIAN TUMORS IN DARBHANGA MEDICAL COLLEGE AND HOSPITAL, LAHERIASARAI, BIHAR

Pathology

Dr. Megha Jha	M.B.B.S., M.D. (Pathology), Tutor, Department of Pathology, Darbhanga Medical College, Laheriasarai, Bihar.
Dr. Hari Shankar Mishra*	M.B.B.S., M.D. (Pathology), Associate Professor, Department of Pathology, Darbhanga Medical College, Laheriasarai, Bihar. *Corresponding Author
Dr. [Prof.] Ajit Kumar Chaudhary	M.B.B.S., M.D. (Pathology), Professor and Head of Department, Department of Pathology, Darbhanga Medical College, Laheriasarai, Bihar.
Dr. Debarshi Jana	Young Scientist (DST) Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India

ABSTRACT

Introduction and Objective: Ovary is one of the complex organs in terms of its embryology, histology, steroidogenesis and tumor incidence. The present study highlights the correlation between the clinical and pathological features of neoplastic lesions of ovary in order to arrive at a better understanding of the disease process. The objective of study is to assess the overall incidence of various histological types of ovarian neoplasm, the pattern of occurrence of ovarian tumors in relation to age, parity and modes of presentations and to study the histopathological findings of various ovarian tumors. **Materials and Methods:** Biopsies received at the Department of Pathology, Darbhanga Medical College and Hospital, Laheriasarai, Bihar were subjected for histopathological study. Specimens were then fixed in 10% formalin for 24-48 hour, processed and embedded in paraffin blocks. Sections of 4-6 microns thickness were taken and stained with routine haematoxylin and eosin and studied. IHC was done whenever necessary. **Result:** Out of 92 cases of ovarian tumors 68 cases were benign, 3 were borderline and 21 were malignant. Serous cystadenoma formed the majority (38%) among benign lesions and among malignant lesions serous cystadenocarcinoma (5.4%) were common. Majority were surface epithelial tumors and constituted 64.13% (59 cases) with germ cell tumors 26% (24 cases) and sex cord stromal tumors 8.6% (8 cases). **Conclusion:** The correct histopathological diagnosis of ovarian tumor is of prime importance in view of their behavioural predictability, clinical correlation and the proper management of patient.

KEYWORDS

Ovary, Tumor, Histopathology.

Introduction

An adnexal mass (mass of the ovary, fallopian tube, or surrounding connective tissues) is a common gynaecologic problem. In the United States, it is estimated that there is a 5 to 10 percent lifetime risk for women undergoing surgery for a suspected ovarian neoplasm. Ovarian cancer is the leading cancer in women (affecting about 1/70) and leading cause of death from gynaecological cancer. It is the 5th leading cause of cancer fatalities in women² after lung cancer, breast cancer, colon cancers and cervical cancer. Ovarian neoplasms remain asymptomatic until massive ovarian enlargement cause compression of pelvic structures, ascites, abdominal distension and distal metastasis. The ovary not only gives rise to a wide variety of malignancies but is also a favourite site for metastases from many other organs.

Ovarian neoplasms are notorious for wide range of histologic differentiation that can be found in different areas of the same neoplasm. The present study is undertaken to analyse the ovarian tumours and their characteristics with regard to age, parity and mode of presentation, further a detailed study of various histological types of ovarian tumours is also done.

Materials and Methods

92 cases of ovarian tumours received in the Department of Pathology, Darbhanga Medical College and Hospital, Laheriasarai, Bihar were studied. The study was undertaken during the period October 2019 to September 2020. The materials were received as surgically resected specimen from patients operated in the hospital. Clinical details like age, obstetric history, menstrual irregularities and other constitutional symptoms were collected in the proforma. CA 125 level was measured for 20 patients who were clinically and sonologically suspected to have ovarian tumour. On receiving the specimen, gross features such as size, shape, colour, external appearance, findings on cut section and contents were noted. Then the tumours were cut at various levels depending on the individual cases and they were allowed to fix in 10% formalin for 24-48 hours. After formalin fixation, multiple bits were taken from representative areas of tumours and the accompanying tissues. They were processed for histopathological examination and paraffin blocks were made. The blocks were cut at 3-5 microns thickness and stained with Haematoxylin and Eosin. Detailed

microscopic examination of the tumour was done to arrive at a histopathological diagnosis following the WHO classification of the ovarian tumours.

For a case of Granulosa cell tumour IHC with Inhibin was done. (Fig. 6) The data compiled was analysed for various parameters like age, parity, oral contraceptive use, clinical signs and symptoms, gross features of the tumours and the incidence of the different histological types.

Results

Ninety-two cases of ovarian tumours received in the Department of Pathology, DMCH, Laheriasarai, Bihar during the period of 1 year from October 2019 – September 2020 were studied. Out of 92 cases of ovarian tumours, 68 were benign, 3 were tumours of low malignant potential and 21 were malignant. (Table 1).

Table 1 : Distribution of ovarian tumours

Type of tumour	No. of cases	Percentage
Benign tumour	68	73.9%
Borderline tumour	3	3.2%
Malignant tumour	21	22%
Total	92	100%

The surface epithelial tumours were the commonest tumours accounting for 64.13%, germ cell tumours were 26% of cases and sex cord stromal tumours formed 8.6% and metastatic tumour 1% In the present study, the youngest patient was 14 months and the oldest was 75 years forming a range of 14 months to 75 years. Highest incidence of ovarian tumour was noted in the third decade i.e. 28 cases out of 92 cases accounting for 30.43%. Highest incidence of benign ovarian tumour was noted in third decade i.e. 25 cases out of 68 accounting for 36.76%. Highest incidence of malignant tumour was noted in the fourth decade i.e. 10 out of 21 cases accounting for 47.61% (Table 2).

Table 2 : Age group distribution of benign, borderline and malignant ovarian tumours

Age in yrs.	Benign	Borderline	Malignant	Total	Percentage

0-10	2	-	2	4	4.3%
11-20	2	-	-	2	2.1%
21-30	12	-	1	13	14.1%
31-40	25	-	3	28	30.4%
41-50	18	3	10	31	33.6%
51-60	4	-	3	7	7.6%
61-70	5	-	-	6	6.5%
71-80	-	-	2	2	2.17%
Total	68	3	21	92	100%

Presenting Complaints: The distribution of symptoms is varied. 35% of the patients complained of dull aching lower abdominal pain, 27% complained of abdominal mass and 6% of the patients gave history of menstrual disturbance like menorrhagia. History of postmenopausal bleeding was given by 4 patients with benign tumours. (Table 3).

Table 3: Symptoms of ovarian tumours

Clinical presentation	No. of cases Benign Tumour	No. of cases Borderline Tumour	No. of cases Malignant Tumour	Percentage
Pain abdomen	27	2	4	35.86%
Mass per abdomen	20	1	4	27.13%
Pain abdomen with mass	7	-	4	11.95%
Menstrual disturbance	4	-	2	6.52%
Urinary disturbance	2	-	4	6.52%
Constitutional symptoms	4	-	2	6.52%
White discharge per vagina	4	-	1	5.43%

Physical signs: 27% of the patients had a pelvic mass. 12% of patients complained of dull aching lower abdominal pain and tenderness. Ascites was seen in 2 patients with mucinous cystadenocarcinoma and 2 patient in serous cystadenocarcinoma.

Parity: 6 patients were not married and all were below twenty years of age. Among married, 81 were parous and remaining were nulliparous. (Table 4).

Table 4: Distribution of ovarian tumours in parous women

Types of tumours	Unmarried	Married	
		Nulliparous	Parous
Benign	4	5	59
LMP	0	0	3
Malignant	2	0	19
Total	6	5	81

Associated Conditions: Out of 92 cases of ovarian tumors, 22 were associated with appendicitis and 12 were associated with uterovaginal prolapse. (Table 5).

Table 5: Conditions associated with ovarian tumours

Sl. No.	Associated conditions	No. of cases
1	Utero Vaginal prolapse	12
2	Leiomyoma	5
5	Appendicitis	22
6	Calculus cholecystitis	1
7.	Pregnancy	1

Oral Contraceptive use: None of the above 92 patients gave history of OCP use.

Family History: This may not be significant as the patients were not aware of the cause of death of their relatives.

The tumours were classified according to the WHO histological classification of the ovarian tumours and the incidence of different histological types noted. (Table 6).

Table 6: Incidence of various histological types of the ovarian tumours

Types of Tumour	No. of cases	Percentage
I. Common Epithelial Tumours		
A. Serous tumours	59	64.13%
a) Benign	35	38%
b) Low Borderline malignancy	2	2.17%
c) Malignant	5	5.4%
B. Mucinous tumours		
a) Benign	7	7.6%
b) LBM	1	1%
c) Malignant	4	4.3%
C. Mixed epithelial tumours Benign		
Malignant	1	1%
D. Endometrioid carcinoma	4	4.3%
E. Transitional cell carcinoma	-	
F. Undifferentiated carcinoma		
II. Sexcord stromal tumours	08	8.6%
A. Granulosa cell tumour	3	2.17%
B. Fibroma/thecoma	5	5.4%
III. Germ cell tumours	24	26%
A. Dysgerminoma	1	1%
B. Endodermal Sinus tumour	1	1%
C. Embryonal carcinoma	-	
D. Teratoma, mature cystic	21	22.8%
E. Immature teratoma	1	1%
IV. Metastatic tumours		
Krukenberg tumour	1	1%

Fig. 1: Hair shaft and greasy material in mature cystic teratoma

Fig. 2: Benign cystic teratoma- 10x showing keratin, sebaceous glands and cartilage

Fig. 3: Papillary excrescences in the cystic cavity

Fig. 4: Papillary serous cystadenocarcinoma

Fig. 5: Granulosa cell tumor – 10x shows cuboidal to polygonal cells having coffee bean nuclei with folds

Fig. 6: Inhibin positivity

CHARACTERISTICS OF THE TUMOURS:

Site of Involvement: Left sided tumours of ovary (51.08%) were more common than right sided tumours (43.47%). 5 cases were bilateral out of which 2 were borderline and 3 malignant.

Size of the Tumours: The largest tumour (mucinous cystadenoma) in the present study measure 34x26x28cm in size and weighed 11 kg while the smallest tumour (serous cystadenoma) was 2x2x1cm in size and weighed 50gms. (Fig. 3).

CA 125 level was measured for 20 patients who were clinically and ultrasonographical suspected to have ovarian tumours. Of that, only 12

patients had elevated levels ($>35\text{U/ml}$). Also in that group of 12, it was noted that CA125 level was above 35 u/ml but not beyond 100 u/ml in benign tumours. The mucinous carcinomas showed moderately elevated levels of 105 u/ml and 360u/ml . relatively much elevated levels were noted in serous carcinoma and malignant mixed mulleriantumour, 678 u/ml and 575 u/ml respectively.

Discussion

Ovarian/tubal neoplasms may arise from stem cells, which typically give rise to the surface epithelium, fallopian tube epithelium, germ cells, or sex cord-stromal cells. The incidence, clinical appearance and the behaviour of the different types of ovarian tumours is extremely variable. Through physical examination, imaging studies like pelvic ultrasound and laboratory studies like serum biomarkers and immunological tests have been reported to be of some help in predicting the nature of the pathology. It is generally impossible to diagnose the nature of the ovarian tumours preoperatively. Surgical evaluation allows a definitive histologic diagnosis. Microscopic examination and diagnosis is essential for further management of the tumour.

Out of 92 ovarian tumours in the present study 73.9 % were benign tumour, 3.2% were borderline tumours and 22% were malignant tumours.

Nature of the Tumors: Among the different histopathological patterns the surface epithelial tumours formed the largest group of tumour (59 cases, 64.13%) followed by the germ cell tumour (24 cases, 26%), sex cord stromal tumours of (8 cases, 8.6 %) and metastatic tumours (1%).

Nalini et al in 2007 and Mondal et al ⁵ in 2011 observed that the epithelial tumours were the most frequent tumour followed by germ cell tumours and sex cord tumours.

The commonest epithelial tumours were serous cystadenoma (35 cases) and the commonest germ cell tumour was benign cystic teratoma (21 cases) in the present series. Similar observations were made by Mondal SK et al in 2011 and Nalini et al in 2007.

Among the benign lesions, serous cystadenoma was the commonest (35 cases, 38%) (Fig. 3) followed by mature cystic teratoma (21 cases, 22.8%) (Fig. 1, Fig. 2) in the present study. This was similar to the observations of Di Bonito et al, Nalini et al and Mondak SK et al. But Ahmed et al, in his paper, stated that mature cystic teratoma (35.17%) was the commonest benign tumours followed by surface epithelial tumours. The commonest malignant tumours in the present study is serous cystadenocarcinoma (5.4%) (Fig. 4) and the next commonest being Mucinous cystadenocarcinoma and endometrioid carcinoma each constituting 4.3%, this is followed by Granulosa cell tumour 3.2 % (Fig. 5) and Metastatic tumour 1%. In a study from eastern India, the same was found to be 5% of all malignant tumors.

Laterality: 5 cases (5.43 %) of bilateral ovarian tumour were seen in the present study, of which 3 cases were malignant serous cystadenocarcinoma. similar findings have been reported by Couta F et al whereas Ramachandran G et al, Gupta SC et al and Kapas MM et al reported more number of bilateral tumours compared to the present study.

Bilaterality in malignancies implies spread to the opposite ovary as part of extension throughout the pelvis and abdomen as seen in advanced cases.

Site of Involvement: In the present study, there were 51.08% of left sided tumours and 43.47% of right sided tumours, the remaining 5.4% were bilateral.

Ramachandran G at el found 46.04 % of all ovarian tumours on the right side and 38.5 % on the left side. The rest were bilateral.

Age: The youngest age is a 14 months old child with mature cystic teratoma. Similarly Pilli et al reported the youngest patient of 8 months. In the present study majority of benign tumours occurred in the 31–40 years of age group and the carcinomas were more common at 41–50 years of age group. Surface epithelial tumours varied widely involving all age groups. The present findings concurred with those of

Ashley DJB(1990) and Herbst A (1994). Similar observations were also made by Ramachandran G et al and Mondal SK et al in 2011.

Parity: In 2006 Gunnar et al did a prospective study onreproductive factors and risk of ovarian cancer in 6565 females in Norway and found that highest risk of ovarian tumours was observed among nulliparous women. The risk decreased significantly with increasing parity.

In 2006 Marine et al did a study on the incidence of ovarian cancer on 87,929 grand multiparous women and concluded that the risk of ovarian cancer was low in all grand multiparous women, no matter how many children and at which age they delivered or contracted cancer.

Out of 92 cases in the present study, 81 were multiparous, 11 were nulliparous of whom 6 were unmarried

Oral Contraceptive Pills Use: In a review of epidemiological study of ovarian carcinoma, Herbst al stated that nulliparous and infertile women particularly those who had used fertility drugs, had an increased risk of ovarian carcinoma. Surprisingly no patients gave the history of any oral contraceptive use in the present study. This might be because of the fact that majority of the patients had come from a rural background.

Family History: Very few patients in present study gave family history of ovarian and breast cancer majority of the patients were ignorant of the exact cause of the death of their parents, siblings and other relatives.

However, Gershenson DM stated that familial ovarian cancer probably accounts for no more than 3% of all epithelial ovarian cancers, with the sporadic form comprising the vast majority of cases. According to Anderson MC, 5% of women who get ovarian cancer have no family history of the disease.

Symptoms: The commonest clinical features in the present study were pain in the lower abdomen (35.86%) followed by mass per abdomen (27.17%).

Present study concorded with Pilli et al¹³ where abdominal pain was the commonest symptoms. But cases presenting as mass per abdomen were less in the present study when compared to other studies.

Size of the Tumors: The largest tumours encountered in the present study was mucinous cystadenoma measuring $35\times 28\times 26\text{cms}$ in size. Similar observations were made in Tyagi et al and Gupta et al who reported a mucinous cystadenoma with maximum diameter of 44.5cms. Majority of the cases were uni/multilocular with a cystic appearance. The tumours with mixed solid and cystic areas and completely solid tumours were mostly malignancies. Similar observations were made by Gupta SC et al¹¹ and Maheshwari et al.

Serum CA 125 Level: Serum CA 125 is the most commonly used laboratory test for the evaluation of adnexal masses for Epithelial Ovarian Carcinomas. CA 125 is a high molecular weight glycoprotein which is widely used prospectively to evaluate the therapeutic efficacy and to monitor the disease status among ovarian cancer patients. The sensitivity of serum CA125 level for stage I disease is lower (50%) than for stage II (90%) or higher disease. The prevalence of an elevated serum CA125 is highest in women with serous histology and lower in those with mucinous tumours, many of which are associated with normal level of CA125. In premenopausal women, elevated serum CA 125 levels are nonspecific and may be elevated due to non-gynaecological reasons and benign abdominal disorders like pregnancy, PID, endometriosis, uterine fibroid, spontaneous abortions etc. In contrast, an elevation of serum CA 125 level in postmenopausal women indicates the necessity for prompt surgical exploration as it is most likely associated with ovarian carcinoma.

References

1. Ahmed Z, Kalyani N, Hassan SH, Muzaffar S, Gilli MS Histological pattern of ovarian neoplasm. J Pakistan medical association 2000;50:46-49.
2. American College of Obstetricians and Gynecologists committee on Gynecologic practice. Committee opinion no.477: the role of obstetrician-gynecologist in the early detection of epithelial ovarian cancer. Obstetgynecol 2011;117:742.
3. Anderson MC: Female Genital tract, systemic pathology, vol 6, symmers W, st.c,3 Edition 1991, Churchill, livingstone.
4. Basu P, DeP, Mandal S, Ray K, Biswan J, study of patterns of Ovarian cancer patients in a

- specialized Ca institute n Kolkata, Eastern India Use the "Insert Citation" button to add citations to this document. Indian J Cancer 2009;46:28-23.
5. Batcharya MM, Shinde SD, Purrandare VN. A clinicopathological analysis of 270 ovarian Tumors J. Postgradmed, 1980;26:103-7.
 6. Carlson KJ, SkatesSJ, Singer DE. Screening for Ovarian cancer. Ann Intern med 1994;121:124.
 7. Couto F, Naolkami N.S, Jose M: Ovarian tumours in Goa, A clinicopathologic study J obstGynec India 43:408-412,1993.
 8. Di Bonito L, Patriarca S, Delendi M, Alberico S, Ovariantumours: anatamohistopathological contribution to their interpretation, Eur J Gynaecol Oncol.1988;9(4):324-30.
 9. Gershenson DM: Epithelial ovarian cancer, Textbook of gynecology, cope land LJ,1993, WB Sauder Co.
 10. Gunnar Kvale,IvarHeach, SteinarNilssen.Reproductive factors and risk of ovarian cancer. International Journal of cancer 2006;42(2):246-251.
 11. Gupta SC Singh PA, Mchrotra T.N. A Clinicopathological study of ovarian tumours India J Path Micro 29,354-362,1986.
 12. Herbst L:The epidemiology of ovarian carcinoma and the current status of tumour markers to detect disease Amjobst and gynec 170:1099-1107,1994.
 13. Jemal A, Siegal R, Xu J, Ward E. Cancer statistics, 2010. CA Cancer J Clin 2010; 60:277.
 14. Kapas MM and RalMC: Varities of ovarian neoplasm. Jobst and Gynec India 32,810-815,1987.
 15. Maheswari V, Tyagi S.P, Saxena et al: surface epithelial tumours of the ovary. Ind J Patho Microbiology 37:75-85,1994.
 16. Mondal SK et al Histological pattern, Bilaterality and Clinical evaluation of 957 cases of ovarian tumours-A 10 year study in a tertiary care hospital of Eastern India, Journal of cancer Research and Therapies 2011
 17. Nalinigupta, DiptiBisht, Anil Kumar Agarwal, Veena K Sharma, Retrospective and prospective study of ovarian tumours and tumour like lesions Indian J patholMicrobiol 2007;50(3):525-527.
 18. National Institutes of Health Consensus Development Conference Statement. Ovarian cancer: screening, treatment, and follow-up. GynecolOncol 1994;55:S4.
 19. Pilli GS, Suneetha KP, Phaded AV, Yenni VV. Ovarian tumours: a study of 282 cases J Indian Medical association 2002;423-4.
 20. Ramachandran G, Harilal KR, Chinnamma KK and ThangaVelu H: Ovarian neoplasm-a study of 903 cases, JObst and Gynec India:309-315,1972.
 21. Tyagi SP, Maheswari V, Tyagi N, Saxena K, Sharme R and Hamced F: Solid tumours of the ovary, JIMA;91:227-230,1993.