CLINICAL PROFILE OF ISOLATED SYSTOLIC HYPERTENSION AND IT'S CARDIOVASCULAR AND RENAL COMPLICATIONS AT AIIMS, PATNA, BIHAR

Medicine

Dr. Rajeev Ranjan Kumar

M.B.B.S., M.D. (Med.), Senior Resident, Department Of Medicine, AIIMS, Patna, Bihar.

Dr. Shiv Kumar*

M.B.B.S., M.D. (Med.), Senior Resident, Department Of Medicine, AIIMS, Patna, Bihar.
*Corresponding Author

Dr. Ravikirti

Additional Professor, Department Of Medicine, AIIMS, Patna, Bihar.

Dr. Debarshi Jana

Young Scientist (DST), Institute Of Post-Graduate Medical Education And Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India.

ABSTRACT

Hypertension, its risk factors and complications is very poorly understood therefore it goes undiagnosed and untreated for a long time. The present study was undertaken to study the clinical profile of isolated systolic hypertension (systolic > 140 and diastolic ≤ 90 mmHg) in elderly (above the age of 60 years) and to find out any other associated risk factors, any end organ complications, specially cardiac through electrocardiography and echocardiography and on kidney through creatinine clearance. The present cross-sectional study was undertaken on patients attending the outdoor department and indoor department of medicine. The study group comprised of 75 elderly patients who had isolated systolic hypertension (ISH). Using Chi square test correlation of age with stage of BP, association of risk factors with ISH and Stage III BP and ECG changes with LVH calculated. It concluded that ISH associated with risk factors has definite effect mainly on cardiac in terms of LV hypertrophy and on kidney in terms of reduced creatinine clearance and is an even better predictor of mortality and morbidity than is diastolic blood pressure.

KEYWORDS

Clinical Profile, Isolated Systolic Hypertension, Cardiovascular, Renal Complication.

INTRODUCTION

In India awareness of hypertension, its risk factors and complications is very poor. Hence, hypertension goes undiagnosed and untreated for a long time. Indeed the JNC- 7 and JNC- 8 report states that in persons older than 60 years, systolic blood pressure is a much more important cardiovascular disease risk factor than diastolic blood pressure. Hence the present study was undertaken to study the clinical profile of isolated systolic hypertension (systolic > 140 and diastolic ≤ 90 mmHg) in elderly (above the age of 60 years), to find out any other associated risk factors, any end organ complications, specially on cardiac through electrocardiography and echocardiography and on kidney through creatinine clearance

AIM AND OBJECTIVES

The main aim of the study is to evaluate the clinical profile of isolated systolic hypertension (SBP > 140 mmHg and DBP ≤ 90 mmHg) in elderly patients (60 Yrs and above).

To detect left ventricular hypertrophy in such patients by electrocardiography and echocardiography and to study the effect of isolated systolic hypertension on kidney.

MATERIALS AND METHODS

The present cross-sectional study was undertaken on patients attending the outpatient department as well as indoor-patients of All India Institute of Medical Sciences (AIIMS), Patna, Bihar during the period from October 2019 to September 2020. The study group comprised of 75 elderly patients (Male=49, Female=26) of the age group of 60 to 89 years. All of these patients had isolated systolic hypertension i.e. (SBP > 140 mmHg and DBP ≤ 90 mmHg), detected first time. Patients below 60 years of age, patients with previous history of hypertension or hypertensive's on medication, patients having chronic kidney disease due to etiology other than hypertension like diabetes, glomerulonephritis, patients having secondary hypertension, patients on any herbal medication and HIV were excluded from the study.

Detailed evaluation of medical status of these patients were done. The evaluation comprised of detailed clinical history, thorough physical examination with relevant investigations. Blood pressure recording was done as per JNC VII recommendation. Standard mercury sphygmomanometer properly calibrated and validated instrument was used. An appropriate sized cuff (Cuff bladder encircling at least 80% of the arm) used. Patients were seated quietly for at least 5 minutes in a chair, with feet on the floor and arm supported at heart level. Standing

blood pressure was recorded. At least two measurements on each of 2 or more visits and mean of the these readings were taken. The first appearance of the sound (phase I, Korotkoff) is the SBP and the disappearance of the sound (phase V) is the DBP. Blood pressure recordings were documented and staging of ISH was done. General physical examination and systemic examination was done. Height and weight measured for each patient. Body surface area was calculated using the DuBois formula: $BSA (m^2) = 0.007184 \times [\text{weight (kgs)}]^{0.425} \times [\text{Height (cm)}]^{0.725}$. Waist and hip ratio calculated. Detailed fundoscopic examination done and hypertensive retinopathy was graded to assess target organ damage. Fundoscopic findings were confirmed by an ophthalmologist, who was blinded to clinical and echocardiographic findings. Routine investigations Hb%, TC, DC, urine – albumin, sugar, microscopy done. Hb% less than 12 gm were excluded from study. Blood tests, Fasting BS, PPBS, B.urea, S.creatinine and lipid profile was done. Creatinine above 1.4 mg% was excluded FROM study. ECG was recorded at 25 mm/sec speed and 10 mm/mv standardization.

The following were analyzed for LVH

1. Sokolow-Lyon criteria (SLC): SV1+RV5 or V6 (whichever is greater) ≥ 35 mm
2. Modified Romhilt-Estes point score system.

These criteria were selected since they are most commonly used in diagnosis of LVH.

Echocardiography: 2-D guided M-mode echocardiography was done by an experienced echocardiographer on a Hewlett Packard Color Doppler machine using 3.5 MHz phased array sector transducer. The echocardiographer was blinded to ECG results. M-mode echocardiographic measurements were obtained using parasternal long axis view, at the mid cavity level as defined by the tip of papillary muscle. The measurements of Left Ventricular Internal Dimension (LVIDd), Inter-Ventricular Septal Thickness (IVSTd) and Posterior Wall Thickness (LVPwd) were made at the end. Diastole is defined by the peak of R-wave on a simultaneously recorded ECG.

Measurements were done using the ASE convention (Standard convention). In ASE convention, the thickness of the endocardial surfaces is excluded from the measurement of LVIDd and included in the measurement of interventricular septum and posterior wall thickness. The mean measurement of three consecutive beats was taken. ASE cube LV mass was calculated by the formula.
 $ASE \text{ Cube LV mass} = 1.04 (LVIDd + IVSTd + LVPWD)^3 - (LVIDd)^3$

Since this method overestimates the LV mass by a mean of 25%, correction of the overestimation is done by using the equation:
 $LV\ mass = 0.80\ (ASE-cube\ LV\ mass) + 0.6\ gms.$ The LV mass was indexed (LVMI) to body surface area.
 $LVMI = LV\ mass/BSA$ and expressed as gms/m^2 .

All the values were expressed as mean $\pm$ SD. Sex specific partition values were defined based on the normal values suggested for Indian population. Echocardiographically determined LV mass was taken as the gold standard and each electrocardiographic criteria were correlated with the echo LV mass.

The ejection fraction is a global index of left ventricular fibre shortening and is generally considered as one of the most meaningful measures of LV pump function. Angiographic studies have shown that the normal ejection fraction averages 72%-78% and ranges from 56% to 78%. Ejection fraction, which is the ratio of the stroke volume to the end diastolic volume noted for each patient $EF = EDV-ESV \times 100\ EDV$ Regional wall motion abnormality, if present was noted.

Normal range

LVIDd = 3.6 to 5.4cms

LVPWd = 0.6 to 1.1cms

IVSTd = 0.6 to 1.1cms

LVDs = 2.4 to 4.2 cms

LV dilatation: Cavity volume $\geq 90\ ml/m^2$

Patients having cavity volume $> 90\ ml/m^2$ were noted.

Statistical Analysis

Chi square and Fisher exact test have used to find the significance of proportion of risk factors between increased LVMI and Normal LVMI.

RESULTS

- 53.33% of the patients are in the age group of 66-75 years.
- Females are more in the age group of < 70 years than males with $p=0.063$.
- Correlation of age with stage of blood pressure

Table – 1

Age in years	Stage I (140-159)	Stage II (160-179)	Stage III (>180)
≤ 60	1	3.70	-
61-65	12	44.44	1
66-70	11	40.74	8
71-75	1	3.70	13
76-80	1	3.70	5
> 80	1	3.70	-
Total	27	36.00	27
Mean $\pm$ SD	66.67 $\pm$ 5.56	72.56 $\pm$ 4.01	± 6.72

57.3% of patients are asymptomatic, 42.7% are symptomatic. Symptomatic patients are more in the stage III BP group with $p<0.001$. Symptoms include breathlessness, swelling of lower limbs, dizziness, headache, blurred vision, palpitations, and motor weakness.

Table - 2 Associated risk factors with ISH

Associated Risk factors (n= 75)	Number	%
BMI (> 25)	23	30.7
Waist/Hip Ratio (Males >0.9 , Females >0.8)	33	44.0
Diabetes mellitus	31	41.3
Smoking	20	26.7
Dyslipidemia	29	38.7

Patients with stage III BP are more likely to develop abnormal retinal changes with $p=0.087$.

Table – 3 Electrocardiographic changes regarding LVH

ECG (n= 75)	Number	%
Sokolow-Lyons Positive [SLC] ($\geq 35\ mm$)	27	36.0
Romhilt-Estes scores [RES] (≥ 5 points)	21	28.0

It was observed in the present study that female patients are more likely to develop increased LVMI compared to male counterparts with $p=0.028$.

Table – 4 Comparison SLC and RES with Echo LVH

Diagnostic statistics	SLC as compared to echo LVH	RES as compared To echo LVH
Sensitivity	47.22	41.67
Specificity	62.96	84.62

PPV	74.36	71.43
NPV	60.42	61.11
Accuracy	61.33	64.00
Kappa	0.22	0.27

Both SLC and RES are having less diagnostic value in comparison to Echo for LVH, however the RES is better than SLC in term of Accuracy and Kappa co-efficient of agreement.

Table - 5 Association of GFR with stage of blood pressure

Stage of Blood Pressure	GFR ≤ 60	GFR > 60
Stage I	5	22
Stage II	8	19
Stage III	14	7

The P value is 0.00181, which is statistically significant.

DISCUSSION

Isolated systolic hypertension is the commonest cause of raised blood pressure in older population. A common misconception among patients and practitioners is that elevated diastolic blood pressure is more important than elevated systolic pressure. In fact one of the key messages of JNC VII is in persons older than 50 years, systolic blood pressure of more than 140 mmHg is a much more cardiovascular risk factor than DBP.

The present study aims at detecting end organ complications of ISH mainly cardiac through ECG & ECHO in term of left ventricular hypertrophy, regional wall abnormalities, reduced ejection fraction, which may result /manifest as CAD, MI & CCF, and on kidney by measuring creatinine clearance.

Study comprised of 75 patients out of which 49 were males and 26 females. Male constituted 65.33% females constituted 34.67%. Mean age was 71.236. 47%, with male to female ratio 1.91: 1. 53.33% of the patients presented were in the age group of 66-75 years.

Symptomatology: In the present study 57.5% of the patient's were asymptomatic. Breathlessness was the major symptom (34.7%) followed by swelling of the limbs (25.3%), chest pain (20%), headache (14.7%), blurred vision, giddiness (5.3%), palpitation motor weakness (2.7%). Stage III blood pressure group constituted more of the symptomatic patients (7.6 times significantly more).

Anthropometry: In the present study average weight for females was 58.08 kg, average BMI was 25.99 $\pm$ 5.06 compared to male (Weight 61.98 kg and BMI 23.17 $\pm$ 5.09). Average waist/hip ratio for females was 0.92 $\pm$ 0.15 compared to males 0.82 $\pm$ 0.16 BMI and waist hip ratio were significantly more in the females.

Associated Risk Factors: 30.7% of the patients had BMI more than 25 (overweight). 44% of the patients had waist by hip ratio more than (0.9) in males and 0.8 in females. 41.3% of the patients had diabetes mellitus.

Dyslipidaemia: 38.7% of the patients had dyslipidaemia. Among these patients, 28% of the patients had isolated low HDLC (normal HDLC $> 50\ mg/dl$ for females & $\geq 40\ mg/dl$ for males). 20% had high isolated triglycerides ($> 150\ mg/dl$) remaining patients had combined low HDLC and high triglycerides and LDL $> 100\ mg/dl$.

Association of risk factors with increased left ventricular mass index: In the present study waist/hip ratio was positively related with the increased LVMI, with $p=0.004$. The patients with DM were 4.06 times more likely to develop increased LVMI, with $p=0.150$.

Retinal Changes: Out of 75 patients, 30 patients had developed retinal changes. 13.3% for focal arteriolar narrowing, 12% had arteriovenous nipping and about 8% for retinopathy. Patients with stage III blood pressure were found to be 3.06 times more likely to develop abnormal retinal changes.

Renal changes: Out of 75 patients, 27 patients had GFR $< 60\ ml/min/1.73\ m^2$. Prevalence of CKD in present study was 36%, out of which 6.6% of patients were in stage 1, 10.6% in stage 2 and 18.6% in stage 3 hypertension group.

Echocardiographic changes: In the present study echocardiographic

changes seen are, increased Left Ventricular Mass Index in 48%, among them 16% had LV volume >90ml/m². 28% of the patients had regional wall motion abnormalities. 26.7% of the patients had reduced ejection fraction (< 56%).^[10] Devereux RB et al found echocardiographic evidence of LVH in the general population. Left ventricular mass indexed to body surface area gives left ventricular mass index.

Overall sensitivity of echo is nearly 100%; specificity is 86% contrasting with ECG sensitivity of 20-50%. In Penn convention the thickness of endocardial surfaces is excluded from measurement of septal and posterior wall thickness and is included in measurement of LVID. In ASE convention, the thickness of endocardial surface is included in the measurement of septal and posterior wall thickness and is excluded from the measurement of LVID. By Penn Convention - left ventricular mass is calculated by

Left Ventricular Mass = $1.04 [(LVIDD + LVPWTD + IVSTD)^3 - (LVIDD)^3] - 13.6$

ASE-Cube Left Ventricular Mass = $1.04 [(LVIDD + LVPWTD + IVSTD)^3 - (LVIDD)^3]$

The constant 13.6 is subtracted to correct for absence of myocardium at the base. Penn-cube left ventricular mass correlated closely with necropsy left ventricular mass and over-estimated it by only 6%; sensitivity of 100% and specificity of 86%. ASE-Cube left ventricular mass correlates similarly to necropsy left ventricular mass, but over estimates it by 25%. The over estimation can be corrected by the equation. Left Ventricular Mass = 0.80 (ASE-Cube Left Ventricular Mass) + 0.6 g. Use of ASE measurements in the volume correction formula underestimates necropsy left ventricular mass by a mean of 30%.

Association of stage of blood pressure with increased LVMI: It is observed in the present study that 19.4% of the patients in stage I blood pressure were having increased LV mass, 41.7% of the patients in stage II and 38.9% in stage III blood pressure. It is observed that as the stage of blood pressure increases, incidence of increased LV mass also increases significantly with $p=0.044$. It is also observed that female patients were more likely to develop increased LVMI compared to male counterparts with $p=0.028$.

CONCLUSION

ISH is the commonest cause of high blood pressure in the elderly. The incidence increases with age advancement. It is more risky in nonwhite population. Waist/hip ratios, DM, dyslipidaemia are significant associated risk factors. ISH associated with risk factors has definite effect mainly on cardia in terms of LV hypertrophy. And on kidney in terms of reduced creatinine clearance is an even better predictor of morbidity and mortality than is diastolic blood pressure. Several large trials have documented a clear benefit to treating ISH. Even small reductions in BP have a substantial impact on patient outcome. Hence, ISH in elderly to be detected early, treated promptly so as to prevent / reduce cardiovascular, renal morbidity and mortality in our growing elderly population ISH associated with risk factors has definite effect mainly on cardia in terms of LV hypertrophy and on kidney in terms of reduced creatinine clearance is an even better predictor of morbidity and mortality than is diastolic blood pressure. Several large trials have documented a clear benefit to treating ISH. Even small reductions in BP have a substantial impact on patient outcome. Hence, ISH in elderly to be detected early, treated promptly so as to prevent / reduce cardiovascular, renal morbidity and mortality in our growing elderly population.

REFERENCES

1. Boon D. ST segment depression criteria the prevalence of silent cardiac ischaemia in hypertension. *Hypertension*, 2003; 41: 476-81.
2. Devereux RB. Cost effectiveness of echocardiography & electrocardiography for detection of LVH in hypertension. *Hypertension*, 1987; 9 (Suppl 2): 69.
3. Dunn FG, Chandraratna P, De Carvalho JGR, Basta LL, Frohlich ED. Sudden cardiac death, ventricular arrhythmias and hypertensive left ventricular hypertrophy. *JHypertens*, 1993; 11: 1003-1010
4. Dwivedi S, Singh G, Agarwal MP. Profile of hypertension in elderly subjects. *JAPI*, 2000; 48 (11): 1045-6.
5. Frohlich ED, Tarzi RC, Miller A et al. Risk mechanism in hypertension heart disease. *Hypertension*, 1999; 34: 782-89.
6. Hazarika NC, Biswas D, Mahata J. Hypertension in the elderly populations of Assam *JAPI*, June 2003; 51.
7. J. Hunter Young, Michael J. Klag, Paul Muntner, Joanna L. Whyte, Marco Pahor, and Josef Coresh. Blood Pressure and Decline in Kidney Function: Findings from the Systolic Hypertension in the Elderly Program (SHEP) *J. Am. Soc. Nephrol.* 2002 13: 2776-2782.
8. JNC on prevention, detection, Naomi DC, Fisher, Gordon H. Williams. The Hypertensive vascular disease: 16ed, Harrison's Principles of Internal Medicine Vol. II,

1436.

9. Malhotra P, Savitakumari Singh S, Verma S. Isolated lipid abnormalities in rural and urban normalities and hypertension. *Northwest Indians JAPI*, 51: May 2003.
10. Messerli FH. Hypertension in special populations. *Med Clin North Am*, 1997; 81(6): 1335-45
11. Michael A. Colandrea, Gary. D. Systolic hypertension in the elderly. An epidemiologic assessment circulation. *Am J Med* 1991; XL190 (4B): 145-205.
12. Nwachukwu, D. C., Agu, U. F., & Nwachukwu, N. Z. (2016). Pattern of renal impairment among hypertensive subjects in Umuahia, South East, Nigeria *International Journal of Medicine and Medical Sciences*, 8(5), 45-50.
13. Thomas Kuruvilla. Prevalence of hypertension in the Parsi community of Mumbai. *JAPI*, Nov 2001; 49: 1130
14. Vrinda Kulkarni, Bhagwat N, Avi Hakim, Sandhya Kamat, Soneji SL. Hypertension in elderly, *JAPI*, 49: September 2001.
15. Wilking SV, Belanger A, Kannel WB et al. Determinants of isolated systolic hypertension. *JAMA*, 1988; 260 (23): 3451-5.
16. Wong TY, Klein R, Sharrett AR. The prevalence & risk factors of retinal microvascular abnormalities in older persons: The cardiovascular health study. *Ophthalmology*, April 2003; 110 (4): 658-64.