FIRST TRIMESTER MATERNAL BMI AND PREGNANCY OUTCOME AT DMCH, LAHERIASARAI, BIHAR

Gynaecology

Dr. Supriya Kumari	MBBS, M.S. (Obs.&Gynae.), Senior Resident, Department of Obstetrics and Gynaecology, Darbhanga Medical College & Hospital, Laheriasarai, Bihar.
Dr. Surya Narayan*	MBBS, M.D. (Radio-diagnosis), Senior Resident, Department of Radio-diagnosis, Darbhanga Medical College & Hospital, Laheriasarai, Bihar. *Corresponding Author
Dr. [Prof.] Kumudini Jha	Professor and Head of Department, Department of Obstetrics and Gynaecology, Darbhanga Medical College & Hospital, Laheriasarai, Bihar.
Dr. Debarshi Jana	Young Scientist (DST) Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata - 700020, West Bengal, India

ABSTRACT

AIM: The aim of the study was to find out relationship of maternal BMI to pregnancy outcome. **METHODOLOGY:** The study carried out in Department of Obstetrics and Gynaecology, Darbhanga Medical College and Hospital, Laheriasarai, Bihar from February 2019 to January 2020 and pregnancy outcomes were analyzed in relation BMI recorded in first trimester of primigravida. **RESULTS:** Mean age of pregnant women was 26.2 years. At first booking obese women were significantly older (28.0 years) than others. Family history of diabetes was significantly higher among obese women (8.25%) compared to others. Rates of cesarean sections were higher in obese compared to others. The macrosomia rates were higher in obese compared to other group. The preeclampsia (1.89%), were significantly higher in obese women than others. **CONCLUSIONS:** Obese women were at a high risk of developing adverse pregnancy outcomes in terms of gestational diabetes, macrosomia, preeclampsia more of cesarean section. Normal weight women have low risk for cesarean section and macrosomia. These results highlight the need for preconception counseling, especially for obese and overweight and have beneficial outcomes in Asian Indian women.

KEYWORDS

Body Mass Index, Primigravida, Macrosomia, Preterm Birth, Preeclampsia, Cesarean Section

INTRODUCTION

Now a day's increase in the prevalence of obesity is seen. High prepregnancy BMI and/or excessive gestational weight gain (GWG) have negative implications on pregnancy outcomes, and this amplifies the burden of chronic diseases putting health of both mother and the infant at risk. Recent studies have reported that prepregnancy BMI is associated with birth weight of the infant. Women who are overweight/obese are at high risk of developing preeclampsia and gestational diabetes mellitus. This in turn is found to influence the neonatal outcomes such as perinatal mortality, macrosomia, and congenital anomalies. In addition, maternal obesity leads to higher cesarean sections and increased risk of anesthesia. The long-term impact of maternal obesity includes maternal weight retention and exacerbation of obesity. In 2009, the Institute of Medicine (IOM), USA, published revised GWG guidelines that are based on prepregnancy ranges for underweight, normal weight, overweight, and obese women. These recommendations were, however, based on American women and therefore its generalizability to other populations is unclear as maternal anthropometry varies across different populations. The IOM later published a commentary acknowledging that they based their recommendations primarily on the basis of primigravida mothers of high social status and those with no physical activity. Some recent studies from Asia have concluded that IOM guidelines are suitable for the Asian population. Whereas others have reported that GWG among Asians is different from what has been recommended by the IOM. There are very few studies from India that have looked at the applicability of the IOM guidelines in pregnant women. This is mainly because the BMI classification for Asians is different from the World Health Organization (WHO) BMI cutoffs recommended for the West.

MATERIAL AND METHODS

A total of 1506 records of pregnant women attending Obstetrics and Gynaecology Department of Darbhanga Medical College and Hospital, Laheriasarai, Bihar from February 2019 to January 2020 were selected. Of the total 1506, a total of 1018 (67.60%) records were available for data analysis after data cleansing. Weight and BMI at early pregnancy in primigravida which was the only feasible option for obtaining reliable information was classified according to WHO Asia Pacific BMI cut points.

Table – 1: WHO Asia Pacific criteria of BMI

Weight category	BMI (kg/m ²)
Under weight	<18.5
Normal weight	18.5-22.9
Over weight	23-24.9
Obese	>25

BMI = Body mass index, WHO = World health of organization
Pregnancy outcomes were analyzed in relation to BMI.

Definitions

Preterm birth or premature birth is one that occurs before the start of 37 weeks of pregnancy. Macrosomia is used to describe a newborn who is significantly larger than average; a infant weighing >3.5 kg is considered macrosomic.

Statistical analysis

All analyses were done using Windows-based SPSS statistical package estimates were expressed as mean $\pm$ standard deviation. To compare adverse pregnancy outcomes Chi-square tests were used to test differences in proportions. $P < 0.05$ was considered statistically significant.

RESULTS

Mean age of pregnant women was 26.2 years, mean BMI was 25.8 kg/m², and mean weight gain during pregnancy was 8.1 kg. Family history of type 2 diabetes was seen in 7.1%.

Table – 2: Distribution of BMI in study group

WHO Asia Pacific criteria of BMI	Distribution (n=1018)
Under weight	57 (5.60%)
Normal weight	321 (31.53%)
Over weight	216 (21.21%)
Obese	424 (41.66%)

As shown in Table 2, 5.60% (n = 57) were underweight, 31.53% (n = 321) normal weight, 21.21% (n = 216) overweight, and the rest 41.66% (n = 424) were obese.

Table 3: Clinical characteristics in study group

Clinical parameter	Underweight (n = 57)	Normal (n = 321)	Overweight (n = 216)	Obese (n = 424)
--------------------	----------------------	------------------	----------------------	-----------------

Mean age	25.1	27.3	27.5	28.0
Mean weight at booking (Kg)	46.2	53.4	59.3	69.5
Mean weight gain during pregnancy	9.2	9.0	8.8	8.4
Mean BMI at booking (Kg/m ²)	17.1	21.1	23.9	28.8
Mean birth weight (Kg)	2.89	3.03	3.05	3.08
Preterm deliveries (%)	3 (5.26)	19 (5.92)	14(6.48)	30(7.08)
Gestational diabetes (%)	2 (3.51)	15 (4.67)	12 (5.56)	35 (8.25)
Cesarean rates (%)	6 (10.53)	82(25.55)	65(30.09)	170(40.09)
Low birth weight (%)	7 (12.28)	37(11.53)	23(10.64)	43(10.14)
Macrosomia (%)	3 (5.26)	25(7.79)	25(11.57)	59 (13.92)
Preeclampsia (%)	1(1.75)	1(0.31)	2(0.93)	8 (1.89)

Table 3 shows the clinical characteristics and pregnancy outcomes of the women classified according to their BMI at first booking in primigravida. Obese women were significantly older (28.0 years) than overweight, normal weight, and underweight women (27.5, 27.3, and 25.1 years, $P < 0.0001$, respectively). Weight at booking (kg) was significantly higher in obese women when compared to the other three groups.

The mean birth weight of the infants born to normal weight, overweight, and obese women was significantly higher than birth weight of infants of underweight women (3.03 kg, 3.05 kg, 3.08 kg, and 2.89; $P = 0.006$, respectively). Gestational diabetes was significantly higher among obese women (8.25%) compared to the overweight, normal weight, and underweight women (5.56%, 4.67%, and 3.51%, $P < 0.001$, respectively). Rates of cesarean sections were (40.09%, 30.9%, 25.55%, and 10.53%, $P=0.001$) respectively in obese, overweight, normal and underweight women. The low birth weight rates were 12.28%, 11.53%, 10.64%, 10.14% respectively in underweight, normal, overweight and obese women. The macrosomia rates were (13.92%, 11.57%, 7.79%, and 5.26%, $P = 0.001$), and preeclampsia (1.89%, 0.93%, 0.31%, and 1.75%, $P = 0.004$) were significantly higher in obese women than overweight, normal weight, and underweight women, respectively.

The risk of preterm birth was higher in all the groups although this did not reach statistical significance.

Table – 4: IOM recommended weight gain in pregnancy.

Weight category/BMI	IOM recommended weight gain (kg)
Under weight	12-18
Normal weight	11.5-16
Over weight	7-11.5
Obese	5-9

BMI = Body mass index, IOM = Institute of Medicine

Table – 5: Shows weight gain during pregnancy across the different BMI categories in our study group

Weight category	Recommended weight gain (kg)	More than recommended weight gain (kg)	Less than recommended weight gain (kg)
Under weight (n=57)	17(29.84%)	3(5.26%)	37(64.90%)
Normal weight (n=321)	89(27.73%)	27(8.41%)	205(63.86%)
Over weight (n=216)	52(20.08%)	18(8.33%)	146(67.59%)
Obese (n=424)	154(36.32%)	128(30.19%)	142(33.49%)

It was seen that 29.84% of underweight, 27.73% of normal weight, 20.08% overweight, and 36.32% obese women met the recommendations for weight gain. While majority of underweight (64.90%), normal weight women (63.86%), and overweight women (67.59%) gained weight less than recommended, among obese women, 30.19% of them gained more weight and 33.49% of them gained less than the recommended weight.

DISCUSSION

Our study shows the following findings:

1. Obese women were at a higher risk of preterm delivery, cesarean section, macrosomia, and preeclampsia.
2. Normal weight women had less risk of a cesarean section and macrosomia.
3. Only 30% of women gained weight as per the recommendations.

In Asian population, the prevalence of obesity may be lower than in Europe; the health risks associated with obesity occur at a lower BMI in Asians than compared to the West, thereby making WHO BMI categorization, less relevant to the Asian population. Hence, in 2000, the Regional Office for the Western Pacific of WHO, the International Association for the Study of Obesity, and the International Obesity Task Force together released, the Asia-Pacific Perspective for redefining obesity suggesting diagnostic criteria to identify overweight and obesity in the Asian population^[16]. Maternal obesity is a risk factor for several pregnancy-related complications which may have adverse effects on both the mother and her infant. Obese women have an increased risk of miscarriage early in pregnancy^[19]. A prospective multicenter study of more than 16,000 pregnancy women showed that obese women were 2.5 times and 1.6 times more likely to develop gestational hypertension and preeclampsia, respectively. In addition to these obstetric problems, obese women are also at an increased risk of undergoing cesarean sections. Fetal overgrowth also is another major concern in obese women. In Denmark, the percentage of macrosomic infants increased from 16.7% to 20.9% in 10 years, and increasing maternal BMI has been implicated to be one major factor. Several other studies have also shown an association between increasing BMI, cesarean section, and macrosomia.

Cesarean section is usually influenced by several factors, such as practice behavior of the obstetrician or other pregnancy complications in obese women, may necessitate the need for cesarean section. In our study show that obese women were at a significantly higher risk of cesarean delivery. Our results also show that obese women were at a high risk of preterm labor and macrosomia. Not many studies have reported its association to preterm labor. However, a few studies that have shown association between obese women and preterm labor lacked statistical significance. Some studies have also reported a negative association between obesity and preterm labor. These differences may be due to inconsistency in the definition of preterm birth and effect of other confounding variables. Underweight women are known to deliver preterm infants. In addition, under weight women were shown to be at two-fold risk of delivering low birth weight infants than those who met the recommendations. Our results showed that though the risk for low birth weight in underweight women was high, it was not statistically significant. This could be because the number of underweight women studied is less. Although previous studies have reported a higher percentage (26%) of low birth weight infants. the scenario seems to be fast changing owing to the quality of antenatal care available to pregnant women in the recent past. The other important finding in this study is that though a major proportion of normal weight had a less risk for cesarean sections and macrosomia which is in contrast with previous studies which showed that these women had increased risk for such complications. This difference could be attributed mainly due to the different BMI criteria used. Limitations of the study are like small sample size and BMI measurement done in first trimester where some women will lose the weight because of excessive vomiting. Ideally should be based on prepregnancy weight. However, as seen in several other studies that data are seldom available in routine antenatal records. Some studies rely on the woman's recall of prepregnancy weight.

CONCLUSION

Preconception counseling, especially for obese and overweight women, emphasizing the importance of proper physical activity and healthy eating to avoid excess weight and its adverse effects, could have beneficial outcomes in pregnant Indian women.

REFERENCES

1. American College of Obstetricians and Gynecologists. ACOG Committee opinion number 315, September 2005. Obesity in pregnancy. *Obstet Gynecol*. 2005; 106:671-5.
2. Frederick IO, Williams MA, Sales AE, Martin DP, Killien Pre-pregnancy body mass index, gestational weight gain, and other maternal characteristics in relation to infant birth weight. *Matern Child Health J*. 2008; 12:557-67.
3. Gunderson EP, Abrams B, Selvin S. Does the pattern of postpartum weight change differ according to pregravid body size? *Int J Obes Relat Metab Disord*. 2001; 25:853-62.
4. Kelly A, Kevany J, de Onis M, Shah PM. A WHO collaborative study of maternal anthropometry and pregnancy outcomes. *Int J Gynaecol Obstet*. 1996; 53:219-33.

5. Kim SY, Dietz PM, England L, Morrow B, Callaghan WM. Trends in pre-pregnancy obesity in nine states, 1993-2003. *Obesity (Silver Spring)*. 2007; 15:986-93.
6. Lashen H, Fear K, Sturdee DW. Obesity is associated with increased risk of first trimester and recurrent miscarriage: Matched case-control study. *Hum Reprod*. 2004; 19:1644-6.
7. Last accessed on 2016 Dec 18. Available from: <http://www.who.int/mediacentre/factsheets/fs363/en/>.
8. Last accessed on 2016 Oct 20. Available from: <http://www.wpro.who.int/nutrition/documents/docs/Redefiningobesity.pdf>.
9. Liu Y, Dai W, Dai X, Li Z. Prepregnancy body mass index and gestational weight gain with the outcome of pregnancy: 13-year study of 292,568 cases in China. *Arch Gynecol Obstet*. 2012; 286:905-11.
10. Lynch CM, Sexton DJ, Hession M, Morrison JJ. Obesity and mode of delivery in primigravid and multigravid women. *Am J Perinatol*. 2008; 25:163-7.
11. Nohr EA, Vaeth M, Baker JL, Sørensen Tia, Olsen J, Rasmussen KM. Combined associations of prepregnancy body mass index and gestational weight gain with the outcome of pregnancy. *Am J Clin Nutr*. 2008; 87:1750-9.
12. Ota E, Haruna M, Suzuki M, Anh DD, Tho le H, Tam NT et al. Maternal body mass index and gestational weight gain and their association with perinatal outcomes in Viet Nam. *Bull World Health Organ*. 2011; 89:127-36.
13. Paul VK, Deorari AK, Singh M, Parthasarathy A, editor. Management of low birth weight babies. IAP Textbook of Pediatrics. 2nd ed. New Delhi: Jaypee Brothers, 2002.
14. Radhakrishnan U, Kolar G, Nirmalan PK. Cross-sectional study of gestational weight gain and perinatal outcomes in pregnant women at a tertiary care center in Southern India. *J Obstet Gynaecol Res*. 2014; 40:25-31.
15. Rasmussen KM, Abrams B, Bodnar LM, Butte NF, Catalano PM, Maria Siega-Riz A. Recommendations for weight gain during pregnancy in the context of the obesity epidemic. *Obstet Gynecol*. 2010; 116:1191-5.
16. Rasmussen KM, Yaktine AL, editors. Institute of Medicine (US) and National Research Council (US) Committee to Reexamine IOM Pregnancy Weight Guidelines. Weight Gain during Pregnancy: Reexamining the Guidelines. Washington, DC: National Academies Press (US), 2009.
17. Sebire NJ, Jolly M, Harris JP, Wadsworth J, Joffe M, Beard RW et al. Maternal obesity and pregnancy outcome: A study of 287,213 pregnancies in London. *Int J Obes Relat Metab Disord*. 2001; 25:1175-82.
18. Weiss JL, Malone FD, Emig D, Ball RH, Nyberg DA, Comstock CH et al. Obesity, obstetric complications and cesarean delivery rate – A population-based screening study. *Am J Obstet Gynecol*. 2004; 190:1091-7.
19. Wong W, Tang NL, Lau TK, Wong TW. A new recommendation for maternal weight gain in Chinese women. *J Am Diet Assoc*. 2000; 100:791-6.
20. Yang YD, Yang HX. Investigation into the clinical suitability of Institute of Medicine 2009 guidelines regarding weight gain during pregnancy for women with full term singleton fetus in China. *Zhonghua Fu Chan Ke Za Zhi*. 2012; 47:646-50.
21. Young TK, Woodmansee B. Factors that are associated with cesarean delivery in a large private practice: The importance of prepregnancy body mass index and weight gain. *Am J Obstet Gynecol*. 2002; 187:312-8.
22. Orskou J, Kesmodel U, Henriksen TB, Secher NJ. An increasing proportion of infants weigh more than 4000 grams at birth. *Acta Obstet Gynecol Scand*. 2001; 80:931-6.
23. Zhong Y, Cahill AG, Macones GA, Zhu F, Odibo AO. The association between prepregnancy maternal body mass index and preterm delivery. *Am J Perinatol*. 2010; 27:293-8.
24. Hendler I, Goldenberg RL, Mercer BM, Iams JD, Meis PJ, Moawad AH et al. The Preterm Prediction Study: Association between maternal body mass index and spontaneous and indicated preterm birth. *Am J Obstet Gynecol*. 2005; 192:882-6.
25. Masho SW, Bishop DL, Munn M. Pre-pregnancy BMI and weight gain: Where is the tipping point for preterm birth? *BMC Pregnancy Childbirth*. 2013; 13:120.
26. Jeric M, Roje D, Medic N, Strinic T, Mestrovic Z, Vulic M. Maternal pre-pregnancy underweight and fetal growth in relation to institute of medicine recommendations for gestational weight gain. *Early Hum Dev*. 2013; 89:277-81.
27. Yu Z, Han S, Zhu J, Sun X, Ji C, Guo X. Pre-pregnancy body mass index in relation to infant birth weight and offspring overweight/obesity: A systematic review and meta-analysis. *PLoS One*. 2013; 8:e61627.
28. Wills AK, Chinchwadkar MC, Joglekar CV, Natekar AS, Yajnik CS, Fall CH et al. Maternal and paternal height and BMI and patterns of fetal growth: The Pune Maternal Nutrition Study. *Early Hum Dev*. 2010; 86:535-40.
29. Thorsdottir I, Torfadottir JE, Birgisdottir BE, Geirsson RT. Weight gain in women of normal weight before pregnancy: Complications in pregnancy or delivery and birth outcome. *Obstet Gynecol*. 2002; 99(5-1):799-806.