COMPARISON OF MATERNAL AND NEONATAL OUTCOME IN FORCEPS VERSUS VENTOUSE ASSISTED VAGINAL DELIVERY IN OBSTETRICS AND GYNAECOLOGY DEPARTMENT OF PMCH, PATNA, BIHAR

Gynaecology

Dr. Archana Kumari

MBBS, M.S. (Obs. & Gynae.), Assistant Professor, Department Obstetrics and Gynaecology, Patna Medical College and Hospital, Patna, Bihar.

Dr. [Prof.] K. Manju*

Professor, Department Obstetrics and Gynaecology, Patna Medical College and Hospital, Patna, Bihar. *Corresponding Author

Dr. Debarshi Jana

Young Scientist (DST) Institute of Post-Graduate Medical Education and Research, A.J.C. Bose Road, Kolkata-700020, West Bengal, India.

ABSTRACT

Objectives: To compare the maternal and neonatal outcome following ventouse and forceps- assisted vaginal deliveries in singleton term pregnancies. **Study Design:** A cross-sectional study. Duration and Place of study: It was conducted at the Obstetrics and Gynaecology Department of Patna Medical College and Hospital, Patna, Bihar from May 2020 to December 2020. **Patients and Methods:** All patients who underwent instrumental vaginal delivery during this period, fulfilling the inclusion criteria were studied. Main maternal outcomes were perineal tears while the main neonatal outcomes were Apgar score at 1 minute and at 5 minutes after birth. Neonatal outcome included record of cephalhematoma, shoulder dystocia and need for admission to NICU. In addition to the above, maternal age, gestational age, parity and booking status were also recorded along with the indication for instrumental vaginal delivery for both ventouse and forceps delivery groups. Data was entered into SPSS-15. Percentages were calculated for all the above variables except maternal age and gestational age for which mean + SD were calculated. Chi-square test was used for qualitative variables to assess any difference between the two groups. P-value of equal to or <0.05 considered to be significant. **Results:** A total of 80 patients underwent ventouse delivery while 120 patients with forceps delivery were included. Mean maternal ages were 25.7+4.33 yrs and 25.05+3.47yrs in ventouse and forceps group respectively. 54(67%) and 95(79.16%) were primiparous in ventouse and forceps group respectively. 35(43.7%) were booked cases in ventouse group while this number was 53(44.2%) in forceps group. Fetal distress was the commonest indication of instrument application in 60(75%) and 108(90%) cases in ventouse and forceps group respectively. 16(20%) and 35(29.16%) neonates had Apgar score of <7 at 1 minute in ventouse and forceps group respectively. Apgar score of <7 at 5 minute was observed in 10(12.5%) of ventouse group while 14(11%) of forceps group. 2.5 % (n=2) and 1.66% (n=2) of neonates had shoulder dystocia in ventouse and forceps group respectively. Cephalhematoma was observed in 4(5%) of ventouse group while it was 1.5% (n=2) in forceps group. 11% (n=9) and 10% (n=12) of neonates admitted to NICU in ventouse and forceps group respectively. No case of maternal 3rd degree perineal tear observed in ventouse group while 3(2.5%) of patients had 3rd degree perineal tears in forceps group. **Conclusion:** Forceps deliveries are more associated with maternal morbidity while neonatal trauma is commoner among those delivered by vacuum extraction however there seems to be no difference in neonatal admission to NICU.

KEYWORDS

Maternal And Neonatal Outcome, Forceps, Ventouse.

INTRODUCTION

The history of forceps delivery dates back as far as 1500 BC, being originally used following fetal demise to save the mother's life. The ventouse however was first described in 1954 and modified in 1969. The incidence of instrumental vaginal delivery ranges between 10-15% in UK and depends upon institution and the population. Commonest indications of instrumental vaginal delivery include delay in the second stage of labor, poor maternal effort and fetal distress including cord prolapse in the second stage of labor.

Maternal indications include severe cardiac, respiratory or hypertensive disease or intracranial pathology where bearing down effort may be detrimental for her health. The use of forceps in practice since decades however the trend shift is seen in the direction of ventouse. The rapid acceptance of the ventouse is due to perceived lesser incidence of maternal trauma and minimal training requirements.

Review of the literature suggests differential maternal and neonatal outcomes and complication rates between the two methods. Both are associated with increased risk of maternal and neonatal injury when compared to normal spontaneous vaginal deliveries. There are studies favoring the time tested outlet forceps to be better instrumental method but on the other hand there is some evidence which suggest its pitfalls. The literature also include studies in which forceps delivery has been termed as better modality of operative vaginal delivery in terms of neonatal outcome. The earlier work focused on the maternal side of problems, including the increased frequency of maternal tears, soft tissue damages and post-delivery scarring resulting in the development of procedures like vacuum extraction. However, recent studies show that maternal soft tissue injury rates are similar in vacuum and forceps assisted deliveries. Thus, there is considerable debate concerning the preferred instrument for assisted vaginal delivery. The objective of this study was to compare the short term neonatal and maternal outcomes after ventouse and forceps deliveries.

MATERIAL AND METHODS

All patients who underwent instrumental vaginal delivery during the period May 2020 to December 2020, having singleton pregnancy at gestation of 37 weeks or more with cephalic presentation were included in the study. Patients having fetal anomalies and diagnosed IUD were excluded from the study. Main maternal outcomes were presence of 3rd or 4th degree perineal tears while the main neonatal outcomes were Apgar score at 1 minute and at 5 minutes after birth. Neonatal outcome also included presence of cephalhematoma, shoulder dystocia and need for admission to NICU. In addition to the above, maternal age, parity and booking status were also recorded along with the indication for instrumental vaginal delivery for both ventouse and forceps delivery groups. Data was entered into SPSS-15. Percentages were calculated for all the above variables except maternal age for which mean + SD were calculated. Chi-square test was used for qualitative variables to assess any difference between the two groups. P-value of equal to or <0.05 considered to be significant.

RESULTS

A total of 80 patients who underwent ventouse delivery while 120 patients who underwent forceps delivery were included in the study. The mean ages in ventouse and forceps groups were 25.7+4.33yrs, and 25.05+3.47yrs respectively. Majority of patients belonged to the age groups >25years in both groups. Age distribution is depicted in table 1. 67% of patients were primigravidas in ventouse group while this percentage was 79.16% in forceps group. Regarding the booking status of patients, they were predominantly unbooked in both the groups i.e.: 45(56.2%) in ventouse while 67(55.8%) in forceps group.

Table 1 : Maternal features in vacuum and forceps group

Maternal features	Ventouse (n=80)	Forceps (n=120)	P value
Maternal age (yrs)	25.7±4.33	25.05±3.47	
<25 years n (%)	31 (37.5%)	42 (35%)	0.89
≥25 years n (%)	49 (61.25%)	78 (65%)	0.99
Parity			

Primipara n (%)	54 (67%)	95 (79.16%)	0.10
Multipara n (%)	26 (33%)	25 (20.84%)	0.40
Booking status			
Booked cases n (%)	35 (43.7%)	53 (44.2%)	0.97
Unbooked cases n (%)	45 (56.2%)	67 (55.8%)	0.97

Table 2 : Maternal features in vacuum and forceps group

Indication	Ventouse (n=80)	Forceps (n=120)
Fetal distress n (%)	60 (75%)	108 (90%)
Maternal exhaustion n (%)	14 (17.5%)	7 (5.8%)
Prolonged second stage of labor n (%)	6 (7.5%)	5 (4.16%)

Table 3 : Maternal and Neonatal morbidity

Morbidity	Ventouse (n=80)	Forceps (n=120)	P value
Apgar score <7 at 1 minute n (%)	16 (20%)	35 (29.16%)	0.99
Apgar score <7 at 5 minute n (%)	10 (12.5%)	14 (11.66%)	0.76
NICU admission n (%)	9 (11%)	12 (10%)	0.70
S Shoulder dystocia n (%)	2 (2.5%)	2 (1.66%)	0.36
Cephalhematoma n (%)	4 (5%)	2 (1.5%)	0.13
Maternal 3rd degree tear	0	3 (2.5%)	0.00

The indications for ventouse application were fetal distress (75%), maternal exhaustion (17.5%) and prolonged second stage of labor (7.5%). Indications for forceps delivery were fetal distress (90%), maternal exhaustion (5.83%) and prolonged second stage of labor (4.16%). Apgar score of <7 at 1 minute after birth was observed in 16(20%) and 35(29.16%) of cases in ventouse and forceps delivered neonates. Apgar score of <7 at 5 minutes after delivery was noted in 10(12.5%) and 14(11.6%) in ventouse and forceps group respectively. Shoulder dystocia was observed in 2.5% (n=2) of ventouse deliveries while it was 1.66% (n=2) of forceps deliveries. Cephalhematoma was noted in 5 % (n=4) of ventouse delivered newborns while it is 1.5% (n=2) in forceps delivered newborns. 11% (n=9) of ventouse delivered neonates while 10% (n=12) of those delivered by forceps were admitted in NICU. 2 (n=3) of mothers who underwent forceps delivery had 3rd degree perineal tears while no patient had 3rd degree perineal tear in ventouse group.

DISCUSSION

Vacuum extraction and obstetric forceps are operative procedures used during complicated deliveries and it remains an important clinical procedure to be performed by obstetricians. The aim of instrumental vaginal delivery is to assist the vaginal birth ensuring minimum maternal and neonatal morbidity. So, an adequate clinical skill is required for use of both vacuum and forceps. This study was performed to focus especially on the short term maternal and neonatal morbidity associated with forceps and vacuum assisted vaginal deliveries. In the present study, mean ages of the patients undergoing for forceps and vacuum were 25.05±3.47yrs and 25.74±3.33yrs respectively which is comparable with the study performed by N Prapas et al.

43.7% and 44.2% of patient were booked in vacuum and forceps group respectively. The higher frequency of un-booked cases depicts the type of patient coming to tertiary care hospital. It also indicates the attitude of society towards care of pregnant women.

Fetal distress was the commonest indication of forceps deliveries making 90% of all forceps deliveries followed by maternal exhaustion (5.8%) and prolonged second stage (4.16%) respectively. Among the vacuum delivery, 75% were due to fetal distress, 17.5% were due to maternal exhaustion and 7.5% were due to prolonged 2nd stage of labor. These finding were supported by local studies performed by Arshad A and also by Akhtar S.

Several studies documented differential maternal and neonatal outcomes and complication rates between the two methods of operative vaginal delivery. In this study, the Apgar scores were calculated at 1 minute and 5 minutes after birth. 20% and 29.16% of neonates had Apgar score of <7 at 1 min in vacuum and forceps group respectively which is comparable to the study performed by Akhtar S. The Apgar score at 5 minutes of <7 were 12.5% in vacuum while 11.66% in forceps group. These findings were consistent with a study conducted by Singh A and Rathore P. 10% of neonates in forceps group while 11% of neonates in vacuum group had to be admitted in NICU. Supporting this, Singh has reported an equal incidence of neonatal unit admission among the two groups.

Shoulder dystocia is an obstetrical emergency and found to be high with vacuum use compared to forceps. The frequency of shoulder dystocia was 2.5% in vacuum and 1.66% in forceps group. This finding is consistent with the results of research done by Caughey AB et al. Cephalhematoma occurred more frequently in vacuum i.e; 5% as compared to 1.5% in forceps deliveries. These results are the supported by studies conducted by Singh A, N Prapas et al and Caughey AB et al. Maternal third degree perineal tears are more common with the use of forceps. The frequency of 3rd degree tears was 0% with vacuum while 2.5% with forceps in our study.

The present study was not without its shortcomings such as retrospective nature of the data and the absence of research on long term complications associated with instrumental vaginal delivery.

CONCLUSION

Forceps deliveries are more associated with maternal morbidity while neonatal trauma is commoner among those delivered by vacuum extraction however there seems to be no difference in neonatal admission to NICU. In conclusion, our study shows that vacuum and forceps are safe alternatives to each other rather than rivals. The main factor determining the safety of instrument is the obstetrician rather than the instrument itself.

REFERENCES

- Walsh J, Murphy DJ. Operative vaginal delivery- a dying art? Recent advances in Obstetrics and Gynecology, 24. 1 Wimpole street, London W1G 0AE, UK. Royal society of medicine press Ltd.2008. P.75
- Ambreen A , Anwar K, Ayub MR, Rashid N. Comparison of Neonatal and Maternal Outcome between Forceps and Vacuum Delivery. Pakistan Journal of medical and health sciences.vol.5, issue.4, Oct-Dec 2011.
- Arshad H, Baqai S, Abdullah KN, Iftikhar B. Neonatal outcome with instrumental vaginal delivery: a study at PNS Shifa. pakistan Armed forces Medical Journal 2011; 2 June ISSN 0030-9648.
- N Prapas, I kalogiannidis, S Masoura, et al. Operative vaginal delivery in singleton term pregnancies: Short-term maternal and neonatal outcomes. Hippokratia. 2009 Jan-Mar; 13(1): 41–45.
- Oliszewski J, Oliszewska H, Królikowska B, Preis K. The analysis of complications in forceps delivery in Institute of Obstetrics and Women Disease Medical University of Gdańsk. Ginekol Pol. 2006; 77(9):700-4.
- Melamed N, Yogev Y, Staimetz S, Ben-Haroush A. What happens when vacuum extraction fails? Arch Gynecol Obstet. 2009; 280 (2): 243-8.
- Lurie S, Glezerman M, Sadan O. Maternal and neonatal effects of forceps versus vacuum operative vaginal delivery. Int J Gynecol Obstet. 2005; 89: 293–294.
- O'Grady JP, Pope CS, Hoffman DE. Forceps delivery. Best Pract Res Clin Obstet Gynaecol. 2002; 16: 1–16.
- Murphy DJ, Liebling RE, Verity L, Swingle R, Patel R. Cohort study of the early maternal and neonatal morbidity associated with operative delivery in the second stage of labour. Lancet.2001; 358: 1203–1207.
- Demissie K, Rhoads GG, Smulian JC, et al. Operative vaginal delivery and neonatal and infant adverse outcomes: population based retrospective analysis. BMJ. 2004; 329: 24–29.
- Singh A, Rathore P. A comparative study of feto-maternal outcome in instrumental vaginal delivery. J Obstet Gynaecol India 2011 December; 61(6): 663-666.
- Caughey AB, Sandberg PL, Zlatnik MG, Thiet MP, Parer JT, Laros RK., Jr Forceps compared with vacuum: rates of neonatal and maternal morbidity. Obstet Gynecol. 2005; 106: 908–912.
- Bofill JA, Rust OA, Schorr SJ, et al. A randomized prospective trial of obstetric forceps versus the M-cup extractor. Am J Obstet Gynecol. 1996; 175: 1325–1330.
- Johnson JH, Figueroe R, Garry D, Elimian A, Maulic D. Immediate maternal and neonatal effects of forceps and vacuum - assisted deliveries. Obstet Gynecol. 2004; 103: 513–518.
- Vacca A. Vacuum-assisted delivery. Best Pract Res Clin Obstet Gynaecol. 2002; 16: 17–30.
- Pope CS. Vacuum extraction [online] 2008 [cited 2006 June 7]. Available from: URL: <http://www.e.med>