Temperature influence on infrared spectra of brown coal humic substances

Volkov D.S.,^{1,2} Rogova O.B.,² Proskurnin M.A.¹

¹Chemistry Department of M.V. Lomonosov Moscow State University, Moscow, Russia, dmsvolkov@gmail.com

²Department of Chemistry and Physical Chemistry of Soils, V.V. Dokuchaev Soil Science Institute, Pyzhevsky, Moscow, Russia

Keywords: humic substances, FTIR, infrared spectroscopy, temperature infrared spectra, temperature-induced evolution of band frequencies

DOI:10.36291/HIT.2021.058

IR spectroscopy is a developed method to study humic substances (HS). However, the bands of different HS functional groups overlap, and a resulting HS IR spectrum shows a small number of relatively uninformative broad bands. Changes in the vibrational spectra of solid samples upon heating or cooling provide information on polymorphic transformations, structural phase transitions, intramolecular interactions, and second-order phase transitions. However, even if such changes are absent, temperature-dependent IR spectra may reveal alterations in the band intensities, positions, shape, and width [1]. According to Hooke's law, the decreased nearest neighbor distances imply larger force constants and higher frequencies. Transitions between higher levels are less energetic than transitions between lower, which causes a redshift of band maxima with temperature and band broadening [2]. Thus, the temperature dependence of band parameters in the IR spectrum of a solid has a fundamental nature that is mainly associated with the structural deformation of its crystal lattice. This effect has long been observed and studied in various minerals. According to our data, no similar studies with humic substances have been carried out.

Humic acid sodium salt (Sigma-Aldrich) and potassium humate from German leonardite and Sakhalin leonardite were used. The analysis by IR spectroscopy was carried out on an FTIR spectrometer with a single attenuated total internal reflection attachment with a diamond crystal heated in the range from room temperature to 215 °C.

It was shown that when heated in the range of 25–215 °C in air, the IR spectra of coal humic substances undergo several significant changes. Almost all bands, both belonging to mineral fragments and purely organic, shifted in frequency; this effect is reversible and, when cooled, the bands return to their original frequency. When comparing the samples, some bands demonstrate an almost complete identity of the temperature-induced change, while other bands behave differently. This approach provides a more detailed HS structure analysis without total HS decomposition or destruction.

Funding: This work was supported by The Russian Foundation for Basic Research, project number 19-13-00117.

Acknowledgments: This research was performed according to the Development program of the Interdisciplinary Scientific and Educational School of Lomonosov Moscow State University, "The future of the planet and global environmental change".

References

1. Katon, J.E.; Phillips, D.B. Infrared Spectroscopy at Subambient Temperatures. *Appl. Spectrosc. Rev.* 1973, 7, 1-45.
2. Koike, C.; Mutschke, H.; Suto, H.; Naoi, T.; Chihara, H.; Henning, T.; Jäger, C.; Tsuchiyama, A.; Dorschner, J.; Okuda, H. Temperature effects on the mid-and far-infrared spectra of olivine particles. *A&A* 2006, 449, 583-596