The Potential to use Chloroquine and other 4-Aminoquinoline Analogues to Modulate Persisting Inflammation in Old Age

Stephen Allen^{1,2*} and Divya Tiwari¹

¹Department of Medicine, The Royal Bournemouth Hospital, UK

²Centre for Postgraduate Medical Research and Education, Bournemouth University, UK

Article Information

Received date: Feb 13, 2019

Accepted date: Mar 05, 2019

Published date: Mar 08, 2019

Corresponding author

Stephen Allen, Department of Medicine,
Royal Bournemouth Hospital, Castle
Lane East, Bournemouth, Dorset,
BH3 7DG, UK, Tel: +44 1202 303626;
Fax: +44 1202 704542;
Email: drscallen@aol.com

Distributed under Creative Commons
CC-BY 4.0

Keywords Systemic inflammation;
Ageing; Frailty; Sarcopenia; Immune
modulation; Theophylline; Chloroquine

Article DOI 10.36876/smgr.1020

Abstract

Low-amplitude persisting systemic inflammation is frequently found in elderly people and is causally linked to several markers of frailty, including sarcopenia, low mood and higher dependency, as well as higher all-cause mortality. Older patients with such "inflammation" have raised baseline blood levels of several pro-inflammatory cytokines, such as tumor necrosis factor alpha and interleukin-1 beta, and chronically raised C-reactive protein. There is a clear need to identify interventions, including drugs that can ameliorate such inflammation by helping to re-set the innate immune system to a less inflamed baseline. Several classes of drugs have such properties. In this review the authors have summarized the background science with respect to the relationship between ageing, inflammation and frailty, then described the established role of methyl-xanthines, particularly theophylline, as immune modulating drugs before positing a case for the use of 4-aminoquinolines, such as chloroquine, in a similar role. The probable mechanisms of anti-inflammatory action for those classes of drugs are compared, leading to a proposal that formal clinical trials should be conducted of chloroquine as an adjunctive immune modulator for "inflammaging" and persisting post-acute inflammation in old age.

Introduction

A strong imperative is emerging to search for pharmacological agents that can effectively modify the consequences of persisting low amplitude systemic inflammation, particularly in aged people in whom it has been shown to be causally associated with a pre-frail or overtly frail condition, and is a probable direct etiological factor in the development of sarcopenia [1,2]. This is especially so for the many patients who are physically or mentally unable to perform exercise of sufficient regularity, work-rate and duration to benefit from the anti-inflammatory effects of aerobic exercise [2,3]. This is a familiar clinical phenotype in geriatric medicine and represents a major challenge to clinicians, health service planners and economists. Our proposition is that opportunities should be taken to identify pharmacological and non-pharmacological interventions that can help to reduce the burden of frailty and preserve function. To that end, a case can be made for taking a fresh look at a number of drugs, many of which are in common use for other purposes, that have been shown to modulate inflammatory cellular and biochemical responses, largely by resetting immune networks toward a less inflamed baseline state. To date, these effects have been arguably explored most avidly for methyl-xanthines and statins [4-7]. In this paper the authors will summarize the prevailing evidence for 4-aminoquinolines (4AQs), mainly chloroquine and hydroxychloroquine that have been shown to have immune modulating effects similar to methyl-xanthines, though apparently through different mechanisms. We draw on that evidence to propose a need for properly conducted clinical trials of 4AQs to ascertain any measurable clinical benefits.

Ageing of the Innate Immune System

Innate immunity in older people is fundamentally the same as that in younger adults but does exhibit some different characteristics, particularly with respect to baseline settings and response amplitudes. For example, commonly measured biochemical markers of inflammation, such as interleukin-1beta (IL-1 β), interleukin-6 (IL-6) and Tumor Necrosis Factor Alpha (TNF α) at baseline between acute inflammatory episodes have been shown to be often 2-4-fold higher in the peripheral plasma of aged subjects, and to become increasingly so with advancing age, most steeply in people older than 80 years, when compared to young and middle-aged adults [1,8,9]. In one high quality study of healthy individuals across the adult age range there was mean baseline venous plasma TNF α concentration of 0.6 pg/ml in adults below the age of 30 years compared to a mean of 1.5 pg/ml in those above the age of 70 years [10]. This is one of the manifestations of the clinical state frequently referred to as "inflammaging", a term that is now widely used in clinical geriatric medicine and by researchers focusing on the immunology of ageing [1]. In many individual patients this state can

OPEN ACCESS

ISSN: 2576-5434

be attributed to clinically obvious co-morbid chronic inflammatory diseases such as Rheumatoid Arthritis (RA) or Chronic Obstructive Pulmonary Disease (COPD) [9]. Similar raised plasma levels of the biochemical indicators of inflammation, particularly TNF α and IL- β have been reported in elderly people with other conditions that appear to shift the innate immune web towards a relatively pro-inflammatory setting; common examples are central obesity, atheromatous arterial disease, and type 2 diabetes [11-13]. Further, a sustained low-level rise in plasma pro-inflammatory markers has been found in people with Alzheimer's Disease (AD), and in association with a sedentary lifestyle, chronic renal disease and osteoarthritis [11,14-16]. For reasons that are uncertain, some but not all elderly people have raised baseline plasma inflammatory markers that appear to have no identifiable pathological cause, and age itself appears to be accompanied by an up-regulated baseline inflammatory state [17,18]. Taking C-Reactive Protein (CRP), a non-specific marker of inflammation, as an example, healthy young adult participants were found to have mean venous blood CRP levels of 0.9 mcg/ml compared to 3.0 mcg/ml in healthy people over the age of 65 [17]. Baseline peripheral blood IL-6 and IL-1 β also appear to be significantly correlated with age [19]. These observations have relevance for clinical practice and pathophysiological research because persisting low-amplitude inflammation, as measured by such blood markers, is associated with higher all-cause mortality, lower muscle strength as assessed by handgrip and quadriceps femoris performance, impaired instrumental function, thereby affecting activities of daily living, and lower subjective well being, mood scores and self-reported health status [20,21]. The observed plasma levels of inflammatory markers, such as CRP, IL-1 β , IL-6 and TNF α vary between studies, depending on assay methods, but overall the adverse outcomes considered were broadly associated with 1.5 to 3-fold elevation above those found in healthy age-matched controls. Another factor of importance to clinical geriatric medical practice that is relevant to the argument we are positing in this paper is the persisting pro-inflammatory state that often does not resolve, or does so slowly and incompletely, after an acute inflammatory event such as pneumonia or septicemia. This clearly has implications for the rate and completeness of clinical recovery and the return to pre-event mobility and instrumental function, and therefore offers a clear and clinically important target for research into effective drug interventions [22-29].

It is likely that disease progression in a number of chronic pathological states is to some extent driven by persisting inflammation and not just indicated by the presence of raised inflammatory markers [25]. It appears that a complex interactive relationship exists between cause and effect. An illustrative example is the endothelial inflammation found in atheromatous disease which has been extensively researched and is frequently cited as being typical of the inter-relationship between inflammation, age-related changes and pathological mechanisms [26]. In many elderly people, the inflammatory response to an acute stimulus, such as infection or injury, settles back to baseline more gradually than it does in younger adults. There is a rise in plasma TNF α , IL-1 β and IL-6 last longer and the corresponding rise in the anti-inflammatory cytokine interleukin-10 (IL-10) occurs later and is of lower amplitude. This effect of ageing on innate immune responses has been most clearly shown for pneumococcal and Gram-negative endotoxin antigens [27,28], in which cases an approximately doubling of the time taken to return to pre-stimulus baseline levels in elderly subjects, after

comparable pro-inflammatory peak levels. This observation strongly suggests that regulatory anti-inflammatory function is impaired, with a consequent delay in re-establishing a baseline surveillance state of the innate immune system. It is a plausible argument that this altered mechanism is a likely reason for the slower post-acute functional recovery and persisting low-level inflammation often seen in older patients after acute inflammatory illnesses [2]. The evidence for cytokines having a central role in age-associated inflammation is certainly well established.

Sustained Inflammation and Frailty

Systemic inflammation is closely, and probably causally, associated with some clinically important features of frailty, including sarcopenia, cachexia and low mood [2]. The molecular mechanisms are inevitably complex, incompletely understood and likely to be dependent on the patho-physiological context in individuals. Various components of the innate and adaptive immune systems, notably chemokines, the catecholamine-cortisol axis, complement reactions, interferons, immune cells and other somatic cells communicate with and influence each other in a non-linear manner best pictured as a network or web. A full account is outside the scope of this paper, but the authors have published detailed descriptive reviews elsewhere [2-4]. Effective drug treatments for the subtle dys-immune state described above are therefore likely to be those that cause a corrective multi-component shift in the innate immune network, a process often termed immune modulation, and consequently ameliorate the inflammatory burden that contributes to frailty.

Drugs that Modulate Systemic Inflammation: Particularly Theophylline

Drugs in several pharmacological classes have been shown to dampen systemic inflammation through a range of complex mechanisms that are only partially understood. The list includes, but is not limited to, statins, methyl-xanthines, salicylates, 4AQs, beta-adrenoreceptor blockers, corticosteroids, non-steroidal anti-inflammatory drugs (NSAIDs), metformin and mono-clonal antibodies (MCAs) [2]. Though many of these are licensed for use as anti-inflammatory agents, some are not despite having received considerable attention to explore immune modulating properties and therapeutic potential. For example, theophylline, a methyl-xanthine drug used as a broncho-dilator in patients with asthma and COPD, was noted to improve certain outcomes in patients with COPD, for example improved walking performance, even when there was little or no measurable change in spirometry or arterial blood gas tensions. Further, this occurred at plasma levels well below (<10 mg/L) those likely to cause toxic effects [30]. Follow up studies found that the anti-inflammatory properties of theophylline acted not only locally on airways but also systemically [31,32]. Despite a series of studies, the set of mechanisms through which theophylline modulates innate immune reactions remain incomplete. Further, the patterns and sequences of reactions that constitute the innate immune web and are likely to vary from episode to episode, within and between individuals, and with the prevailing pathophysiological context. Importantly however, theophylline has consistently been found to reduce post-acute TNF α and IL-1 β , and increase IL-10, by around 25-50 per cent. Similarly, in vivo testing of comparable concentrations of pentoxifylline, another methyl-xanthine, caused a progressive fall in the release of pro-inflammatory IL-1 β , IL-6, IL-8 and TNF α of between 20 and 80

percent by peripheral venous blood monocytes over 4 days [32]. This effect appears to be mediated by activation of histone deacetylase-dependent gene switches to set a more anti-inflammatory phenotype [31-33]. These properties are apparently due to theophylline-induced re-direction of the immune web toward an anti-inflammatory state in macrophages and other immune cells. Such cells have been shown to possess several dose-dependent gene switches that up- and down-regulate various cytokines. This is therefore a plausible explanation for the observation that patients with COPD treated with the addition of theophylline to a standard regimen had lower baseline CRP levels and better functional scores compared with control subjects. It also appears that theophylline at normative therapeutic doses reduces inflammation without compromising an appropriate inflammatory response to infection. In a critical care context, it has been shown that a reduction in mortality occurs in patients with severe sepsis treated with theophylline. These effects can be viewed as the clinical manifestations of immune “modulation” [5,7,34-43].

A Possible Role for Chloroquine

Chloroquine and other 4AQs analogues are in established use as anti-inflammatory drugs, mainly in the field of rheumatology for the management of conditions such as rheumatoid arthritis and systemic lupus erythematosus. The anti-inflammatory properties of 4AQs were discovered as a beneficial side effect when chloroquine was in widespread deployment as an anti-malarial drug. Subsequent research established the efficacy of 4AQs in a range of disorders and some progress has been made to identify the mechanisms of action. Early research suggested that the principal anti-inflammatory action was mediated by the changes in macrophage and other monocyte intra-lysosomal pH, and consequent effects on antigen proteolysis and enzymatic activation, as chloroquine and its analogues are concentrated in lysosomes [44]. The accumulation of chloroquine in lysosomes appears to be irreversible and elevates lysosomal pH by trapping of H⁺ ions in association with chloroquine molecules. Hydroxychloroquine, also a lysosomotropic amine, has a similar effect on cellular function. By that apparent mechanism, chloroquine and its analogues interfere with lysosomal pH, which in turn inhibits phagocytosis, antigen processing and proteolysis and, thereby indirectly, chemotaxis, antigen presentation and the phenotypic expression of inflammation. Further research showed that chloroquine and its analogues reduce the release, and probably the production, of the pro-inflammatory cytokines TNF α , IL-1 β and IL-6 by monocytes in vitro, and in vivo during treatment of patients with inflammatory conditions [45]. However, confirmatory in vitro studies have shown that it is likely that the chloroquine-induced reduction of production by monocytes is due to slower conversion of TNF precursors to the active molecule, whereas the reduced production of IL-1 β and IL-6 appears to be caused by a pH-dependent inhibition of cytokine release from cells consequent mainly upon reduced molecular stability but possibly due to acid-base effects on gene transcription [45,46]. These effects appear to be mediated by alterations in antigen processing, as described above, and by an acid-base sensitive reduction in messenger-ribonucleic acid (mRNA) involved in IL-1 β and IL-6 transcription [46], rather than through an epigenetic gene-switching mechanism. It appears, therefore, that the biochemical mechanisms of the immune modulating properties of 4AQs (for example chloroquine) and methyl-xanthines (for example theophylline) are different.

The Need for Research

We have, in previous papers, contended that a case can be made for well conducted studies of the use of low-dose theophylline to modulate persisting inflammation, when it is inappropriately prolonged after stimuli such as sepsis, particularly in pre-frail or overtly frail elderly patients [2,4]. We also advocate similar well designed trials of 4AQs, especially chloroquine and hydroxychloroquine. A role for the use of 4AQs might consequently be established in the reduction of post-acute and chronic inflammation in an attempt to ameliorate the progression from pre-frailty to established frailty, either as a sole treatment or as an adjunct to use of, for example, methyl-xanthines. The principal targets for further research should be specifically: 1) to conduct placebo-controlled trials of anti-inflammatory doses of chloroquine or hydroxychloroquine, given orally, as short-term (4-6-weeks) adjunctive treatment in elderly patients recovering from sepsis or trauma. 2) To establish whether chronic low-grade inflammation, as measured by peripheral blood biochemical markers such as TNF α and IL-1 β can be modified by such treatment, and functional outcomes improved. 3) Include as defined outcomes measures of mortality, mobility scores, functional scores, return to independent living, measurements of muscle strength and other indices of sarcopenia, cognition, mood, wellbeing scores, and duration of hospital treatment.

References

1. Franceschi C, Campisi J. Chronic inflammation (inflammaging) and its potential contribution to age-associated diseases. *J Gerontol. A Biol. Sci. Med. Sci.* 2014; 69: S4-S9.
2. Allen SC. Systemic inflammation in the genesis of frailty and sarcopenia: an overview of the preventive and therapeutic role of exercise and the potential for drug treatments. *Geriatrics.* 2017; 2: 6.
3. Allen SC. Ageing, exercise and the chemistry of inflammation. *Rev Clin Gerontol.* 2015; 25: 73-80.
4. Allen SC, Tiwari D. Theophylline as a systemic anti-inflammatory agent: the need for its revival as a possible adjunctive treatment for “inflammaging”. *BiolEng Med.* 2019; 4: 1-3.
5. Spatafora M, Chiappara G, Merendino AM, D'Amico D, Bellia V, Bonsignore G. Theophylline suppresses the release of TNF alpha by blood monocytes and alveolar macrophages. *Eur Respir J.* 1994; 7: 223-228.
6. Yoshimura T, Usami E, Kurita C, Watanabe S, Nakao T, Kobayashi J, et al. Effect of theophylline on the production of IL-1 beta, TNF alpha and IL-8 by human peripheral blood mononuclear cells. *Biol Pharm Bull.* 1995; 18: 1405-1408.
7. Shih YN, Chen YT, Seethala R, Aisiko I, Frenzl G, Hou P. Effect of the use of theophylline and sepsis outcomes. *Crit Care Med.* 2015; 43: 274.
8. Wener MH, Daum PR, McQuillan GM. The influence of age, sex and race on the upper reference limit of serum C-reactive protein concentration. *J Rheumatol.* 2000; 27: 2351-2359.
9. Chung HY, Cesari M, Anton S, Marzetti E, Giovannini S, Seo AY, et al. Molecular inflammation: Underpinnings of aging and age-related diseases. *Ageing Res Rev.* 2009; 8: 18-30.
10. Bruunsgaard H, Pedersen BK. Age-related inflammatory cytokines and disease. *Immunol Allergy Clin N Am.* 2003; 23: 15-39.
11. Caspersen CJ, Pereira MA, Curran KM. Changes in physical activity patterns in the United States, by sex and cross-sectional age. *Med Sci Sports Exerc.* 2000; 32: 1601-1609.

12. Driver JA, Djousse L, Logroscino G, Gaziano JM, Kurth T. Incidence of cardiovascular disease and cancer in advanced age: Prospective cohort study. *BMJ*. 2008; 337: a2467.
13. Cowie CC, Rust KF, Byrd-Holt DD, Gregg EW, Ford ES, Geiss LS et al. Prevalence of diabetes and impaired fasting glucose in adults in the US population: NHANES survey 1999-2002. *Diabetes Care*. 2006; 29: 1263-1268.
14. Kalaria RN, Maestre GE, Arizaga RN, Friedland RP, Galasko D, Hall K, et al. Alzheimer's disease and vascular dementia in developing countries: Prevalence, management, and risk factors. *Lancet Neurol*. 2008; 7: 812-826.
15. Coresh J, Selvin E, Stevens LA, Manzi J, Kusek JW, Eggers P, et al. Prevalence of chronic kidney disease in the United States. *JAMA*. 2007; 298: 2038-2047.
16. Dagenais S, Garbedian S, Wai EK. Systematic review of the prevalence of radiographic primary hip osteoarthritis. *Clin Orthop Relat Res*. 2009; 467: 623-637.
17. Ballou SP, Lozanski FB, Hodder S, Rzewnicki DL, Mion LC, Sipe JD, et al. Quantitative and qualitative alterations of acute-phase proteins in healthy elderly persons. *Age Ageing*. 1996; 25: 224-230.
18. Ershler WB, Sun WH, Binkley N, Gravenstein S, Volk MJ, Kamoske G, et al. Interleukin-6 and aging: blood levels and mononuclear cell production increase with advancing age and in vitro production is modifiable by dietary restriction. *Lymphokine Cytokine Res*. 1993; 12: 225-230.
19. Wei J, Xu H, Davies, JL, Hemmings GP. Increase in plasma IL-6 concentration with age in healthy subjects. *Life Sci*. 1992; 51: 1953-1956.
20. Kabagambe EK, Judd SE, Howard VJ, Zakai NA, Jenny NS, Hsieh M, et al. Inflammation biomarkers and risk of all-cause mortality in the RCARDS cohort. *Am J Epidemiol*. 2011; 174: 284-292.
21. DeMartini M, Franceschi C, Monti D, Ginaldi L. Inflammation markers predicting frailty and mortality in the elderly. *Exp Mol Pathol*. 2006; 80: 219-227.
22. Jensen GL. Inflammation: Roles in aging and sarcopenia. *J. Parenter. Enter. Nutr*. 2008; 32: 656-659.
23. Penninx BW, Kritchevsky SB, Newman AB, Nicklas BJ, Simonsick EM, Rubin, et al. Inflammatory markers and incident mobility limitation in the elderly. *J Am Geriatr Soc*. 2004; 52: 1105-1113.
24. Christian LM, Glaser R, Porter K, Malarkey WB, Beversdorf D, Kiecolt-Glaser JK. Poorer self-related health is associated with elevated inflammatory markers among older adults. *Psychoneuro endocrinology*. 2011; 36: 1495-1504.
25. Michaud M, Balardy L, Moulis G, Gaudin C, Peyrot C, Vellas B et al. Proinflammatory cytokines, aging, and age-related diseases. *J Am Med Dir Assoc*. 2013; 14: 877-882.
26. Golbidi S, Laher I. Exercise and the aging endothelium. *J Diabetes Res*. 2013.
27. Bruunsgaard H, Skinhoj P, Qvist J, Pedersen BK. Elderly humans show prolonged in vivo inflammatory activity during pneumococcal infections. *J Infect Dis*. 1999; 180: 551-554.
28. Krabbe KS, Bruunsgaard H, Hansen CM, Møller K, Fonsmark L, Qvist J, et al. Ageing is associated with a prolonged fever in human endotoxemia. *Clin Diagn Lab Immunol*. 2001; 8: 333-338.
29. Wu J, Xia S, Kalonis B, Wan W, Sun T. The role of oxidative stress and inflammation in cardiovascular aging. *Biomed Res Int*. 2014.
30. Barnes PJ. Theophylline for COPD. *Thorax* 2006; 61: 742-744.
31. Culpitt SV, de Matos C, Russell RE, Donnelly LE, Rogers DF, Barnes PJ. Effect of theophylline on induced sputum inflammatory indices and neutrophil chemotaxis in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. 2002; 165: 1371-1376.
32. Neuner P, Klosner G, Schauer E, Pourmojib M, Macheiner W, Grünwald C, et al. Pentoxifylline in vivo down-regulates the release of IL-1 beta, IL-6, IL-8 and TNF alpha by human peripheral blood mononuclear cells. *Immunology*. 1994; 83: 262-267.
33. Mascali JJ, Cvetusa P, Negri J, Borish L. Anti-inflammatory effects of theophylline: Modulation of cytokine production. *Ann Allergy Asthma Immunol*. 1996; 77: 34-38.
34. Ito K, Lim S, Caramori G, Cosio B, Chung KF, Adcock IM, et al. A molecular mechanism of the action of theophylline: Induction of histone deacetylase activity to decrease inflammatory gene expression. *Proc Natl Acad Sci USA*. 2002; 99: 8921-8926.
35. Ichiyama T, Hasegawa S, Matsubara T, Hayashi T, Furukawa S. Theophylline inhibits NF-kappa activation and I kappa B alpha degradation in human pulmonary epithelial cells. *Arch Pharmacol*. 2001; 364: 558-561.
36. Scheller J, Chalaris A, Schmidt-Arras D, Rose-John S. The pro- and anti-inflammatory properties of the cytokine interleukin-6. *Biochim Biophys Acta Mol Cell Res*. 2011; 1813: 878-888.
37. So B, Kim H-J, Kim J, Song W. Exercise-induced myokines in health and metabolic diseases. *Integr Med Res*. 2014; 3: 172-179.
38. Vasunilashorn SM, Ngo L, Inouye SK, Libermann TA, Jones RN, Alsop DC, et al. Cytokines and postoperative delirium in older patients undergoing major elective surgery. *J Gerontol Biol Sci Med Sci*. 2015; 70: 1289-1295.
39. Roubenoff R, Parise H, Payette HA, Abad LW, D'Agostini R, Jacques PF, et al. Cytokines, insulin-like growth factor 1, sarcopenia, and mortality in very old community-dwelling men and women: the Framingham Heart Study. *Am J Med*. 2003; 115: 429-435.
40. Stenvinkel P, Ketteler M, Johnson RJ, Lindholm B, Pecoits-Filho R, Riella M, et al. IL-10, IL-6 and TNF-alpha: Central factors in the altered cytokine network of uraemia the good, the bad and the ugly. *Kidney Int*. 2005; 67: 1216-1233.
41. Payette H, Roubenoff R, Jacques PF, Dinarello CA, Wilson WF, Abad LW, et al. Insulin-like growth factor 1 and interleukin 6 predict sarcopenia in very old community-living men and women: the Framingham Heart Study. *J Am Geriatr Soc*. 2003; 51: 1237-1243.
42. Subramanian V, Ragulan AB, Jindal A, Wiswambhar V. The study of tolerability and safety of theophylline given along with formoterol plus budesonide in COPD. *J Clin Diagn Res*. 2015; 9: 10-13.
43. Hancock REW, Nijnik A, Philpott DJ. Modulating immunity as a therapy for bacterial infections. *Nat Rev Microbiol*. 2012; 10: 243-254.
44. Al-Bari MAA. Chloroquine analogues in drug discovery: new directions of uses, mechanisms of actions and toxic manifestations from malaria to multifarious diseases. *J Antimicrob Chemother*. 2015; 70: 1608-1621.
45. Ramos-Avila A, Ventura-Gallegos JL, Zentella-Dehesa A, Machuca-Rodriguez C, Moreno-Altamirano MM, Narvaez V, et al. Immunomodulatory role of chloroquine and pyrimethamine in *Plasmodium yoelii* 17XL infected mice. *Scand J Immunol*. 2007; 65: 54-62.
46. Jang CH, Choi JH, Byun MS, Jue DM. Chloroquine inhibits production of TNF- α , IL-1 β and IL-6 from lipopolysaccharide-stimulated human monocytes/macrophages by different modes. *Rheumatology*. 2006; 45: 703-710.