Original Article

Maxillary Sinus Pathology Detected in Asymptomatic Patients: A Retrospective Cone-beam Computed Tomography Study

Geetika Sobti¹, Jaishree Thanvi², Isha Gaurav³, Nikunj Mathur⁴

^{1,2,3,4} Assistant Professor, Department of Oral Medicine and Radiology, RUHS College of Dental Sciences, Jaipur, Rajasthan, India

ABSTRACT

Introduction: The presence of pathology in maxillary sinus is a frequent finding in cone-beam computed tomography. The aim of this study was to assess the prevalence of pathologic findings in the maxillary sinus by using cone-beam computerized tomography (CBCT).

Methodology: Three hundred CBCT scans were retrospectively inspected for pathologic findings in the maxillary sinus by two observers. All the scans were screened for any variation in size, deviated nasal septum (DNS), opacification, mucosal thickening, retention pseudocyst, polypoidal mucosal thickening, and septa in the maxillary sinus. Statistical data analysis was done using chi-square test. The measure of agreement between the two observers was evaluated by Kappa statistics.

Results: Pathology were found in 63% of the patients with presence of septa (42%) being the most prevalent finding followed by polyposis and deviated nasal septum (39%). No significant association was found between age and pathology while the association of gender and sinus pathology was found to be significant in two parameters i.e. opacification and mucositis where males were predominant.

Conclusion: The interpretation of the complete volume of CBCT images by a dentomaxillofacial radiologist is important not only to avert any complication during surgery but also to identify or diagnose patients who require further referral for their abnormal findings.

INTRODUCTION

Out of the four paranasal sinuses, maxillary sinus is of utmost importance to a dental surgeon. Ascertaining any normal anatomic variation or any coincidental finding in

maxillary sinuses is very crucial, especially prior to surgical procedures such as placement of dental implants and sinus lifting. Moreover, pulpal inflammation/infection can cause sinusitis like symptoms and sinusitis can mimic tooth pain in maxillary posterior region, thus posing diagnostic difficulty.

The panoramic radiograph is a standard two dimension (2D) radiograph used in dental offices to provide an overview of oral and maxillofacial hard tissues, which also include the maxillary sinuses. This view is useful for evaluating the relationship of the maxillary dentition to the sinus and even identify any pathology in them.^{1,2} However, panoramic images commonly present distortions, structural superimpositions, and image enlargement which are inherent to the technique. These factors considerably limit the evaluation and diagnostic accuracy of the maxillary sinus.³

Cone-beam computed tomography (CBCT) was originally developed for angiography in the early 1980s. Since then CBCT technology has been integrated into the field of dentistry for visualization of the maxillofacial region in three-dimensional cone-beam volumetric images. CBCT overcomes all the shortcoming of 2D radiography i.e. geometric distortion and superimposition of surrounding anatomical structures.⁴ It even has several advantages over its comparable counterpart i.e. computed tomography (CT), including reduced radiation dose, higher spatial resolution, lower equipment cost, and potentially a reduced examination expense.⁵

Moreover, CBCT even allows the radiologist to evaluate the entire volume of dental and maxillofacial structures which are frequently neglected when interpreting images of areas at a distance from the dentoalveolar region, such

as the full anatomical aspect of the maxillary sinus. Sometimes such in-depth analysis of CBCT images or maxillary sinus per se may reveal pathologic findings which require further evaluation or have direct significance to the treatment. It is pivotal to be familiar with different anatomic and pathologic findings in sinus to minimize the risk of postoperative complications of maxillary sinus floor lift and other surgeries in this region. As maxillary sinuses are significant anatomic structures in dental practice, their exact and definitive radiological assessment is necessary, and considering CBCT as an important diagnostic image modality in dentistry, the recognition of anatomic variations of the maxillary sinuses in CBCT is noteworthy.⁶ This CBCT study was thus taken up to determine the prevalence of anatomic variations and pathological findings in maxillary sinus in asymptomatic patients visiting the Department of Oral Medicine and Radiology for CBCT.

METHODS

The study sample size comprised of 300 CBCT scans obtained from the archives of Dental Radiology department over a span of 6 months using non-probability (purposive) sampling. As a prescan protocol, the purpose of the CBCT scan was always clearly noted.

Inclusion criteria

- Patients in the age range of 12 to 81 years
- Patients with the entire maxillary sinuses visualized bilaterally in the scan

The patients in the study had been referred for scans for reasons such as implant assessment, evaluation of maxillary or mandibular pathology, temporomandibular joint (TMJ), orthodontic treatment, trauma, calcifications of styloid ligament etc. and none of them had been primarily referred for the scan of the maxillary sinus area because of any sinus pathology.

Exclusion criteria

- Patients under 12 years of age because of their incomplete sinus development. Moreover, certain pathologies such as mucosal thickening and opacification are common findings in early childhood and are not indicative of sinus disease
- CBCT scans that did not have the entire sino-nasal complex in the field of view
- CBCT scans with patient movement
- Scans with fixed orthodontic appliances, implants

or metallic prosthesis in the maxillary posterior region that could generate metallic artifact and thus impaired sinus visualization

All CBCT images were taken using Carestream CS9300 imaging system consisting of an X-ray tube, a source collimator and an amorphous-silicon flat-panel detector (CS 3D Imaging v3.5.7; Carestream Health, Inc New York, USA) with the field of view (FOV) of 17x11 cm or 17x13 cm. Before every scan a scout image was acquired to assess and adjust for proper positioning of the subject to include the region of interest in the scan according to his complaint and requirement. Image volume was reconstructed with isotropic isometric 300x300x300 μm voxels. The tube voltage ranged from 68-90 KVp, tube current was 4 mA, and an exposure time of 11.30 seconds was used. Tube voltage was altered depending on the physique of the patient and presence of metallic structures such as crowns, implants etc, and resulted in change in exposure which ranged from 697-1585 mGy cm².

A written informed consent form was obtained from each patient prior to imaging which included a clause of use of images for academics and research. Two dento-maxillofacial radiologists evaluated the CBCT scans independently in a darkened quiet room on the Dell Precision workstation which comprised of dual monitors (Dell 24-inch nonglossy monitor (1920 $\times$ 1200 resolution) using CS 3D software (v3.5.7). Each viewing session lasted 30 minutes and care was taken to ensure that the next viewing session was not before a period of 24 hours for the examiners. For intra examiner reliability and reproducibility of the measurements, the images were evaluated for the second time by the same observers two weeks later. Observers were allowed to zoom and adjust brightness and contrast of the images using the image processing tool in the software to ensure optimal visualization. Any divergences or conflicting views between the examiners were resolved by discussion and reaching a consensus.

The parameters which were evaluated in the present study were developed on the basis of the published literature and indicated the presence of an incidental finding or abnormality in maxillary sinus.⁷⁻¹¹ It included evaluation of:

- (i) Size i.e. any increase or decrease in dimension of the sinus, in accordance to the average dimension (34 $\times$ 33 $\times$ 25 mm) (Figure 1a, 1b).

Figure 1: Variation in the size of maxillary sims. Coronal section shown (a) bilateral hypoplasia (b) unilateral right maxillary sinus hypoplasia.

Figure 2: Coronal view showing deviated nasal septum towards right.

Figure 3: Coronal section shows (a) partial opacification in both the sinuses (b) complete opacification of right maxillary sinus.

Figure 4: Sagittal view showing mucosal thickening on the floor of sinus.

Figure 5: Coronal view showing dome shaped pseudocyst attached partly on the floor and lateral wall of right maxillary sinus.

Figure 6: Coronal view showing bilateral polypoid alongwith increased thickness of sinus lining on its all walls.

Figure 7: Axial section showing variations in septae (a) incomplete bilateral septae (b) complete bilateral septae.

- (ii) Deviated nasal septum (DNS) (Figure 2).
- (iii) Any opacity in the sinus cavity: the designation of partial opacification was given when at least 1/3rd of the maxillary sinus was opacified and the term complete opacification was used when the complete maxillary sinus was opacified in all axial and coronal slices either unilaterally or bilaterally (Figure 3a, 3b).
- (iv) Mucosal thickening: Pathology was considered when thickening of >2 mm was present in at least one wall of the sinus; Coronal and sagittal views were used for evaluation and the thickest area was recorded (Figure 4).
- (v) Retention pseudocyst: it appears as non-corticated, smooth, dome-shaped radiopaque mass (Figure 5).
- (vi) Polyp and polypoidal mucosal thickening: Polyp also appear as homogenous round opacity similar to retention pseudocyst but can be differentiated with the fact that polyp is accompanied with a thickened mucous membrane lining as it is no more than an accentuation of the sinus mucosa¹² (Figure 6).
- (vii) Septa: Axial images were used for identification of

presence of septa. They were either incomplete or complete dividing the sinus into locules (Figure 7a, 7b).

For statistical analysis, patients were separated into seven age groups: 12-21, 22-31, 32-41, 42-51, 52-61, 62-71, and 72-81 years. Statistical analysis was done using SPSS (version 19) and the test applied was chi-square.

RESULTS

The age of the participants in the study ranged from 12-81 years with 157 males and 140 females. Out of the seven parameters which were evaluated in the study, 63% of the participants manifested one or more positive findings with the distribution of 104 males and 85 females (Table 1).

The age group with most prevalent finding was 12-21(23%) followed by 22-31 and 42-51(17.5%) which had equal percentage of findings. Presence of septa (42%) was the most prevalent incidental finding followed by polyposis and DNS (39%) and mucositis (20.6%). The prevalence of sinus hypoplasia and pseudocyst was 7% and 4% respectively (Table 2).

The prevalence of unilateral and bilateral septae was found to be equal i.e. 40 in each; however, variation in gender distribution was seen wherein unilateral septa

Table 1: Gender wise distribution of incidental findings in maxillary sinus of study participants (n=189)

Gender	Variation in size	DNS	Partial opacification	Complete opacification	Mucositis	Retention pseudocyst	Polypoidal mucosal thickening	Septa
Males	9	47	14	16	18	8	44	43
Females	5	28	18	4	21	1	32	37

DNS: deviated nasal septum

Table 2: Age-wise distribution of incidental findings in maxillary sinus of study participants (n=189)

Age groups	Variation in size	DNS	Partial opacification	Complete opacification	Mucositis	Retention pseudocyst	Polypoidal mucosal thickening	Septa
12-21	3	15	5	3	7	3	25	22
22-31	1	13	5	3	8	2	14	16
32-41	1	9	3	0	3	2	5	9
42-51	4	11	4	3	9	1	10	15
52-61	1	12	7	4	3	0	10	6
62-71	1	9	7	6	9	0	10	10
72-81	3	6	1	1	0	1	2	2
Total	14	75	32	20	39	09	76	80
%	7.4	39	16.9	10.5	76	4.7	40	42

DNS: Deviated nasal septum

were more common in females (55%) while bilateral septa were more in males (62.5%). Out of the 49 cases of mucositis, 14 could be attributed to odontogenic cause i.e. it was referred to as odontogenic mucositis. Another inference which could be made from the data was that out of the 75 cases of DNS, 59 (78.6%) were associated with some kind of inflammation of sinus i.e. either mucositis, polyposis or opacification and it was found to be statistically significant ($p < 0.05$).

No significant association was found between age groups and sinus pathology. Although the prevalence of partial and complete opacification and mucositis were more common in patients above 50 years of age but it was not statistically significant. Regarding the association between gender and sinus pathology, it was found to be significant ($p < 0.05$) in two parameters i.e. opacification and mucositis where males were predominant. The inter-observer agreement between the observers was assessed by Kappa statistics and it was found to be 0.72.

DISCUSSION

The prevalence of pathologic findings in one or both the maxillary sinuses was found to be 63% in the present study. This data varies among the previous studies and was found to be in the range of 27% to 61%.^{8,13-15} A systematic review by Vogiatzi et al reported that frequency of maxillary sinus disease varied widely from 14.3% to 82%.⁵ The reasons which can be attributed for this discord include the sampling and the diagnostic criteria applied to evaluate the pathosis in maxillary sinuses.¹⁶ A low prevalence was noticed when the authors selected only a particular age group like patients undergoing orthodontic treatment or patients scheduled for sinus lift procedures.¹⁵ While higher prevalence was observed when a wider age group and numerous parameters were evaluated simultaneously as in the study by Raghav et al¹⁷ and our own study where we investigated seven possible causes of alterations in maxillary sinuses. Another reason for this observation can be the application of varied definitions or lack of any clear cut criteria for certain pathology such as mucosal thickening. There is no consensus in the literature for the cut off value of mucosal thickening and the measurements range from 2-6 mm.^{8,16} In the present study, the diagnostic criteria were pre defined and the observers were calibrated to strengthen the reproducibility of the results.

There was a higher frequency of incidental findings in

males which is in agreement with the literature but the difference was not statistically significant in the study.^{8,10, 18}

It has been proposed that males are more vulnerable to the occurrence of mucosal thickening due to irritation of the sinus mucosa because of the greater prevalence of dental pathology in them.¹⁸ Although there has been a study which showed no correlation between the findings in maxillary sinus and gender.⁹

In the present study, presence of septa was the most prevalent incidental finding which is in accordance with the studies by Tadinada A et al¹⁹ and Lana JP et al.²⁰ According to a systematic review by Maestre-Ferrín L et al²¹, the septal prevalence ranged from 13-56%. Its prime distribution in the younger age group of 12-21 years in the study implies that they are mainly primary septa. According to Krennmer et al²², primary septa are the ones which arise during the development of maxilla and secondary septa arise from irregular pneumatization of the sinus floor following tooth loss. The present study emphasizes the significance of not ignoring this crucial finding as it increases the chance of membrane perforation leading to profuse bleeding during implant surgery. Moreover, septa can also obstruct proper visualization of the sinus and its accessibility during sinus floor augmentation.²³

Deviated nasal septum, polypoidal mucosal thickening, and mucositis were the next most common findings in the present study. The radiographic presentation of thickening of the mucosal lining of maxillary sinus can be the outcome of many causes like bacterial, viral or fungal infection, allergy, polyps, DNS or odontogenic. Nasal polyps and anatomical variation such as DNS can block the middle meatus and impede the drainage of the sinuses resulting in mucositis.²⁴

Odontogenic mucositis was 28.5% with floor of the sinus being the most affected location and maxillary first molar being the most common tooth involved. Its prevalence in literature ranges from 10-40%. Sinus inflammation and infection from a dental source is more common in pneumatized sinus in which the pathogens can easily penetrate through the thin Schneiderian membrane from any periapical or periodontal infection or perforation of sinus iatrogenically during extractions, implant surgery etc. The accurate identification of odontogenic sinusitis is crucial because directing the medical and surgical therapies only to treat sinus per se without addressing the

dental pathology will result in failure of the treatment and persistence of symptoms.^{1,7}

No statistical significance of dental disease and polyposis was found in the study and thus the reason for polypoidal mucosal thickening can be partially attributed to DNS.

Complete opacification accounted for only 10% of the total findings and this data was in agreement with the previous literature.²⁵ Although chronic rhinosinusitis was the most common diagnosis, other differentials should also be ruled out for this condition such as fungal disease and malignancy in older individuals and antrochoanal polyp in younger patients especially with unilateral opacification.²⁶ Patients with a history of mechanical or barotrauma can also present with complete opacification due to hemorrhage as blood and pus appear radiographically identical.¹⁷

The limitation of the study was that it was a retrospective prevalence study lacking in depth clinical history and follow up of the patients. Etiology of the pathologic findings or association among the different parameters can only be presumed but cannot be established with confirmation.

CONCLUSION

In view of the high prevalence rate of maxillary sinus findings or pathology in the CBCT scans in asymptomatic patients, it is the responsibility of every dento maxillo-facial radiologist to be vigilant and comprehensively review the entire volume. This will not only help them to evade complications during implant surgery but also lead to diagnosis of certain conditions like pansinusitis, blocked osteomeatal complex, concha bullosa etc. which necessitate appropriate referrals and are beyond the scope of the dentist.

REFERENCES

1. Simuntis R, Kubilius R, Vaitkus S. Odontogenic maxillary sinusitis: A review. *Stomatologija* 2014;16(2): 39-43.
2. Allareddy V, Vincent SD, Hellstein JW, Qian F, Smoker WR, Ruprecht A. Incidental findings on cone beam computed tomography images. *Int J Dent* 2012;9:1-9.
3. Danda TF, Santos KC, Oliveira LB, Shintaku W, Oliveira JX. Comparison of CBCT and panoramic radiography for the evaluation of opacification of the maxillary sinuses. *Clin Lab Res Den* 2014;20(1):23-30.
4. Rheem S, Nielsen IL, Oberoi S. Incidental findings in the maxillofacial region identified on cone-beam computed tomography scans. *J Orthod Res* 2013;1(1):33-39.
5. Vogiatzi T, Kloukos D, Scarfe WC, Bornstein MM. Incidence of anatomical variations and disease of the maxillary sinuses as identified by cone beam computed tomography: A systematic review. *Int J Oral Maxillofac Implants* 2014;29(6):1301-14.
6. Shahidi S, Zamiri B, Momeni Danaei SH, Salehi S, Hamedani SH. Evaluation of anatomic variations in maxillary sinus with the aid of cone beam computed tomography (CBCT) in a population in south of Iran *J Dent (Shiraz)* 2016;17(1):7-15.
7. Maillet M, Bowles WR, McClanahan SL, John MT, Ahmad M. Cone-beam computed tomography evaluation of maxillary sinusitis. *J Endod* 2011;37(6):753-57.
8. Rege IC, Sousa TO, Leles CR, Mendonça EF. Occurrence of maxillary sinus abnormalities detected by cone beam CT in asymptomatic patients. *BMC Oral Health* 2012;12:30.
9. Dobelev I, Kise L, Apse P, Kragis G, Bigestans A. Radiographic assessment of findings in the maxillary sinus using cone-beam computed tomography. *Stomatologija* 2013;15(4):119-22.
10. Gracco A, Parenti SI, Ioele C, Bonetti GA, Stellini E. Prevalence of incidental maxillary sinus findings in Italian orthodontic patients: A retrospective cone-beam computed tomography study. *Korean J Orthod* 2012;42(6):329-34.
11. Jafari-Pozve N, Sheikhi M, Ataie-Khorasgani M, Jafari-Pozve S. Aplasia and hypoplasia of the maxillary sinus: a case series. *Dent Res J* 2014;11(5):615-17.
12. White SC, Pharoah MJ. Oral radiology principles and interpretation. 6th ed. St. Louis: Mosby;2009.
13. Ritter L, Lutz J, Neugebauer J, Scheer M, Dreiseidler T, Zinser MJ, et al. Prevalence of pathologic findings in the maxillary sinus in cone-beam computerized tomography. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2011;111(5):634-40.
14. Pazera P, Bornstein MM, Pazera A, Sendi P, Katsaros C. Incidental maxillary sinus findings in orthodontic patients: A radiographic analysis using cone-beam computed tomography (CBCT). *Orthod Craniofac Res* 2011;14:17-24.
15. Beaumont C, Zafiroopoulos GG, Rohmann K, Tatakis DN. Prevalence of maxillary sinus disease and abnormalities in patients scheduled for sinus lift procedures. *J Periodontol* 2005;76(3):461-67.
16. Bozdemir E, Görmez O, Yıldırım D, Erik AA. Paranasal sinus pathoses on cone beam computed tomography. *J*

- Istanbul Univ Fac Dent 2016;50(1): 27-34.
17. Raghav M, Karjodkar FR, Sontakke S, Sansare K. Prevalence of incidental maxillary sinus pathologies in dental patients on cone-beam computed tomographic images. *Contemp Clin Dent* 2014; 5(3):361-65.
 18. Vallo J, Suominen-Taipale L, Huuonen S, Soikkonen K, Norblad A. Prevalence of mucosal abnormalities of the maxillary sinus and their relationship to dental disease in panoramic radiography: Results from the health 2000 health examination survey. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod* 2010;109(3):e80-e87.
 19. Tadinada A, Jalali E, Al-Salman W, Jambhekar S, Katechia B, Almas K. Prevalence of bony septa, antral pathology, and dimensions of the maxillary sinus from a sinus augmentation perspective: A retrospective cone-beam computed tomography study. *Imaging Sci Dent* 2016;46(2):109-15.
 20. Lana JP, Carneiro PM, Machado Vde C, de Souza PE, Manzi FR, Horta MC. Anatomic variations and lesions of the maxillary sinus detected in cone beam computed tomography for dental implants. *Clin Oral Implants Res* 2012; 23(12): 1398-403.
 21. Maestre-Ferrín L, Galán-Gil S, Rubio-Serrano M, Peñarrocha-Diogo M, Peñarrocha-Oltra D. Maxillary sinus septa: a systematic review. *Med Oral Patol Oral Cir Bucal* 2010;15(2):e383-e386.
 22. Krennmair G, Ulm CW, Lugmayr H, Solar P. The incidence, location, and height of maxillary sinus septa in the edentulous and dentate maxilla. *J Oral Maxillofac Surg* 1999; 57: 667-71.
 23. Gandhi KR, Wabale RN, Siddiqui AU, Farooqui MS. The incidence and morphology of maxillary sinus septa in dentate and edentulous maxillae: a cadaveric study with a brief review of the literature. *J Korean Assoc Oral Maxillofac Surg* 2015;41(1):30-36.
 24. Bell GW, Joshi BB, Macleod RI. Maxillary sinus disease: Diagnosis and treatment. *Br Dent J* 2011; 210(3):113-18.
 25. Lim CG, Spanger M. Incidental maxillary sinus findings in patients referred for head and neck CT angiography. *Singapore Dent J* 2012;33(1):1-4.
 26. Chen HJ, Chen HS, Chang YL, Huang YC. Complete unilateral maxillary sinus opacity in computed tomography. *J Formos Med Assoc* 2010;109(10):709-15.

Corresponding Author

Dr Geetika Sobti, 53, Vishnupuri, opposite Dak Colony, Malviya Nagar, Jaipur-302017, Rajasthan, India.
email: geetikamohitmidha@gmail.com