BIBLIOGRAPHIC INFORMATION SYSTEM

Journal Full Title: Journal of Biomedical Research & Environmental Sciences

Journal NLM Abbreviation: J Biomed Res Environ Sci

Journal Website Link: <https://www.jelsciences.com>

Journal ISSN: 2766-2276

Category: Multidisciplinary

Subject Areas: Medicine Group, Biology Group, General, Environmental Sciences

Topics Summation: 128

Issue Regularity: Monthly

Review Process: Double Blind

Time to Publication: 21 Days

Indexing catalog: [Visit here](#)

Publication fee catalog: [Visit here](#)

DOI: 10.37871 ([CrossRef](#))

Plagiarism detection software: iThenticate

Managing entity: USA

Language: English

Research work collecting capability: Worldwide

Organized by: [SciRes Literature LLC](#)

License: Open Access by Journal of Biomedical Research & Environmental Sciences is licensed under a Creative Commons Attribution 4.0 International License. Based on a work at SciRes Literature LLC.

Manuscript should be submitted in Word Document (.doc or .docx) through

Online Submission

form or can be mailed to support@jelsciences.com

**IndexCopernicus
ICV 2020:
53.77**

Vision: Journal of Biomedical Research & Environmental Sciences main aim is to enhance the importance of science and technology to the scientific community and also to provide an equal opportunity to seek and share ideas to all our researchers and scientists without any barriers to develop their career and helping in their development of discovering the world.

Risk Factors Associated with Mortality among Chronic Kidney Disease Patients on Regular Hemodialysis Presenting in Emergency Services

Tirtha Man Shrestha^{1*}, Laxman Bhusal¹, Ram Prasad Neupane¹, Rajan Ghimire² and Pratap Narayan Prasad³

¹Department of General Practice and Emergency Medicine, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal

²Clinical Coordinator, District Hospital, Terhathum, Province 1, Nepal

³Campus chief, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal

ABSTRACT

Background: With the increasing number of Chronic Kidney Disease (CKD), emergency visit of these patients is also increasing. This study tried to find some of the reasons for which patients with CKD visit the emergency room and the reasons for their mortality.

Method: A cross-sectional study was done in the emergency room of Tribhuvan University Teaching Hospital, Kathmandu, Nepal. We conducted this study from 1 May 2018 to 31 October 2018 among the adult CKD patients under regular hemodialysis. We used a convenience sampling method. Three hundred patients were included. We studied the following variables: patient's age, sex, risk factors, laboratory parameter during the emergency visit (viz. hemoglobin, pH, serum bicarbonate level, and potassium level), emergency hemodialysis, blood transfusion, and clinical outcome during emergency room stay.

Result: We enrolled 300 patients in the study. The mean age was 45.04 years in the mortality group and 45.69 years in the survival group. 152 (50.7%) of patients had hypertension. Mean hemoglobin was 6.52 gm% (SD = 1.93). Mean hemoglobin in survivor and the non-survivor group was 6.59 gm% and 5.58 gm% respectively. Serum creatinine was 1220.87 micromol/l and 1064.01 micromol/l in mortality and survivor group respectively. Likewise, serum potassium was 6.13 mEq/l and 5.74 mEq/l among mortality and survivor groups respectively. Binary logistic regression showed significant association ($p < 0.05$) of anemia, emergency dialysis and presence of sepsis with the mortality. There was significant correlation of presence of comorbidities, anemia, serum creatinine, serum potassium level, and sepsis with mortality. Area under the Receiver Operating Curve to predict mortality among CKD patients was 0.660 for potassium and 0.598 for serum creatinine.

Conclusion: Anemia, increased serum creatinine, and hyperkalemia was significantly correlated with mortality in chronic kidney disease and were causes of frequent visits in the emergency room. Therefore, we should address these factors during the management of CKD patients.

INTRODUCTION

In Nepal, the cases of chronic kidney disease are increasing. In the same ratio complications related to CKD and hemodialysis are also increasing. The management of this disease is expensive. 10.6% of Nepalese had CKD during a community screen [1]. In contrast, 56.02% of outpatient and in-patient cases of a tertiary hospital in Nepal had chronic kidney disease [2]. Approximately 13.1% of United States adults had CKD and it led to ESRD, cardiovascular disease, and

*Corresponding author

Tirtha Man Shrestha, Department of General Practice and Emergency Medicine, Maharajgunj Medical Campus, Institute of Medicine, Tribhuvan University, Kathmandu, Nepal

Tel: +977-985-101-7345

E-mail: tirtha_48@yahoo.com.au

DOI: 10.37871/jbres1178

Submitted: 25 December 2020

Accepted: 04 January 2021

Published: 05 January 2021

Copyright: © 2021 Shrestha TM, et al. Distributed under Creative Commons CC-BY 4.0

OPEN ACCESS

Subject: Medicine Group

Topic & Subtopic(s): Emergency Medicine, Anesthesiology, Nephrology

Keywords

- Chronic kidney disease
- Emergency presentation
- Mortality
- Risk factors

premature death [3]. The main reasons for frequent visits of CKD patients in emergency services were high blood pressure, anemia, high potassium, increased metabolic acidosis, pulmonary edema, hypoxia, hypocapnia, cardiac failure, and other cardiovascular problem [4]. Diabetes, hypertension, glomerulonephritis and other disorder are major causes of CKD [5-7].

Early detection and treatment can often prevent CKD from getting worse. The risk of death from cardiovascular disease is higher than requiring dialysis among CKD patients [3]. A study was done in Haryana, India in 2015 which showed chronic glomerulonephritis as the commonest presentation (34.72%) followed by diabetic nephropathy (23.44%) as a cause of CKD [5]. Similar study in Nepal showed the presence of overweight, obesity, hypertension, diabetes, and proteinuria were 20%, 5.0%, 38.6%, 7.5%, and 5.1% respectively [8]. Cancer (31.9%) and cardiovascular disease (30.2%) were the two most common causes of mortality [9]. In patients with CKD having comorbidities like cardiovascular problems and malignancy, hyponatremia was associated with all-cause mortality. Hyponatremia was associated with all-cause and non-cardiovascular/non-malignancy related deaths [10]. Though metabolic acidosis is common among hemodialysis patients, respiratory acid-base disorders are also important to consider. Therefore, complete acid-base measurements should be done for a complete analysis of acid-base disorder in CKD patients [11]. Patients with CKD have an increased risk of atrial fibrillation and stroke [12].

There are few centers for the diagnosis and management of kidney diseases in Nepal. CKD is one of the common causes of the Emergency visit. Therefore, the emergency physician should update their knowledge on this matter. Despite the maintenance hemodialysis, frequent emergency visits of patients point towards either inadequate hemodialysis or inadequate management of other medical conditions like hypertension, diabetes, anemia, hyperkalemia, etc. Therefore, this study aims to analyze the factors associated with the mortality of chronic kidney disease cases under regular maintenance hemodialysis during an emergency visit. This will help to improve the management of patients and decrease the emergency room visit of CKD patients.

METHODOLOGY

This was a prospective cross-sectional study. It was conducted in the emergency room of Tribhuvan University Teaching Hospital, Maharajgunj; Kathmandu; Nepal from 1 May 2018 to 31 October 2018.

Sample size

The sample size was calculated using the Daniel method [8]. Sample size was 300. We collected samples without randomization by purposive non-probability sampling method. The inclusion criteria for the study were: (a)

Patients of age > 16years. (b) Patients with chronic kidney disease under maintenance hemodialysis. (c) Patients visiting the emergency room. Pregnant women and patients with 'do not resuscitate' order were excluded from the study.

Data analysis

We studied variables in the following sectors:

Demographic data: Age, sex.

Comorbid health conditions: Diabetes, hypertension, glomerulonephritis, heart failure, liver failure, malignancy, chronic pulmonary disease, management in an emergency: emergency dialysis and blood transfusions.

Laboratory values: Hemoglobin, serum potassium, serum creatinine, serum bicarbonate.

Clinical outcome: Improved or died.

We collected data from the clinical record file of the patient after getting proper consent. Patients were followed until their stay in the emergency room. Clinical outcomes (survived and mortality) were studied during emergency stay only. These study variables were analyzed to search for any association with mortality among the patients using Pearson's Correlation. We did binary logistic regression to find the association of confounding factors. Receiver Operating Curve (ROC) analysis and area under the ROC were used to calculate the efficacy of various laboratory tests to predict mortality in CKD patients [13]. We used IBM SPSS 25.0 for all calculations. Institutional Review Board, Institute of Medicine, Tribhuvan University; Kathmandu; Nepal provided the ethical clearance.

RESULTS

During the study period from 1 May 2018 to 31 October 2018, 300 patients met the criteria to participate in the study. Among them 190 (63.30%) were male and 110 (36.70%) were female. Among these patients, 152 (50.7%) had hypertension, 63 (21%) had diabetes, 63 (21%) had glomerulonephritis, and the remaining seven (2.33%) had other comorbidities like nephrolithiasis, benign enlargement of the prostate, renal tuberculosis, and polycystic kidney disease. 269 (89.7%) of these patients underwent emergency hemodialysis and 239 (79.7%) required blood transfusion, 20 (6.7%) patients had sepsis and 23 (7.7%) participants died during the emergency stay.

The mean age of all patients was 45.64 years (SD = 17.15). Mean hemoglobin was 6.52 gm% (SD = 1.93), pH was 7.17 (SD = 0.154), serum creatinine was 1076.3 micromol/l (SD = 367.25), serum potassium was 5.77 mEq/l (SD = 0.76), pCO₂ was 22.48 (SD = 6.82), and serum bicarbonate was 10.81 mmol/L (SD = 4.36). Data for pH and serum potassium were negatively skewed (-0.192 and -0.271 respectively).

Out of 23 (7.7%) mortality, 16 (5.33%) were male and 7 (2.33%) were female. Among the non-survivors, 19 (6.33%) had other comorbidities like hypertension, diabetes, and glomerulonephritis. 18 (6.0%) patients who had sepsis died and 2 (0.67%) survived (Table 1).

The mean age in the survivor and non-survivor groups were 45.04 (SD = 18.96) and 45.96 (SD = 17.03) respectively. Patients who could not survive had lower hemoglobin (5.58 vs 6.59), pH (7.11 vs 7.18), and serum bicarbonate (9.89 vs 10.89) than those who survived. Meanwhile, those who could not survive had higher pCO₂ (24.04 vs 22.35), serum creatinine (1220 vs 1064), serum potassium (6.13 vs 5.74) than those who survived (Table 2).

Binary logistic regression analysis showed significant association of low hemoglobin (OR = 2.679, $p = 0.041$), emergency dialysis (OR = 0.016, $p = 0.015$) and presence of sepsis (OR = 7649.521, $p < 0.001$) to increase mortality rate in chronic kidney disease (Table 3). However, there was significant correlation of mortality with low hemoglobin (Pearson's correlation = 0.141, $p = 0.015$); high serum creatinine (Pearson's correlation = -0.114, $p = 0.049$); high serum potassium (Spearman's correlation = -0.148, $p = 0.01$); presence of sepsis (Pearson's correlation = 0.827, $p < 0.001$). The value of serum potassium was negatively skewed so Spearman's correlation was used.

Receiver Operating Curve (ROC) and the Area Under the Receiver Operating Curve (AUROC) for serum potassium and

creatinine to predict mortality in CKD patients is shown in figure 1 and table 4. AUROC for both serum creatinine and potassium is < 0.7 which means these tests are poor to predict mortality.

Youden index was maximum (0.33) at serum potassium 6.25 mEq/L. So, using serum potassium level of 6.25 mEq/L as a cutoff to predict mortality among CKD patients; sensitivity was 57% (positive likelihood ratio = 2.37) and specificity was 76% (negative likelihood ratio = 0.57). Similarly, the Youden index was maximum (0.30) at serum creatinine level 1195 micromol/l with a sensitivity of 61% (1.96) and specificity of 69% (0.57) (Table 5). Sensitivity decreases and specificity increases for an increase in the value of serum potassium and creatinine.

DISCUSSION

With the advancement in modern medicine, the incidence of communicable diseases is decreasing. Non-communicable diseases like hypertension and diabetes are rising as the prominent health problem [14]. With this, there is an increase in the incidence of CKD also [15]. The mean age of patients was similar to the study done by Sigdel MR and et al in the same center [16]. The mean age in the mortality group was 45.04 years (SD = 18.96) and in the survivor group, it was 45.69 years (SD = 17.03). Similar results were found in studies from Saudi Arabia [17], Africa [18], India [19,20]. In a developed nation, the average age for CKD was 60 years and above [21].

Table 1: Comparison of categorical variables among survivor and non-survivor group.

		Mortality			
		Yes (n = 23)		No (n = 277)	
		Number	%	Number	%
Sex	Male	16	5.33%	174	58.00%
	Female	7	2.33%	103	34.33%
Comorbidities	No	4	1.33%	11	3.67%
	Yes	19	6.33%	266	88.67%
Hypertension	Yes	11	3.67%	141	47.00%
	No	12	4.00%	136	45.33%
Diabetes	Yes	3	1.00%	60	20.00%
	No	20	6.67%	217	72.33%
Glomerulonephritis	Yes	5	1.67%	58	19.33%
	No	18	6.00%	219	73.00%
other comorbidities	Yes	0	0.00%	7	2.33%
	No	23	7.67%	270	90.00%
Emergency hemodialysis	Yes	20	6.67%	249	83.00%
	No	3	1.00%	28	9.33%
Blood Transfusion	Yes	20	6.67%	219	73.00%
	No	3	1%	58	19.33%
Sepsis	Yes	18	6.00%	2	0.67%
	No	5	1.67%	275	91.67%

Table 2: Comparisons of discrete variables using measures of central tendencies among survivors and non-survivor groups.

	Mortality			
	Yes (n = 23)		No (n = 277)	
	Mean; S.D	Range	Mean; S.D	Range
Age in years	45.04; 18.96	62	45.69; 17.03	69
Hemoglobin (gm%)	5.58; 2.02	8	6.59; 1.9	10
pH	7.11; 0.21	0.82	7.18; 0.15	1.1
pCO2(mmHg)	24.04; 13.72	62	22.35; 5.94	39
Serum Bicarbonate (mmol/L)	9.89; 5.06	20.6	10.89; 4.30	23
Serum Creatinine (micromol/l)	1220.87; 481.58	1860	1064.01; 354.59	2210
Serum Potassium (mEq/L)	6.13; 0.89	4.2	5.74; 0.78	6.2

Table 3: Binary logistic regression and correlation analysis of variables in association with mortality in chronic kidney disease.

	p-value	Odds Ratio	95% C.I. for Odds Ratio			
			Lower	Upper	Pearson's correlation	p-value
Age in years	0.423	1.028	0.96	1.101	0.01	0.862
Sex	0.204	0.234	0.025	2.201	0.037	0.52
Hypertension	0.999	5093786.484	0		-0.016	0.778
Diabetes	0.999	5951714.182	0		-0.056	0.331
Glomerulonephritis	0.999	598326.416	0		0.005	0.928
Hemoglobin (gm%)	0.041	2.679	1.039	6.907	0.141	0.015
pH	0.415	92.508	0.002	4886427.644	0.111*	0.054
pCO2	0.747	1.025	0.883	1.19	-0.066	0.256
HCO3	0.342	0.813	0.531	1.246	0.061	0.295
Serum Creatinine	0.293	0.999	0.996	1.001	-0.114	0.049
Serum Potassium	0.144	0.456	0.159	1.307	-0.148*	0.01
Dialysis	0.015	0.016	0.001	0.45	-0.026	0.658
Blood Transfusion	0.291	0.137	0.003	5.487	0.052	0.368
Sepsis	<0.001	7649.521	137.294	426203.855	0.827	<0.001

*Spearman's correlation is used as the data are negatively skewed. Bold values indicate statistically significant values with a $p < 0.05$.

Table 4: Prediction of mortality among CKD patients using Area under the receiver operating curve.

Variables	Area	p-value	95% Confidence Interval	
			Lower Bound	Upper Bound
Serum potassium	0.66	0.011	0.53	0.791
Serum creatinine	0.598	0.119	0.457	0.739

In our study, Hypertension (50.7%) was most common followed by diabetes (21%) and glomerulonephritis (21%). In another study, CKD was seen in newly diagnosed hypertensive patients (51.9%) and normotensive controls (23.6%) which were statistically significant ($p < 0.001$) [22]. Chronic glomerulonephritis (42.3%) and diabetic nephropathy (21.1%) were the leading causes of CKD. Nephrotic syndrome accounted for 22.36% [2].

We found that the mean serum bicarbonate level was 9.89 mmol/L (SD = 5.06) in non-survivors and 10.89 mmol/L (SD = 4.30) in the survivor group. Patients with low eGFR had lower serum bicarbonate levels (<22 mmol/L) which

are associated with adverse renal outcome and mortality [10]. In our study, mean serum potassium in the mortality group was 6.13 (SD = 0.89, range = 4.2) and in survivor, it was 5.74 (SD = 0.78, range = 6.20). The risk of hyperkalemia is increased with CKD, and its occurrence increases the odds of mortality and need of dialysis [4,23-25]. Hypokalemia and hyperkalemia both increase the risk of mortality in CKD patients while hypokalemia is associated with faster CKD progression [25].

In our study, 89.7% of patients underwent emergency dialysis. A south Indian study showed that nearly 87.4% of patients with ESRD required emergency dialysis [26].

Figure 1 ROC curve to test the efficacy of serum Potassium and Creatinine to predict mortality in CKD patients under maintenance hemodialysis.

Table 5: Sensitivity, specificity and Youden index of serum potassium and creatinine to predict mortality in chronic kidney disease.

Potassium (mEq/L)	Sensitivity	Specificity	Youden Index	Creatinine micromol/l	Sensitivity	Specificity	Youden Index
5.95	0.65	0.67	0.32	1121	0.61	0.67	0.28
6.05	0.61	0.68	0.29	1185	0.61	0.69	0.29
6.25	0.57	0.76	0.33	1195	0.61	0.69	0.3
6.35	0.52	0.79	0.32	1217	0.57	0.71	0.27
6.45	0.44	0.85	0.29	1228.5	0.48	0.72	0.2

Scheduled dialysis decreased mortality (3% vs 17%, $P = 0.01$; absolute risk reduction, 14%) in comparison to emergency-only dialysis [27]. In comparison to patients without hemodialysis, patients under maintenance hemodialysis had higher ED utilization and a significantly higher risk of resuscitation during ED visits [27].

Our study showed that overall mean hemoglobin was 6.52 gm% (SD = 1.93). Mean hemoglobin among the non-survivor group was 5.58 gm% (SD = 2.02) and in the survivor group, it was 6.59 gm% (SD = 1.9). The study done by Sigdel MR showed that mean hemoglobin was 9.98 gm% in patients with CKD stage 3 and 8.23 gm% in CKD stage 5 [16]. 79.7% of our study population received blood transfusions. In another study, 20% of non-dialysis patients received at least one transfusion (mean of two units, range = 1-4). The mean hemoglobin before transfusion among these patients was 8.8 g/dl (SD = 1.5) [28].

We found association of low hemoglobin (OR = 2.679, $p = 0.041$), emergency dialysis (OR = 0.016, $p = 0.015$) and presence of sepsis (OR = 7649.521, $p < 0.001$) to increase mortality rate in chronic kidney disease. There was significant correlation of mortality with low hemoglobin (Pearson's correlation = 0.141, $p = 0.015$); high serum

creatinine (Pearson's correlation = -0.114, $p = 0.049$); high serum potassium (Spearman's correlation = -0.148, $p = 0.01$); presence of sepsis (Pearson's correlation = 0.827, $p < 0.001$).

Many studies had shown that hyperkalemia is associated with increased mortality in CKD patients [29]. Despite the normal range of hemoglobin, patients with CKD and low quintile hemoglobin (13.2 g/dl, range 7.6 to 14.6 g/dl), had increased risk for CKD-related death [30]. In patients with CKD not under hemodialysis, Erythropoiesis-Stimulating Agents (ESA), users had higher rates of death and cardiovascular events than non-users of ESA [31]. In contrast, a systemic review and meta-analysis involving ESA users showed an increased risk of mortality in the high Hb levels (RR 1.18; 95% CI 1.02 to 1.37). Overall, compared with low Hb levels (<10 g/dl), high Hb levels (>13.0 g/dl) are associated with increased risk of hypertension (RR = 1.40; 95% CI = 1.11 - 1.75), stroke (RR = 1.73; 95% CI = 1.31 - 2.29), and hospitalizations (RR = 1.07; 95% CI = 1.01 - 1.14) [31]. The National Kidney Foundation Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines recommend target hemoglobin levels in the range 11 to 12 g/dl, whereas hemoglobin >13 g/dl should be avoided [32]. In comparison

to patients without CKD or with other chronic medical conditions, CKD patients had the highest 90-day mortality [33]. Mortality due to sepsis was about 50-fold higher in dialysis patients compared with the general population [34].

CKD increases the number of visits to emergency services and hence increases the resource use of emergency services. Certain factors like anemia, hyperkalemia, acid-base disorder, and the presence of other chronic health conditions like hypertension, diabetes, glomerulonephritis not only leads to the poor outcome of these patients but also increases the emergency visits. The major limitation of this study is that it included laboratory parameters of CKD patients during an emergency visit. We studied mortality during the emergency room stay only. So we recommend to include clinical parameters, longer duration of follow up, and larger studies with randomization during further studies.

CONCLUSION

There were factors like anemia, hyperkalemia, increased serum creatinine, and sepsis which increased emergency visits and mortality of CKD patients. Therefore these factors need to be well controlled to decrease mortality and emergency visits of these patients.

DECLARATIONS

Authorship

Dr. TMS, Dr. PNP and Dr. LB did paper conceptualization, ethical clearance, data collection, and submission to the journal; Dr. RPN and Dr. RG did data analysis and manuscript preparation. All authors consented to the publication of this article.

Funding

Whole research was funded by authors.

Availability of data and materials

All the data and materials will be available from the corresponding author upon reasonable request.

Informed consent

Verbal consent was taken from the patient and/or legal guardian if the patient was critically ill.

References

- Sharma SK, Dhakal S, Thapa L, Ghimire A, Tamrakar R, Chaudhary S, Deo R, Manandhar D, Perico N, Perna A, Remuzzi G, Lamsal M. Community-based screening for chronic kidney disease, hypertension and diabetes in Dharan. *JNMA J Nepal Med Assoc.* 2013 Jan-Mar;52(189):205-212. PMID: 23591297.
- Agrawal RK, Hada R, Khakurel S, Baral A. Analysis of Eight Years Data of Renal Disorders in a Tertiary Care Hospital in Nepal. *Postgrad Med J NAMS* 2009 July-Dec;9:28-34.
- Snyder JJ, Collins AJ. Association of preventive health care with atherosclerotic heart disease and mortality in CKD. *J Am Soc Nephrol.* 2009 Jul;20(7):1614-1622. doi: 10.1681/ASN.2008090954. Epub 2009 May 7. PMID: 19423688; PMCID: PMC2709678.
- Ronskley PE, Tonelli M, Manns BJ, Weaver RG, Thomas CM, MacRae JM, Ravani P, Quinn RR, James MT, Lewanczuk R, Hemmelgarn BR. Emergency Department Use among Patients with CKD: A Population-Based Analysis. *Clin J Am Soc Nephrol.* 2017 Feb 7;12(2):304-314. doi: 10.2215/CJN.06280616. Epub 2017 Jan 24. PMID: 28119410; PMCID: PMC5293336.
- Nand N, Kala V, Sharma M. Spectrum of chronic kidney disease in a tertiary care hospital in Haryana. *Journal, Indian Acad Clin Med* 2015 Jan;16(2):110-113.
- Malekmakan L, Malekmakan A, Daneshian A, Pakfetrat M, Roosbeh J. Hypertension and diabetes remain the main causes of chronic renal failure in Fars Province, Iran 2013. *Saudi J Kidney Dis Transpl.* 2016 Mar;27(2):423-424. doi: 10.4103/1319-2442.178594. PMID: 26997406.
- Ghaderian SB, Beladi-Mousavi SS. The role of diabetes mellitus and hypertension in chronic kidney disease. *J Renal Inj Prev.* 2014 Dec 1;3(4):109-110. doi: 10.12861/jrip.2014.31. PMID: 25610891; PMCID: PMC4301389.
- Pourhoseingholi MA, Vahedi M, Rahimzadeh M. Sample size calculation in medical studies. *Gastroenterol Hepatol Bed Bench.* 2013 Winter;6(1):14-17. PMID: 24834239; PMCID: PMC4017493.
- Thompson S, James M, Wiebe N, Hemmelgarn B, Manns B, Klarenbach S, Tonelli M; Alberta Kidney Disease Network. Cause of Death in Patients with Reduced Kidney Function. *J Am Soc Nephrol.* 2015 Oct;26(10):2504-2511. doi: 10.1681/ASN.2014070714. Epub 2015 Mar 2. PMID: 25733525; PMCID: PMC4587695.
- Dobre M, Rahman M, Hostetter TH. Current status of bicarbonate in CKD. *J Am Soc Nephrol.* 2015 Mar;26(3):515-523. doi: 10.1681/ASN.2014020205. Epub 2014 Aug 22. PMID: 25150154; PMCID: PMC4341479.
- Marano M, Marano S, Gennari FJ. Beyond bicarbonate: complete acid-base assessment in patients receiving intermittent hemodialysis. *Nephrol Dial Transplant.* 2017 Mar 1;32(3):528-533. doi: 10.1093/ndt/gfw022. PMID: 27001688.
- Turakhia MP, Blankstijn PJ, Carrero JJ, Clase CM, Deo R, Herzog CA, Kasner SE, Passman RS, Pecoits-Filho R, Reinecke H, Shroff GR, Zareba W, Cheung M, Wheeler DC, Winkelmayer WC, Wanner C; Conference Participants. Chronic kidney disease and arrhythmias: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Eur Heart J.* 2018 Jun 21;39(24):2314-2325. doi: 10.1093/eurheartj/ehy060. PMID: 29522134; PMCID: PMC6012907.
- Sedgwick P. Receiver operating characteristic curves. *BMJ.* 2011;343:d4302. doi: 10.1136/bmj.d4302.
- WHO. Noncommunicable diseases: The slow-motion disaster. WHO. 2017 June 25.
- Luyckx VA, Tonelli M, Stanifer JW. The global burden of kidney disease and the sustainable development goals. *Bull World Health Organ.* 2018 Jun 1;96(6):414-422D. doi: 10.2471/BLT.17.206441. Epub 2018 Apr 20. PMID: 29904224; PMCID: PMC5996218.
- Sigdel MR, Pradhan R. Chronic Kidney Disease in a Tertiary Care Hospital in Nepal. *J Inst Med.* 2018 Aug; 40(2): 104-111.
- Alsuwaida AO, Farag YM, Al Sayyari AA, Mousa D, Alhejaili F, Al-Harbi A, Housawi A, Mittal BV, Singh AK. Epidemiology of chronic kidney disease in the Kingdom of Saudi Arabia (SEEK-Saudi investigators) - a pilot study. *Saudi J Kidney Dis Transpl.* 2010 Nov;21(6):1066-1072. PMID: 21060175.
- Hamat I, Abderraman GM, Tondi ZMM, Youssef M, Cisse MM, Tara F, Ka EF, Niang A, Diouf B. Epidemiological Profile of Chronic Kidney Disease at the General Hospital of National Reference of N'Djamena (Chad). *Open J Nephrol.* 2016 Sep;6(3):78-85. DOI: 10.4236/ojneph.2016.63010.
- Singh AK, Farag YM, Mittal BV, Subramanian KK, Reddy SR, Acharya VN, Almeida AF, Channakeshavamurthy A, Ballal HS, P G, Issacs R, Jasuja S, Kirpalani AL, Kher V, Modi GK, Nainan G, Prakash J, Rana DS, Sreedhara R, Sinha DK, V SB, Sunder S, Sharma RK, Seetharam S, Raju TR, Rajapurkar MM. Epidemiology and risk factors of chronic kidney disease in India - results from the SEEK (Screening and Early Evaluation of Kidney Disease) study. *BMC Nephrol.* 2013 May 28;14:114. doi: 10.1186/1471-2369-14-114. PMID: 23714169; PMCID: PMC3848478.
- Rajapurkar MM, John GT, Kirpalani AL, Abraham G, Agarwal SK, Almeida AF, Gang S, Gupta A, Modi G, Pahari D, Pisharody R, Prakash J, Raman A, Rana DS, Sharma RK, Sahoo RN, Sakhuja V, Tatapudi RR, Jha V. What do we know about chronic kidney disease in India: first report of the Indian CKD registry? *BMC Nephrol.* 2012 Mar 6;13:10. doi: 10.1186/1471-2369-13-10. PMID: 22390203; PMCID: PMC3350459.

21. Jiwa M, Chakera A, Dadich A, Meng X, Kanjo E. The profile of patients with chronic kidney disease who regularly present at an Australian general practice. *Curr Med Res Opin.* 2016;32(1):183-189. doi: 10.1185/03007995.2015.1109505. Epub 2015 Nov 17. PMID: 26473553.
22. Poudel B, Yadav BK, Shrestha R, Mittal A, Jha B, Raut KB. Assessment of chronic kidney disease in Nepalese people with hypertension. *Nepal Med Coll J.* 2012 Mar;14(1):25-30. PMID: 23441490.
23. Einhorn LM, Zhan M, Hsu VD, Walker LD, Moen MF, Seliger SL, Weir MR, Fink JC. The frequency of hyperkalemia and its significance in chronic kidney disease. *Arch Intern Med.* 2009 Jun 22;169(12):1156-1162. doi: 10.1001/archinternmed.2009.132. PMID: 19546417; PMCID: PMC3544306.
24. An JN, Lee JP, Jeon HJ, Kim DH, Oh YK, Kim YS, Lim CS. Severe hyperkalemia requiring hospitalization: predictors of mortality. *Crit Care.* 2012 Nov 21;16(6):R225. doi: 10.1186/cc11872. PMID: 23171442; PMCID: PMC3672605.
25. Hayes J, Kalantar-Zadeh K, Lu JL, Turban S, Anderson JE, Kovesdy CP. Association of hypo- and hyperkalemia with disease progression and mortality in males with chronic kidney disease: the role of race. *Nephron Clin Pract.* 2012;120(1):c8-16. doi: 10.1159/000329511. Epub 2011 Dec 2. PMID: 22156587; PMCID: PMC3267990.
26. Hemachandar R. Practice pattern of hemodialysis among end-stage renal disease patients in Rural South India: A single-center experience. *Saudi J Kidney Dis Transpl.* 2017 Sep-Oct;28(5):1150-1156. doi: 10.4103/1319-2442.215134. PMID: 28937077.
27. Nguyen OK, Vazquez MA, Charles L, Berger JR, Quiñones H, Fuquay R, Sanders JM, Kapinos KA, Halm EA, Makam AN. Association of Scheduled vs Emergency-Only Dialysis With Health Outcomes and Costs in Undocumented Immigrants With End-stage Renal Disease. *JAMA Intern Med.* 2019 Feb 1;179(2):175-183. doi: 10.1001/jamainternmed.2018.5866. PMID: 30575859; PMCID: PMC6439652.
28. Fox KM, Yee J, Cong Z, Brooks JM, Petersen J, Lamerato L, Gandra SR. Transfusion burden in non-dialysis chronic kidney disease patients with persistent anemia treated in routine clinical practice: a retrospective observational study. *BMC Nephrol.* 2012 Jan 24;13:5. doi: 10.1186/1471-2369-13-5. PMID: 22273400; PMCID: PMC3283448.
29. Kovesdy CP. Updates in hyperkalemia: Outcomes and therapeutic strategies. *Rev Endocr Metab Disord.* 2017 Mar;18(1):41-47. doi: 10.1007/s11154-016-9384-x. PMID: 27600582; PMCID: PMC5339065.
30. Leeder SR, Mitchell P, Liew G, Rochtchina E, Smith W, Wang JJ. Low hemoglobin, chronic kidney disease, and risk for coronary heart disease-related death: the Blue Mountains Eye Study. *J Am Soc Nephrol.* 2006 Jan;17(1):279-284. doi: 10.1681/ASN.2005050553. Epub 2005 Nov 30. PMID: 16319191.
31. Stirnadel-Farrant HA, Luo J, Kler L, Cizman B, Jones D, Brunelli SM, Cobitz AR. Anemia and mortality in patients with nondialysis-dependent chronic kidney disease. *BMC Nephrol.* 2018 Jun 11;19(1):135. doi: 10.1186/s12882-018-0925-2. PMID: 29890958; PMCID: PMC5996482.
32. KDOQI. KDOQI Clinical Practice Guideline and Clinical Practice Recommendations for anemia in chronic kidney disease: 2007 update of hemoglobin target. *Am J Kidney Dis.* 2007 Sep;50(3):471-530. doi: 10.1053/j.ajkd.2007.06.008. PMID: 17720528.
33. Mansur A, Mulwande E, Steinau M, Bergmann I, Popov AF, Ghadimi M, Beissbarth T, Bauer M, Hinz J. Chronic kidney disease is associated with a higher 90-day mortality than other chronic medical conditions in patients with sepsis. *Sci Rep.* 2015 May 21;5:10539. doi: 10.1038/srep10539. PMID: 25995131; PMCID: PMC4650757.
34. Sarnak MJ, Jaber BL. Mortality caused by sepsis in patients with end-stage renal disease compared with the general population. *Kidney Int.* 2000 Oct;58(4):1758-64. doi: 10.1111/j.1523-1755.2000.00337.x. PMID: 11012910.

How to cite this article: Shrestha TM, Bhusal L, Neupane RP, Ghimire R, Prasad PN. Risk Factors Associated with Mortality among Chronic Kidney Disease Patients on Regular Hemodialysis Presenting in Emergency Services. *J Biomed Res Environ Sci.* 2021 Jan 05; 2(1): 001-007. doi: 10.37871/jbres1178, Article ID: JBRES1178