Pale skin: An adrenal cancer?

Ana VALEA^{1,2}, Mara CARSONTE^{3,4}, Eugenia PETROVA^{3,4}, Mihai Cristian DUMITRASCU^{4,5},
Florica SANDRU^{4,6}

¹ Clinical County Hospital, Cluj-Napoca, Romania

² "Iuliu Hatieganu" University of Medicine and Pharmacy, Cluj-Napoca, Romania

³ "C.I. Parhon" National Institute of Endocrinology, Bucharest, Romania

⁴ Carol Davila University of Medicine and Pharmacy, Bucharest, Romania

⁵ Emergency University Hospital, Bucharest, Romania

⁶ Elias Emergency University Hospital, Bucharest, Romania

ABSTRACT

Adrenal carcinoma is a complex orphan disease with a very poor prognosis. The evolution is rapid and in cases with suggestive symptoms the diagnosis may be established earlier in order to improve the outcome. One third of the patients are negative for a clear hormonal expression or they have an atypical presentation. We introduce an adult patient diagnosed with adrenal carcinoma and a rapid lethal outcome who initially presented with pale skin and non-specific abdominal pain. A multidisciplinary team is essential for adequate diagnosis and prompt intervention. Severe bleeding at the level of the adrenal carcinoma causing symptomatic anaemia is a rare event. The prognosis in adrenocortical carcinoma is severe and the presence of humoral hypercalcemia represents a sign of short survival time.

Keywords: pale skin, abdominal tumour, adrenal cancer

Abbreviations

ACTH = adrenocorticotrophic hormone

BMD = bone mineral density

DXA = Dual-Energy X-Ray Absorptiometry

MEN 2A = multiple endocrine neoplasia

PTH = parathormone

PTHrP = PTH related protein

SD = standard deviation

INTRODUCTION

Adrenal carcinoma is a complex orphan disease with a very poor prognosis (1,2). The evolution is rapid and in cases with suggestive symptoms the diagnosis may be established earlier in order to improve the outcome which is still dramatic in most cases (1,2). One third of the patients are negative for a clear hormonal expression or they have an atypical presentation (3,4). The risk of malignancy at the level of an adrenal incidental involving an adrenal carcinoma is 0.1% (5,6). Typically the abdominal pain and mass is associated with Cushing's syndrome but some other compli-

cations are related to anaemia, rapidly developing hirsutim in females, persistent infections of complications of the high blood pressure like stroke and acute myocardial infarct (7,8). After tumour removal the prognosis is improved through Cushing's syndrome control and medical therapy continues with mitotane (9,10,11).

AIM

We introduce an adult patient case diagnosed with adrenal carcinoma and a rapid lethal outcome who initially presented with pale skin and non-specific abdominal pain.

Corresponding author:

Lecturer Mara Carsote, MD, PhD

E-mail: carsote_m@hotmail.com

Article History:

Received: 5 November 2019

Accepted: 25 November 2019

MATERIAL AND METHOD

This is a case presentation type of report. The endocrine and imaging profile is provided. The investigations required a multidisciplinary team.

CASE REPORT

Presentation

This is a 49-year old male coming from endemic area with irrelevant family and personal medical background. He was admitted in different medical services for developing within weeks pale skin, asthenia and non-specific abdominal pain. After several hospitalisations, the diagnosis was not established yet, so he had an abdominal and thorax computed tomography done without contrast and an adrenal tumour of 10 cm was detected.

Endocrine panel

On admission, the investigations confirmed anaemia with hyposideremia (haemoglobin of 8 g/dl, normal above 14 g/dl) with rapid decreased of the haemoglobin (within days). The endocrine profile showed high serum calcium of 11.7 g/dl (normal between 8.5 and 10.2 mg/dl) with suppressed PTH (parathormone) of 1 pg/ml (normal levels between 15 and 65 pg/ml) with an increase of total serum alkaline phosphatase 1.5 time above normal upper limit and normal level of 25-hydroxyvitamin D of 40 ng/ml (normal: 30-100 ng/ml).

He associates multinodular goitre with small nodules normal thyroid function and negative antibodies against thyroid (Figure 1). The serum calcitonin is two times above the normal upper limit.

FIGURE 1. Thyroid ultrasound shows micronodular goitre on an adult male with adrenocortical cancer and mild increase of calcitonin

The morning plasma cortisol is high of 35 µg/dl (normal less than 21 µg/dl) and there is no sup-

pression after the administration of 2 day X 2 mg dexamethasone during test. ACTH (adrenocorticotrophic hormone) is mid-normal of 35 pg/ml (normal: 3-66 pg/ml) associated with normal serum levels of chromogranin A, plasma metanephrines and normetanephrines.

Other imaging scans were done. Whole body bone scintigram was negative for bone metastases (Figure 2). DXA (Dual-Energy X-Ray Absorptiometry) showed normal BMD (bone mineral density) for age (Figure 3).

FIGURE 2. Whole body bone scintigram negative for bone metastasis in a patient with high calcium and suppressed PTH associating a malignant right adrenal tumour

Abdominal ultrasound and then computed tomography was re-done and showed a rapid right adrenal tumour increase to 13 cm maximum diameter and liver invasion in addition to intra-tumour necrosis and secondary lung spreading (Figure 4).

Outcome

The patient was transferred for palliative tumour resection but he died within days before the intervention because of massive intra-tumour haemorrhage.

DISCUSSION

Hipercalcitoninemia

The increase of calcitonin in this case may be considered a non-specific neuroendocrine tu-

FIGURE 3. Normal central DXA (lumbar spine, GE Lunar Prodigy device)

FIGURE 4. Large right adrenal tumour with liver invasion at abdominal ultrasound (different sections)

mour. It is less possible to be associated with a synchronous medullary thyroid cancer since the values of calcitonin were only mildly increased and the thyroid ultrasound pattern was not suggestive. Fine needle aspiration might have been an investigational tool but not a feasible option due to the emergency aspect. MEN 2A (multiple endocrine neoplasia) syndrome also involves at least one adrenal tumour and increased calcitonin but this was not such a case (11). However, adrenocortical carcinoma as this case is usually included in classification of neuroendocrine tumours (12).

Cancer-related hypercalcemia

In this case hypercalcemia was a paraneoplastic expression which was not caused by bone metastasis as revealed by whole body bone scintigram but most probable by adrenal cancer-derivate PTH related protein (PTHrP) (13). The suppression of

PTH seems consecutive to hypercalcemia while DXA exam was irrelevant for the present circumstance (14). Up to 30% of malignancies associate high serum calcium and it usually represents a poor prognosis (14). PTHrP production or humoral hypercalcemia represents the most common subtype of hypercalcemia but PTHrP is not assessed in daily practice (15,16).

Adrenocortical carcinoma-associated anaemia

In this particular case the clinical presentation a part from abdominal pain and mass was marked by rapidly deterioration of haemoglobin levels. All the hematologic tests showed that the only cause was most probably intra-tumour haemorrhage. The potential alternative mechanisms are: primary adrenal insufficiency in cases with adrenal cancer and contralateral metastases that display more than 90% from adrenal parenchyma, male hypog-

onadism, and bone marrow metastasis (17). Cystic transformation of an adrenal tumour because of the necrosis is possible in adrenocortical carcinoma but it is more frequent in pheochromocytoma, especially in children (18,19).

Poor prognosis of adrenal cancer

The present case introduces a malignant adrenal tumour (although there was not pathological confirmation) with a fatal outcome within days since presentation at Endocrinology Department. Generally a poor prognosis is expected in this type of malignancy despite the progress of technology

and therapy (20). Cortisol producing subtype of adrenal cancer has the highest rate of mortality (as was this case) when compared to the subtypes with other endocrine profile (21).

CONCLUSION

Severe bleeding at the level of the adrenal carcinoma causing symptomatic anaemia is a rare event. The prognosis in adrenocortical carcinoma is poor and the presence of humoral hypercalcaemia represents a sign of short survival time as well as active cortisol production.

Conflict of interest: none declared
Financial support: none declared

REFERENCES

- Hodgson A, Pakbaz S, Mete O. A Diagnostic Approach to Adrenocortical Tumors. *Surg Pathol Clin*. 2019 Dec; 12(4):967-995.
- Vaidya A, Nehs M, Kilbridge K. Treatment of Adrenocortical Carcinoma. *Surg Pathol Clin*. 2019 Dec;12(4):997-1006.
- Poiana C, Chirita C, Carsote M, Hortopan D, Ioachim D, Corneci CM, Stanescu B. Adrenal and Pituitary Incidentalomas in a Case of Cushing's Syndrome. *Chirurgia* 2013; 6(108):886-891.
- Gheorghiu M, Hortopan D, Dumitrascu A, Caragheorgheopol A, Stefanescu A, Trifanescu R, Niculescu D, Baci I, Carsote M, Poiana C, Badiu C, Coculescu M. Age-related endocrine tumors: Nonfunctioning adrenal tumors as compared to pituitary adenomas. *Acta Endocrinologica*, 2009;V(3):371-384.
- Gheorghisan-Galateanu AA, Carsote M, Valea A. Incidentaloma: From general practice to specific endocrine frame. *J Pak Med Assoc*. 2017.67(6):917-922.
- Else T, Kim AC, Sabolch A, Raymond VM, Kandathil A, Caoili EM, Jolly S, Miller BS, Giordano TJ, Hammer GD. Adrenocortical carcinoma. *Endocr Rev*. 2014 Apr; 35(2):282-326.
- Hodgson A, Pakbaz S, Mete O. A Diagnostic Approach to Adrenocortical Tumors. *Surg Pathol Clin*. 2019 Dec; 12(4):967-995.
- Wengander S, Trimpou P, Papakokkinou E, Ragnarsson O. The incidence of endogenous Cushing's syndrome in the modern era. *Clin Endocrinol (Oxf)*. 2019 Aug; 9
- Paduraru DN, Nica A, Carsote M, Valea A. Adrenalectomy for Cushing's syndrome: do's and don'ts. *Journal of Medicine and Life*. 2016;4(9):334-341.
- Valea A, Ghevan C, Carsote M, Albu SE, Georgescu CE. Different surgical options in Cushing's disease. *Journal of Surgical Sciences*. 2016; 3(1):39-43.
- Valea A, Radu O, Morar A, Ghemigian A, Carsote M. Synchronous medullary thyroid cancer and primary hyperparathyroidism on a female within the sixth decade of life with positive family history for type 2A MEN syndrome. *Journal of Medical Practice*. 2016; 4(47):11:346-349.
- Strosberg JR Update on the management of unusual neuroendocrine tumors: pheochromocytoma and paraganglioma, medullary thyroid cancer and adrenocortical carcinoma. *Semin Oncol*. 2013 Feb; 40(1):120-33.
- Goldner W. Cancer-Related Hypercalcemia. *J Oncol Pract*. 2016 May;12(5):426-32.
- Radu L, Carsote M, Gheorghisan-Galateanu AA, Preda SA, Calborean V, Stanescu R, Gheorman V, Albulescu DM. Blood Parathyryn and Mineral Metabolism Dynamics. A clinical analyze. *Rev.Chim.* (Bucharest). 2018; 69(10):2754-2758.
- Szymanski JJ, Otrrock ZK, Patel KK, Scott MG. Incidence of humoral hypercalcemia of malignancy among hypercalcemic patients with cancer. *Clin Chim Acta*. 2016 Jan 30; 453:190-3.
- Carsote M, Valea A, Dumitru N, Terzea D, Petrova E, Albu S, Buruiana A, Ghemigian A. Metastases in daily endocrine practice. *Archives of Balkan Medical Union*. 2016; 51(4):476-480.
- Foster M, Nolan RL, Hong HH. Bilateral primary adrenocortical carcinoma complicated by Addisonian crisis: Case report. *Can Assoc Radiol J*. 2001 Aug; 52(4):220-2.
- Poiana C, Carsote M, Chirita C, Terzea D, Paun S, Beuran M. Giant adrenal cyst: Case study. *J Med Life* 2010;3(3):308-313.
- Carsote M, Ghemigian A, Terzea D, Gheorghisan-Galateanu AA, Valea A. Cystic adrenal lesions: Focus on pediatric population (a review). *Clujul Med*. 2017; 90(1):5-12.
- Long SE, Miller BS. Adrenocortical Cancer Treatment. *Surg Clin North Am*. 2019 Aug; 99(4):759-771.
- Vanbrabant T, Fassnacht M, Assie G, Dekkers OM. Influence of hormonal functional status on survival in adrenocortical carcinoma: Systematic review and meta-analysis. *Eur J Endocrinol*. 2018 Dec 1;179(6):429-436.