Poster (E06)

Gestational Trophoblastic Neoplasia

<https://doi.org/10.3802/jgo.2021.32.S1.E06>

Genomic profiles of gestational choriocarcinoma for diagnosis and predicting prognosis

Seongeun Bak, Youn Jin Choi*

The Catholic University of Korea, Seoul St. Mary's Hospital, Seoul, Korea
(yunno@catholic.ac.kr)

Objective: Gestational choriocarcinoma (GC) is one of gestational trophoblastic diseases (GTDs), and a unique cancer that originates in pregnant tissues. GCs are known to be preceded by complete hydatidiform mole (CHM) (50%), previous abortions (25%), or normal/ectopic pregnancies (25%). CHM is considered a preneoplastic condition of GC, and invasive mole (IM) has neoplastic characteristics such as tissue invasion. However, what triggers CHM/IM to progress to GC is not figured out. Genomic alterations of GCs are not well studied because of the rare availability of tumor tissues. The lack of genome-wide alteration data on GTDs, led us to analyze GC genomes by whole exome sequencing, copy number alteration, copy-neutral loss-of-heterozygosity (CN-LOH) profiles in this study.

Methods: Whole-exome sequencing (20 GCs) and/or single-nucleotide polymorphism microarray (29 GCs) were performed. We analyzed CN-LOH in 29 GCs.

Results: Most GCs (25/29) harboring recurrent copy number alterations (CNAs) and gains on 1q21.1-q44 were significantly associated with poor prognosis. We detected five driver mutations in the GCs, most of which were chromatin remodeling gene (*ARID1A*, *SMARCD1*, and *EP300*) mutations but not in common cancer genes such as *TP53* and *KRAS*.

Conclusion: Our data indicate that GCs have unique profiles of CN-LOHs, mutations and CNAs that together differentiate GCs from non-GCs. Practically, CN-LOH and CNA profiles are useful for the molecular diagnosis of GC and the selection of GC patients with poor prognosis for more intensive treatments, respectively.

Poster (E07)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.E07>

Impact of surgical approach on survival and perioperative outcomes in women with stage II endometrial cancer

Xian Li,¹ Gladys Shuk Tak Kwok,² Mandy Man Yee Chu,² Siew Fei Ngu,² Karen Kar Loen Chan,² Hextan Yuen Sheung Ngan,³ Keedon Wong,² Ka Yu Tse^{3,*}
¹The University of Hong Kong - Shenzhen Hospital, Shenzhen, China

²The University of Hong Kong - Queen Mary Hospital, Hong Kong

³The University of Hong Kong, Hong Kong (tseky22@gmail.com)

Objective: To compare the survival and perioperative outcomes between minimally-invasive surgery (MIS) and open surgery in stage II endometrial cancer patients.

Methods: This was a retrospective study involving 205 patients with stage II endometrial cancer in a single institution between 2002–2018. Patients were stratified to either MIS or open surgery groups, and their pathological and clinical data were reviewed.

Results: One hundred fourteen patients had open surgery (55.6%) while 91 patients had MIS (44.4%) including conventional laparoscopic (70.3%), robot-assisted (22.0%) and laparoscopic-assisted vaginal hysterectomy (7.7%). The 5- and 10-year overall survival rate was 94.5% and 93.4% for the MIS group, while 89.4% and 85.8% for the open surgery group ($p=0.219$ and $p=0.127$). The 5- and 10-year progression-free survival rate was 91.2% and 90.1% for the MIS group, while 86.8% and 81.5% for the open surgery group, without statistical significance, $p=0.395$ and $p=0.372$. The uterine size (5.7 ± 2.3 vs. 8.0 ± 4.0 weeks; $p=0.007$) and tumor size (2.5 ± 1.7 vs. 4.5 ± 3.3 cm; $p=0.001$) were significantly smaller in the MIS group than in the open group. MIS was also associated with less blood loss (198.6 ± 4.3 vs. 684.8 ± 614.0 mL; $p=0.001$), fewer wound complications (1.1% vs. 5.6%; $p=0.034$), and shorter hospital stay (3 vs. 5 days; $p=0.031$) compared to open surgery. However, there is no significant difference in operation duration in the two groups (224 ± 97 vs. 227 ± 94 minutes; $p=0.554$).

Conclusion: The oncological outcomes and perioperative complication rates of patients undergoing MIS were not inferior to open surgery for stage II endometrial carcinoma.

Poster (E08)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.E08>

Incorporation of protein biomarkers to the ProMisE molecular classifier for further prognostication in endometrial cancer

Se Ik Kim, Hyunji Lim, Cheol Lee, Jae-Weon Kim, Maria Lee*

Seoul National University College of Medicine, Seoul, Korea
(marialeemd@gmail.com)

Objective: The Proactive Molecular Risk Classifier for Endometrial Cancer (ProMisE) is a clinically applicable prognostication method

in endometrial cancer. In order to classify endometrial cancer more precisely, especially those with nonspecific molecular profile (NSMP), we aimed to determine the prognostic significance of 3 additional proteins in the institution's endometrial cancer cohort: β -catenin, L1 cell-adhesion molecule (L1CAM), and programmed death-ligand 1 (PD-L1).

Methods: Endometrial cancer patients, surgically treated between 2014 and 2018 at our institution, underwent ProMisE classification using immunohistochemical (IHC) staining for p53 and 4 mismatch repair (MMR) proteins (MLH1, PMS2, MSH2, and MSH6) and hot spot sequencing of DNA polymerase epsilon (POLE) exonuclease domain via droplet digital polymerase chain reaction. Expression of β -catenin, L1CAM, and PD-L1 were studied by IHC staining. Then, we investigated the relationship between protein expression and progression-free survival (PFS).

Results: In total, 156 patients were included in this analysis: 136 (87.2%) and 107 (68.6%) had an endometrioid histologic type and early-stage disease, respectively. ProMisE classification assigned 45 (28.8%), 16 (10.3%), 70 (44.9%), and 25 (16.0%) to MMRd, POLEmut, NSMP (p53wt), and p53abn, respectively. Using 5% of nuclear staining as a cut-off, we observed no difference in PFS between positive and negative β -catenin expression groups ($p=0.465$). Expression of PD-L1 was most frequent in the MMRd subgroup (48.9%). Multivariate analysis adjusting confounders revealed that PD-L1 $\geq 1\%$ of tumor cells was identified as a favorable prognostic biomarker for PFS (adjusted hazard ratio [HR]=0.394; 95% confidence interval [CI]=0.182–0.854; $p=0.018$). However, PD-L1 expression was not predictive of prognosis among the NSMP subgroup. Using 10% of staining as a cut-off, high expression of L1CAM was most frequent in the p53abn subgroup (32.0%). High expression of L1CAM was identified as a poor prognostic biomarker for PFS (adjusted HR=3.486; 95% CI=1.577–7.707; $p=0.002$). L1CAM expression was predictive of prognosis among the NSMP subgroup. In subgroup analyses in endometrioid tumors, consistent results were observed.

Conclusion: While β -catenin was not informative for risk stratification, expression of PD-L1 and L1CAM were associated with disease recurrence. Especially, L1CAM expression further stratifies the risk of disease recurrence among the NSMP subgroup.

Poster (E09)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.E09>

Interim analysis of immune-related endpoints of the mismatch repair deficient and proficient endometrial cancer (EC) cohorts from the garnet study

<https://asgo2021.org>

Yoshiaki Kawano,^{1,*} Ana Oaknin,² Lucy Gilbert,³ Anna Tinker,⁴ Renaud Sabatier,⁵ Valentina Boni⁶

¹GlaxoSmithKline, Brentford, UK (yoshiaki.x.kawano@gsk.com)

²Vall d'Hebron Institute of Oncology, Barcelona, Spain

³McGill University, Montreal, Canada

⁴BC Cancer, Vancouver, Canada

⁵Institut Paoli Calmettes, Marseille, France

⁶Hospital Universitario HM Sanchinarro, Madrid, Spain

Objective: GARNET (NCT02715284) is assessing the antitumor activity and safety of dostarlimab, an anti-programmed death-1 receptor therapy, in patients with solid tumors. We report investigator-assessed (IA) efficacy endpoints for endometrial cancer (EC) patients.

Methods: A phase 1, multicenter, open-label, single-arm, dose-escalation and cohort-expansion study. In 2 independent expansion cohorts of patients with recurrent/advanced EC (deficient mismatch repair [dMMR] and mismatch repair proficient [MMRp] determined by immunohistochemistry) that progressed on/after a platinum regimen, patients received dostarlimab 500 mg IV Q3W for 4 cycles, then 1,000 mg Q6W until disease progression, discontinuation, or withdrawal. Primary endpoints of objective response rate (ORR), disease control rate (DCR), and duration of response (DOR) using RECIST v1.1 were reported previously (Oaknin ESMO 2020 LBA36). Immune-related (ir) ORR, irDCR, and irDOR by irRECIST, based on IA, were secondary endpoints.

Results: One hundred twenty-six dMMR and 145 MMRp patients were enrolled and dosed; 110 and 144 patients, respectively, had measurable disease at baseline. After median (range) follow-up of 16.5 (0.03–30.6) and 13.7 (0.03–33.1) months for dMMR and MMRp patients, irORR was 45.5% and 13.9%, irDCR was 63.6% and 42.4%, and irDOR was not reached and 12.2%, respectively.

Conclusion: Efficacy endpoints by RECIST v1.1 and irRECIST were similar. irDCR was particularly interesting in MMRp patients, a group with a poorer prognosis, but the potential benefit awaits confirmation in ongoing randomized controlled studies.

Funding: GSK 213346; editorial support from Fishawack Health, funded by GSK.

This abstract was previously presented at the Society of Gynecologic Oncology 2021 Annual Meeting and is submitted on behalf of the original authors with their permission.

Poster (E10)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.E10>