Surgical film

Surgical Film (F1)

Surgical Techniques & Perioperative Management

<https://doi.org/10.3802/jgo.2021.32.S1.F1>

Removal of metastatic nodes para-aortic area

Mikio Mikami*

Department of Obstetrics and Gynecology, Tokai University School of Medicine, Isehara, Kanagawa, Japan (mmikami@is.icc.u-tokai.ac.jp)

US guideline for cervical cancer recommends CCRT + brachytherapy for IB2 IVA disease if pelvic or para-aortic nodes are positive by radiological imaging or surgical staging. There is no description of lymphadenectomy. ESGO guideline describes that para-aortic lymphadenectomy may be considered in locally advanced cervical cancer with negative paraaortic lymph nodes on imaging for staging purposes and that debulking of suspicious pelvic nodes may be also considered. There is a big difference between the treatment policies for metastatic nodes for cervical cancer. LION trial for advanced ovarian cancer (AOC) concluded systematic pelvic and para-aortic lymphadenectomy, in patients with AOC who had undergone intraabdominal macroscopically complete resection and had normal lymph nodes both before and during surgery, was not associated with longer overall or progression free survival than no lymphadenectomy, and was associated with a higher incidence of postoperative complications. Two randomized controlled trials did not show a survival benefit of lymphadenectomy for patients with low-risk endometrial carcinoma. For patients with intermediate risk, data also has not shown a survival benefit. However, in Japan, Dr. Todo reported from retrospective cohort analysis that combined pelvic and para-aortic lymphadenectomy should be recommended as treatment for patients with endometrial carcinoma of intermediate or high risk of recurrence (Lancet 2010). As described above, especially therapeutic significance of systemic lymphadenectomy is still controversial in gynecological malignancies. What do you think about removing bulky nodes after completing the resection of primary tumor or optimal surgery? In this video, I am going to focus on Bulky nodes, which means clearly identified metastatic nodes. We will try to show you the “meticulous” and “awesome” lymph node dissection by our original video.

Surgical Film (F2)

Endometrial Hyperplasia, Endometrial Intra-epithelial Neoplasia, and Endometrial Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.F2>

Laparoscopic para-aortic lymphadenectomy in endometrial cancer

Atthapon Jaishuen,* Pisutt Srichaikul, Vuthinun Achariyapota, Khemanat Khemworapong

Siriraj Hospital, Bangkok, Thailand (ajaishuen@hotmail.com)

A 46-year-old female underwent total abdominal hysterectomy with right salpingo-oophorectomy and left salpingectomy for her provisionally diagnosed myoma uteri. The pathology report revealed myoma uteri with grade 1 endometrioid carcinoma of endometrium. Myometrial invasion was less than half of the thickness. However, the left fallopian tube had mucosal invasion of cancer. She was referred to our hospital. Six weeks after first operation, she underwent laparoscopic left oophorectomy, pelvic and para-aortic lymphadenectomy. The camera port was set at Li-Huang point with 3 ancillary diamond shaped ports. The procedure started with lysis adhesion and bilateral lymphadenectomy. Para-aortic lymphadenectomy began with opening the peritoneum covering right ureter along common iliac artery and aorta to the level of third part of duodenum. Both sides of peritoneum were sutured with vicryl no 0, hanging with abdominal wall at camera port level. The para-aortic lymph node (LN) was dissected in 4 groups; right below inferior mesenteric artery (IMA), right above IMA, left below IMA and left above IMA, respectively. The postoperative period was uneventful. The patient can be discharged in 2 days. The final pathology was negative of carcinoma in all specimens (0/17 right pelvic LN, 0/30 left pelvic LN, 0/12 right para-aortic below IMA, 0/6 right para-aortic above IMA, 0/11 left para-aortic below IMA and 0/10 left para-aortic above IMA). The pelvic and para-aortic lymphadenectomy in endometrial cancer patient can be achieved. However, this procedure takes times and complication such as; vascular injury, ureteric injury, lymphocyst and lymphedema. We conduct the sentinel LN biopsy study to detect the false negative rate compared to full lymphadenectomy. If this false negative rate is acceptable, we will not do the full lymphadenectomy anymore.

Surgical Film (F3)

Surgical Techniques & perioperative Management

<https://doi.org/10.3802/jgo.2021.32.S1.F3>

Robotic radical hysterectomy without uterine manipulator or vaginal tube

Naery Kim, Eun Jung Yang, A Jin Lee, Kyeong A So, Sun Joo Lee, Tae Jin Kim, Seung Hyuk Shim*

Konkuk University Hospital, Seoul, Korea (20130131@kuh.ac.kr)

The purpose of this study is to introduce robotic radical hysterectomy with tagged uterine suture instead of using a uterine manipulator or vaginal tube. A total of 4 ports were used; first

port was located left at 8 cm from umbilicus, second port was 20 mm sized at umbilicus, third port was located right at 8 cm from umbilicus, and fourth was located right at 8 cm from the third port (near the right flank). Uterus was tied with needle-straightened multifilament Vicryl 2-0 and tagged uterus was manipulated by fourth arm of the robot. If additional traction is required, instrument was inserted through the umbilical trocar site. During operation, the tagged uterus was successfully manipulated and appropriate parametrial space was exposed. Pathologically, all surgical margins were not involved with cancer. No tumor cells were seen in cytologic exam before and after the colpotomy. Robotic radical hysterectomy can be easily and safely done with the traction of tagged uterine suture.

Surgical Film (F4)

Surgical Techniques & Perioperative Management

<https://doi.org/10.3802/jgo.2021.32.S1.F4>

Vaginal assisted laparoscopic radical hysterectomy (nerve sparing) after LACC era

Supachai Raungkaewmanee, Uraiwan Khomphaiboonkij,^{*} Supakorn Pitakkarnkul, Panwad Ratanasrithong, Warangkana Kolaka

National Cancer Institute Thailand, Bangkok, Thailand
(awan1985@hotmail.com)

From Laparoscopic Approach to Cervical Cancer (LACC) trial, patients who underwent minimally invasive radical hysterectomy for early-stage cervical cancer had lower rates of disease-free survival and overall survival and a higher rate of locoregional recurrence. There are potential reasons for the inferior oncologic outcomes, that the routine usage of a uterine manipulator and intracorporeal colpotomy might increase the propensity for tumor spillage, effect of the insufflation gas (CO₂) on tumor-cell growth or spread. We use selection criteria and change some steps of procedure to reduce risk of recurrent disease, such as: 1) tumor size <2 cm, 2) no lymph-vascular space invasion, 3) not using uterine manipulator, 4) transvaginal colpotomy, and 5) closing vaginal cuff transvaginally. Steps of procedure divided into laparoscopic and vaginal parts. Laparoscopic part steps are as follows: 1) after starting pneumoperitoneum with CO₂ pressure 10–12 mmHg, abdominal exploration is performed; 2) bilateral pelvic lymphadenectomy is performed then frozen section; 3) creating paravesical and pararectal space then identifying hypogastric nerve; 4) radical hysterectomy performed as conventional; 5) cutting uterine branch of hypogastric nerve then dissecting laterally; and 6) suturing to mark vaginal wall. Vaginal part follows as: 1) identifying suture marker; 2) opening vaginal cuff by scalpel; 3) removing specimen then painting with povidone

solution; and 4) closing vaginal stump transvaginally. The advantages of minimal invasive surgery should not be abandoned because of the unexpected findings of the LACC trial. We continue to be critically assessed by discussion with patients before surgery. Ongoing and future randomized controlled trials are needed and should be strongly supported.

Surgical Film (F5)

Cervical Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.F5>

Robotic assisted radical vaginal trachelectomy (fertility sparing surgery)

Amany Abdelwadoud Makroum,^{1,*} Jung-Yun Lee²

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Mansoura University, Mansoura, Egypt (amany_makroum@mans.edu.eg)

²Department of Obstetrics and Gynecology, Institute of Women's Life Medical Science, Yonsei University College of Medicine, Seoul, Korea

A 32-year-old nulliparous woman with early-stage invasive cervical adenocarcinoma underwent robotic assisted radical vaginal trachelectomy (fertility sparing surgery). Steps were followed as: 1) Sentinel lymph node biopsy, 2) Paravesical and Pararectal spaces dissection, 3) Ureter & uterine artery identification, 4) Division of anterior vesico-uterine ligaments (bladder pillars), 5) Recto-vaginal space dissection (uterosacral ligaments division), 6) Colpotomy, 7) Cervical amputation, 8) Frozen biopsy, 9) Cervical cerclage, and 10) Utero-vaginal anastomosis.

Surgical Film (F6)

Uterine Sarcoma

<https://doi.org/10.3802/jgo.2021.32.S1.F6>

vNOTE hysterectomy for adenosarcoma in patient with complicated abdominal surgical history

Thanasak Sueblinvong,^{*} Teresa Walsh

Kaiser Permanente, Oakland, CA, USA (Thanasak.Sueblinvong@kp.org)

This is a 72-year-old with postmenopausal bleeding underwent a hysteroscopic dilation and curettage which revealed adenosarcoma. Her underlying medical problems included chronic kidney disease (glomerular filtration rate 19); obesity (body mass index 38) with associated comorbidities; also an abdominal aortic aneurysm (AAA) with a diameter 7.4 cm. Due to newly diagnosis of cancer and vaginal bleeding, her AAA repair process was put on hold. She had a complicated surgical history. In 2016, she had large bowel obstruction due to underlying