port was located left at 8 cm from umbilicus, second port was 20 mm sized at umbilicus, third port was located right at 8 cm from umbilicus, and fourth was located right at 8 cm from the third port (near the right flank). Uterus was tied with needle-straightened multifilament Vicryl 2-0 and tagged uterus was manipulated by fourth arm of the robot. If additional traction is required, instrument was inserted through the umbilical trocar site. During operation, the tagged uterus was successfully manipulated and appropriate parametrial space was exposed. Pathologically, all surgical margins were not involved with cancer. No tumor cells were seen in cytologic exam before and after the colpotomy. Robotic radical hysterectomy can be easily and safely done with the traction of tagged uterine suture.

Surgical Film (F4)

Surgical Techniques & Perioperative Management

<https://doi.org/10.3802/jgo.2021.32.S1.F4>

Vaginal assisted laparoscopic radical hysterectomy (nerve sparing) after LACC era

Supachai Raungkaewmanee, Uraiwan Khomphaiboonkij,^{*} Supakorn Pitakkarnkul, Panwad Ratanasrithong, Warangkana Kolaka

National Cancer Institute Thailand, Bangkok, Thailand
(awan1985@hotmail.com)

From Laparoscopic Approach to Cervical Cancer (LACC) trial, patients who underwent minimally invasive radical hysterectomy for early-stage cervical cancer had lower rates of disease-free survival and overall survival and a higher rate of locoregional recurrence. There are potential reasons for the inferior oncologic outcomes, that the routine usage of a uterine manipulator and intracorporeal colpotomy might increase the propensity for tumor spillage, effect of the insufflation gas (CO₂) on tumor-cell growth or spread. We use selection criteria and change some steps of procedure to reduce risk of recurrent disease, such as: 1) tumor size <2 cm, 2) no lymph-vascular space invasion, 3) not using uterine manipulator, 4) transvaginal colpotomy, and 5) closing vaginal cuff transvaginally. Steps of procedure divided into laparoscopic and vaginal parts. Laparoscopic part steps are as follows: 1) after starting pneumoperitoneum with CO₂ pressure 10–12 mmHg, abdominal exploration is performed; 2) bilateral pelvic lymphadenectomy is performed then frozen section; 3) creating paravesical and pararectal space then identifying hypogastric nerve; 4) radical hysterectomy performed as conventional; 5) cutting uterine branch of hypogastric nerve then dissecting laterally; and 6) suturing to mark vaginal wall. Vaginal part follows as: 1) identifying suture marker; 2) opening vaginal cuff by scalpel; 3) removing specimen then painting with povidone

solution; and 4) closing vaginal stump transvaginally. The advantages of minimal invasive surgery should not be abandoned because of the unexpected findings of the LACC trial. We continue to be critically assessed by discussion with patients before surgery. Ongoing and future randomized controlled trials are needed and should be strongly supported.

Surgical Film (F5)

Cervical Cancer

<https://doi.org/10.3802/jgo.2021.32.S1.F5>

Robotic assisted radical vaginal trachelectomy (fertility sparing surgery)

Amany Abdelwadoud Makroum,^{1,*} Jung-Yun Lee²

¹Department of Obstetrics and Gynecology, Faculty of Medicine, Mansoura University, Mansoura, Egypt (amany_makroum@mans.edu.eg)

²Department of Obstetrics and Gynecology, Institute of Women's Life Medical Science, Yonsei University College of Medicine, Seoul, Korea

A 32-year-old nulliparous woman with early-stage invasive cervical adenocarcinoma underwent robotic assisted radical vaginal trachelectomy (fertility sparing surgery). Steps were followed as: 1) Sentinel lymph node biopsy, 2) Paravesical and Pararectal spaces dissection, 3) Ureter & uterine artery identification, 4) Division of anterior vesico-uterine ligaments (bladder pillars), 5) Recto-vaginal space dissection (uterosacral ligaments division), 6) Colpotomy, 7) Cervical amputation, 8) Frozen biopsy, 9) Cervical cerclage, and 10) Utero-vaginal anastomosis.

Surgical Film (F6)

Uterine Sarcoma

<https://doi.org/10.3802/jgo.2021.32.S1.F6>

vNOTE hysterectomy for adenosarcoma in patient with complicated abdominal surgical history

Thanasak Sueblinvong,^{*} Teresa Walsh

Kaiser Permanente, Oakland, CA, USA (Thanasak.Sueblinvong@kp.org)

This is a 72-year-old with postmenopausal bleeding underwent a hysteroscopic dilation and curettage which revealed adenosarcoma. Her underlying medical problems included chronic kidney disease (glomerular filtration rate 19); obesity (body mass index 38) with associated comorbidities; also an abdominal aortic aneurysm (AAA) with a diameter 7.4 cm. Due to newly diagnosis of cancer and vaginal bleeding, her AAA repair process was put on hold. She had a complicated surgical history. In 2016, she had large bowel obstruction due to underlying

diverticulitis requiring laparotomy sigmoid colectomy, right hemicolectomy, ileocolic anastomosis and sigmoid end colostomy. She subsequently had closure of colostomy, but had a leak with a washout procedure with diverting loop ileostomy. She eventually had closure of ileostomy. A preoperative positron emission tomography scan revealed no distant or locoregional metastasis. Due to her complex abdominal surgery history, decision made to attempt a vaginal approach to avoid extensive adhesions documented. She then underwent vNOTE hysterectomy and bilateral salpingo-oophorectomy (BSO). Intraoperatively extensive laparoscopic lysis of adhesions were able to be performed,

which allowed successful vNOTE hysterectomy and BSO. Due to extensive adhesions a serosal laceration of small bowel was noted and repaired. The entire procedure was completed vaginally. Estimate blood loss was 50 mL. She developed pneumonia on postoperative day (POD) #3 and also congestive heart failure but eventually able to recover and discharge home on POD#10. Her pathology revealed stage IC adenosarcoma of uterus. She has not received post operative adjuvant treatment and now free without disease. vNOTES is a feasible alternative route in oncology patients with complex surgical history requiring hysterectomy.