the ERAS cohort and the legacy non-ERAS cohort (those who received the same types of surgery from the 2 ERAS attending surgeons prior to implementation of the protocols).

Results: A total of 244 patients were evaluated (122 patients in ERAS cohort vs. 122 patients in non-ERAS cohort). The number of vials of opioid analgesia used during the first 2 post-operative days were significantly lower in the ERAS group. The patients in ERAS group reported less post-operative pain, feelings of hunger and thirst, and greater amount of exercise post-operatively. These benefits of the ERAS cohort were more pronounced in the patients who underwent laparotomic surgeries than those who underwent laparoscopic surgeries. The case-matched comparative analysis also showed similar results. The length of hospital stay did not differ between the 2 groups.

Conclusion: The results of this study demonstrated the safety, clinical feasibility and benefits of the ERAS protocols for patients undergoing gynecologic surgeries.

Poster (M03)
Miscellaneous

<https://doi.org/10.3802/jgo.2021.32.S1.M03>

Clinical analysis of multiple primary malignancies in gynecologic cancer patients

Eun Jung Yang, Yoo Jin Kim, A Jin Lee, Nae Ri Kim, Seung Hyuk Shim, Sun Joo Lee, Tae Jin Kim, Kyeong A So*

Konkuk University School of Medicine, Seoul, Korea (20190001@kuh.ac.kr)

Objective: The aim of this study was to evaluate the prevalence and oncologic outcomes in multiple primary malignant tumors (MPMTs) with gynecologic cancers.

Methods: A retrospective study included 1,929 patients diagnosed with gynecologic malignancies at a tertiary institution medical center between August 2005 and April 2021. We analyzed MPMTs in gynecologic malignancies.

Results: The prevalence of MPMTs with gynecologic cancer patients was 8.6% (165/1,929). The median diagnosis period between primary and secondary cancer was 38 months (range 0–420). Among 165 MPMTs patients, 20 patients had multiple primary gynecologic cancers (MPGC), and 145 patients had gynecologic cancer coexisting with non-gynecologic cancer (GNC). The coexistence of endometrial cancer and ovarian cancer (12/20, 60%) was the most common in MPGC patients. The most common non-gynecological cancer in GNC patients was breast cancer (50/145, 34.5%). Synchronous malignancies cases were 48 (29.1%), and metachronous malignancies cases were 117 (70.9%). MPGC patients had more synchronous cancer than GNC patients (50.0% vs. 26.2%, $p=0.030$). In patients with

ovarian cancer, the diagnosis of early stage was significantly higher in MPGC patients than that in GNC patients (87.5% vs. 52.4%, $p=0.031$). The mortality rate during study period was 7.3%, all of which were GNC patients. Gynecologic cancers were a more direct cause of death than non-gynecological cancer (66.7% vs. 33.3%).

Conclusion: Half of MPGC patients had synchronous cancer, which was significantly higher than GNC. Ovarian cancer showed a higher rate of being diagnosed at an early stage in MPGC patients than that in GNC patients. A systematic examination after the diagnosis of primary cancer can help to diagnose secondary primary malignancy at early stage and improve the prognosis of the patients.

Poster (M04)

Miscellaneous

<https://doi.org/10.3802/jgo.2021.32.S1.M04>

Natural killer cell as prognostic factor in Thai gynecologic cancer patients

Suwanit Therasakvichya,* Morakt Kanbun

Faculty of Medicine, Siriraj Hospital, Mahidol University, Bangkok, Thailand (suwanit.the@mahidol.ac.th)

Objective: Natural killer (NK) cells are lymphocytes that are programmed in immune system to recognize altered/ or stressed by malignant transformation. In gynecologic cancers, the prognostic impacts of NK cell have also been published. Thus, this study aims to demonstrate a preliminary report of NK cell levels in correlation with treatment outcomes among Thai gynecologic cancer patients.

Methods: This cohort comparative study was conducted on 122 women with gynecologic cancer who have been treated with chemotherapy at Siriraj Hospital, between March 2011 and March 2012. The assessment of NK cell using flow cytometry (FACscan) with CD16/CD56 surface markers was done prior to the first course of chemotherapy, which reported in the percentage of NK cells and NK cell count. The majority chemotherapy was carboplatin and paclitaxel (PT) regimen every 3 week. The patients were divided into 2 groups based on the outcomes of the treatment: responder group (complete response or partial response), and non-responder group (stable disease, progression of disease, death with disease).

Results: After completion of 6 cycle-chemotherapy, the median follow-up time was 7.3 months (range 1–18 months). The mean percentage of NK cell in responder and non-responder group were $19.5\% \pm 9\%$ and $19.6\% \pm 8\%$, respectively ($p=0.949$). The median NK cell count in responder group was 313 cells/ μ L (range 95–1,255) compared to 282 cells/ μ L (range 66–1,885) in non-responder group ($p=0.276$). The mean of NK cell count in patients with death with disease, alive with disease, and alive without disease were 171 ± 95 ,

326±342, and 400±240, respectively ($p=0.001$). The higher NK cells levels were significantly associated with the survival of the patients in the follow-up time ($p=0.009$).

Conclusion: Pre-chemotherapy NK cell levels failed to demonstrate a significant predictor of response to chemotherapy. However, the higher pre-chemotherapy NK cell levels are associated with favorable patient follow-up outcome. This study is only a preliminary report of NK cell in Thai gynecologic cancer. A well-design and larger population study may be further conducted to clarify the role of this immunologic factor.

Poster (M05)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M05>

Perspective and conception of Thai gynecologic oncologists in palliative care

Panida Mathaveechotikul,* Chuenkamon Charakorn, Arb-aroon Lertkhachonsuk, Navamol Lekskul, Sarikapan Wilailak, Kittiphon Nagaviroj

Ramathibodi Hospital, Bangkok, Thailand (Panidamtv@gmail.com)

Objective: To evaluate the conception and perspective on palliative care (PC) in Thai gynecologic oncologists.

Methods: The online survey was distributed to all certificated Thai gynecologic oncologists. The survey could be accessed via working email address, hyperlink, or QR code during May 2020 and January 2021. A 5-point Likert scale captured the perspectives and concepts of PC. The association between respondents' characteristics and their choices of content in PC, together with their decision making in specified clinical scenarios was analyzed. **Results:** A total of 207 completed surveys from 320 Thai gynecologic oncologists were received (64.6% participation rate). They prospectively a willingness to give the advices to both patients and their families (85.5%), and strongly agreed to introduce PC in any stage of cancer at the time of diagnosis (75.8%). The numbers of their palliative cases per year were 5–20 (57.97%) and the PC teams were available in their hospitals. They decided to offer early PC and do-not-resuscitate (DNR), especially for the elders, or patients with advance stages, or recurrent disease. We found that gynecologic oncologists who previously experienced a PC training did not show any difference in decision making in specified clinical scenarios, compared with who did not.

Conclusion: Thai gynecologic oncologists responded to the conceptions and perspectives in PC. Their concepts of early and willingness to offer a PC especially in the elders, advanced stage, or recurrent patients were proven, regardless of the experience in PC training.

Poster (M06)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M06>

Prevalence of malnourishment and predictive factors associated with the nutritional status of gynecologic cancer patients undergoing chemotherapy: a cross-sectional analysis

Khemmanat Sanguanwongthong, Prapaporn Suprasert*

Chiang Mai University, Chiang Mai, Thailand (psuprase@gmail.com)

Objective: To investigate the prevalence of malnourishment and predictive factors for nutritional status of gynecologic cancer patients treated with chemotherapy.

Methods: One hundred and one gynecologic cancer patients treated with chemotherapy between April 2020 and February 2021 were interviewed for Patient Generated-Subjective Global Assessment (PG-SGA). Clinical data including body mass index (BMI), underlying disease, the diagnosis, FIGO staging, the recurrent status, the surgical intervention, the performance status, the chemotherapy detail, the total number of lymphocytes per cubic milliliter (TLC) and serum albumin were collected. Levels of TLC and serum albumin were calculated for an optimal cut-off point by using receiver operating characteristic curves. The clinicopathological variables were compared using univariate and multivariate analysis to identify the independent predictive factors for malnourishment status.

Results: The prevalence of well, moderate and severe nourishment were 73.3%, 18.8% and 7.9%, respectively. The optimal cut-off points for TLC was 1,450 cells/ μ L and for albumin was 3.9 g/dL. The univariate analysis indicated that the number of present chemotherapy <3 cycles, albumin in level <3.95 g/dL, BMI <25 kg/m² TLC <1,450 cells/ μ L, anemia and no neutropenia were significant associated factors for malnourishment. However, only serum albumin <3.95 g/dL was independent associated factor for malnourishment (adjusted odds ratios = 6.709; 95% confidence interval = 2.113–21.304).

Conclusion: Nearly one-fourth of patients with gynecologic cancer who on going treatment with chemotherapy were found malnourishment. The only independent associated factor related to the nutritional status was serum albumin <3.95 g/dL.

Poster (M07)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M07>

Validity of the Comprehensive Score for Financial Toxicity (COST) tool in gynecologic cancer patients