326±342, and 400±240, respectively ($p=0.001$). The higher NK cells levels were significantly associated with the survival of the patients in the follow-up time ($p=0.009$).

Conclusion: Pre-chemotherapy NK cell levels failed to demonstrate a significant predictor of response to chemotherapy. However, the higher pre-chemotherapy NK cell levels are associated with favorable patient follow-up outcome. This study is only a preliminary report of NK cell in Thai gynecologic cancer. A well-design and larger population study may be further conducted to clarify the role of this immunologic factor.

Poster (M05)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M05>

Perspective and conception of Thai gynecologic oncologists in palliative care

Panida Mathaveechotikul,* Chuenkamon Charakorn, Arb-aroon Lertkhachonsuk, Navamol Lekskul, Sarikapan Wilailak, Kittiphon Nagaviroj

Ramathibodi Hospital, Bangkok, Thailand (Panidamtv@gmail.com)

Objective: To evaluate the conception and perspective on palliative care (PC) in Thai gynecologic oncologists.

Methods: The online survey was distributed to all certificated Thai gynecologic oncologists. The survey could be accessed via working email address, hyperlink, or QR code during May 2020 and January 2021. A 5-point Likert scale captured the perspectives and concepts of PC. The association between respondents' characteristics and their choices of content in PC, together with their decision making in specified clinical scenarios was analyzed. **Results:** A total of 207 completed surveys from 320 Thai gynecologic oncologists were received (64.6% participation rate). They prospectively a willingness to give the advices to both patients and their families (85.5%), and strongly agreed to introduce PC in any stage of cancer at the time of diagnosis (75.8%). The numbers of their palliative cases per year were 5–20 (57.97%) and the PC teams were available in their hospitals. They decided to offer early PC and do-not-resuscitate (DNR), especially for the elders, or patients with advance stages, or recurrent disease. We found that gynecologic oncologists who previously experienced a PC training did not show any difference in decision making in specified clinical scenarios, compared with who did not.

Conclusion: Thai gynecologic oncologists responded to the conceptions and perspectives in PC. Their concepts of early and willingness to offer a PC especially in the elders, advanced stage, or recurrent patients were proven, regardless of the experience in PC training.

Poster (M06)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M06>

Prevalence of malnourishment and predictive factors associated with the nutritional status of gynecologic cancer patients undergoing chemotherapy: a cross-sectional analysis

Khemmanat Sanguanwongthong, Prapaporn Suprasert*

Chiang Mai University, Chiang Mai, Thailand (psuprase@gmail.com)

Objective: To investigate the prevalence of malnourishment and predictive factors for nutritional status of gynecologic cancer patients treated with chemotherapy.

Methods: One hundred and one gynecologic cancer patients treated with chemotherapy between April 2020 and February 2021 were interviewed for Patient Generated-Subjective Global Assessment (PG-SGA). Clinical data including body mass index (BMI), underlying disease, the diagnosis, FIGO staging, the recurrent status, the surgical intervention, the performance status, the chemotherapy detail, the total number of lymphocytes per cubic milliliter (TLC) and serum albumin were collected. Levels of TLC and serum albumin were calculated for an optimal cut-off point by using receiver operating characteristic curves. The clinicopathological variables were compared using univariate and multivariate analysis to identify the independent predictive factors for malnourishment status.

Results: The prevalence of well, moderate and severe nourishment were 73.3%, 18.8% and 7.9%, respectively. The optimal cut-off points for TLC was 1,450 cells/ μ L and for albumin was 3.9 g/dL. The univariate analysis indicated that the number of present chemotherapy <3 cycles, albumin in level <3.95 g/dL, BMI <25 kg/m², TLC <1,450 cells/ μ L, anemia and no neutropenia were significant associated factors for malnourishment. However, only serum albumin <3.95 g/dL was independent associated factor for malnourishment (adjusted odds ratios = 6.709; 95% confidence interval = 2.113–21.304).

Conclusion: Nearly one-fourth of patients with gynecologic cancer who on going treatment with chemotherapy were found malnourishment. The only independent associated factor related to the nutritional status was serum albumin <3.95 g/dL.

Poster (M07)

Quality of Life, Nursing & Supportive Care
<https://doi.org/10.3802/jgo.2021.32.S1.M07>

Validity of the Comprehensive Score for Financial Toxicity (COST) tool in gynecologic cancer patients

Takashi Shibutani,¹ Shiro Suzuki,² Kohshiro Nakao,³
Takahiro Koyanagi,⁴ Shuko Tamaki,⁵ Lida Teng,⁶ Kazunori Honda,²
Yusuke Kajimoto,^{6,*} Ataru Igarashi⁶

¹Hyogo Cancer Center, Akashi, Japan

²Aichi Cancer Center Hospital, Nagoya, Japan

³Gunma University Graduate School of Medicine, Gunma, Japan

⁴Jichi Medical University, Shimotsuke, Japan

⁵Saitama Medical University International Medical Center, Hidaka, Japan

⁶The University of Tokyo, Tokyo, Japan (ykajimoto234@gmail.com)

Objective: Costs related to cancer care negatively affect patients physically and psychologically. This is called financial toxicity (FT). To quantify FT, the “COMprehensive Score for financial Toxicity (COST)” tool has been developed in the United States. The COST tool consists of 11 items as patient-reported outcomes. Our objective is to validate the COST tool for patients with ovarian, cervical, and endometrial cancer. Additionally, we aim to assess reproducibility, which explains consistent scores over time by test-retest reliability.

Methods: In this multi-center study, patients were eligible for enrollment if they had ovarian, cervical or endometrial cancer; were receiving anti-cancer drug treatment for at least 2 months, and had out-of-pocket expenses for treatment at a hospital. Patients answered the COST tool twice and sent the responses to the data center by mail. Internal validity was examined by Cronbach's alpha at the first responses. Test-retest reliability was assessed by intra-class correlation coefficient (ICC).

Results: From April 2019 through July 2021, a total of 112 patients (ovarian cancer: 50, cervical cancer: 26, and endometrial cancer: 36) answered the initial COST questionnaire and 99 patients answered the re-test. Median age was 58 years (range: 28–78 years). Mean COST score was 19.82 (SD=7.28). Cronbach's alpha was 0.83 (95% confidence interval [CI]=0.78–0.87). ICC was 0.85 (95% CI=0.77–0.90).

Conclusion: The COST tool demonstrated good internal consistency and reproducibility among patients with gynecologic cancers. The results strengthen the evidence quality of FT researches using the COST tool in gynecologic cancers.

Poster (M08)

Gynecologic Cancer Screening

<https://doi.org/10.3802/jgo.2021.32.S1.M08>

Association of cervicitis resulting from lower genital tract infections with oncogenic human papillomavirus and Pap smear abnormalities among Indian women

Kavita V Anand,^{*} Sharmila A Pimple, Gauravi A Mishra

Department of Preventive Oncology, Centre for Cancer Epidemiology (CCE),

Tata Memorial Centre, Homi Bhabha National Institute, Mumbai, India
(drkavitaanand@gmail.com)

Objective: To investigate the relationship between cervicitis with oncogenic Human papillomavirus, Bacterial vaginosis and epithelial cell abnormalities.

Methods: Total 508 women; 254 women with established diagnosis of cervicitis (case arm) and 254 asymptomatic women with healthy cervix (control arm) attending the tertiary cancer hospital for cervical cancer screening were enrolled in the study. Gram-stained smears for diagnosing lower genital reproductive tract infections (RTIs), Hybrid Capture-2 (HC2) test to diagnose oncogenic human papillomavirus (HPV) infection and Pap smears to demonstrate epithelial cell abnormalities (EP) were performed for all women enrolled in the study.

Results: Study findings demonstrated the prevalence of bacterial vaginosis (BV), HPV infection and EP to be significantly higher among women in case arm (BV=44.09%; EP=18.47%; and HPV=14.17%) than control arm (BV=3.95%; EP=4.82%; and HPV=5.12%). Case arm demonstrated increased risk for BV (odds ratio [OR]=19; 95% confidence interval [CI]=9.63–37.49), HPV (OR=3.06; 95% CI=1.58–5.92), and EP (OR=4.47; 95% CI=2.30–8.68).

Conclusion: Current study addresses the gaps in literature about the prevalence of HPV infection among Indian women with cervicitis. Our results demonstrate cervical inflammation to be significantly associated with increased risk of harboring HPV and EP. This results enforce the need of strengthening the RTI programs at national level in an attempt to reduce the burden of cervical cancer. Emerging trends of association of BV with cervicitis also warrant the treatment of BV in addition to treatment of recognized pathogens for cervicitis i.e., Gonococcal and Chlamydia infections as a part of syndromic management for women with cervicitis.

Poster (M09)

Miscellaneous

<https://doi.org/10.3802/jgo.2021.32.S1.M09>

Comparison between International Ovarian Tumor Analysis (IOTA)—simple rules and risk malignancy indices in differentiating benign and malignant adnexal masses—prospective comparative study

Vrushti Solanki, Pratibha Singh,^{*} Shashank Shekhar, Navdeep Kaur Ghuman, Garima Yadav, Priyanka Kathuria, Meenakshi Gothwal, Charu Sharma, Binit Sureka

All India Institute of Medical Sciences, New Delhi, India
(drpratibha69@hotmail.com)

Objective: To compare the sensitivity, specificity, and the